

References

- ABLES, J.R. S.L. JONES & D.W. McCOMMAS. (Jr.) (1978). Response of selected predator species to densities of *Aphis gossypii* and *Heliothis virescens* eggs. *Environmental Entomology*. **7**: 402-404.
- ANDREWARTHER, H.G. & L.C. BIRCH. (1954). *The distribution and abundance of animals*. University of Chicago Press, Chicago.
- ANON. (1995). *entomoLOGIC 1995 Manual*. CSIRO Cotton Research Unit, Narrabri, NSW, 49 pp.
- ARDITI, R. & H.R. AKCAKAYA. (1990). Underestimation of mutual interference of predators. *Oecologia*. **83**: 358-361.
- ASANTE, S.K. (1995). Functional responses of the European earwig and two species of coccinellids to densities of *Eriosoma lanigerum* (Hausmann) (Hemiptera: Aphididae). *Journal of the Australian Entomological Society*. **34**: (2) 105-109
- ATIM, A.B. & H.M. GRAHAM. (1984). Predation of *Geocoris punctipes* by *Nabis alternatus*. *The Southwestern Entomologist*. **9**: (2) 227-231.
- AWAN, M.S. (1985). Anti-predator ploys of *Heliothis punctigera* (Lepidoptera: Noctuidae) caterpillars against the predator *Oechalia schellenbergii* (Hemiptera: Pentatomidae). *Australian Journal of Zoology*. **33**: 885-890.
- AWAN, M.S. (1990). Predation by three Hemipterans: *Tropiconabis nigrolineatus*, *Oechalia schellenbergii* and *Cermatulus nasalis*, on *Heliothis punctigera* larvae in two types of searching arenas. *Entomophaga*. **35**: (2) 203-210.
- AWAN, M.S., L.T. WILSON & M.P. HOFFMANN. (1989). Prey location by *Oechalia schellenbergii*. *Entomologia experimentalis et applicata*. **51**: 225-231.
- BAUMGAERTNER, J.U., B.D. FRAZER, N. GILBERT, B. GILL, A.P. GUTIERREZ, P.M. IVES, V. NEALIS, D.A. RAWORTH, & C.G. SUMMERS. (1981). Coccinellids (Coleoptera) and aphids (Homoptera) The overall relationship. *The Canadian Entomologist*. **113**: 975-1048.
- BECHINSKI, E.J. & L.P. PEDIGO. (1982). Evaluation of methods for sampling predatory arthropods in soybean. *Environmental Entomology*. **11**: 756-761.
- BELLOWS, T.S. Jr. & J.G. MORSE. (1993) Toxicity of insecticides used in citrus to *Aphytis melinus* DeBach (Hymenoptera: Aphelinidae) and *Rhizobius lophanthae* (Blaisd.) (Coleoptera: Coccinellidae). *The Canadian Entomologist*. **125**: 987- 994.
- BERRYMAN, A.A. (1982). Biological control, thresholds and pest outbreaks. *Environmental Entomology*. **11**: 544-549.
- BOREHAM, P.F.L. & C.E. OHIAGU. (1978). The use of serology in evaluating invertebrate prey-predator relationships: a review. *Bulletin of Entomological Research*. **68**: 171-194.
- BRAMAN, S.K. & K.V. YEARGAN. (1989). Interplant distribution of three *Nabis* species (Hemiptera: Nabidae), and impact of *N. roseipennis* on green cloverworm populations in soybean. *Environmental Entomology*. **18**: (2) 240-244.

- BRAMAN, S.K. & K.V. YEARGAN. (1990). Phenology and abundance of *Nabis americanoferus*, *N. roseipennis*, and *N. rufusculus* (Hemiptera: Nabidae) and their parasitoids in alfalfa and soybean. *Journal of Economic Entomology*. **83**: (2) 823-830.
- BROADLEY, R.H. (1983). Toxicity of insecticides to *Coccinella repanda* Thunberg and *Harmonia octomaculata* (Fabricius)(Coleoptera: Coccinellidae). *Queensland Journal of Agriculture and Animal Science*. **40**: 125-127.
- BUTLER, G.D. & C.J. MAY. (1971). Laboratory studies of the searching capacity of larvae of *Chrysopa carnea* for eggs of *Heliothis* spp. *Journal of Economic Entomology*. **64**: (6) 1459-1461.
- BYERLY, K.F., A.P. GUTIERREZ, R.E. JONES & R.F. LUCK. (1978). A comparison of sampling methods for some arthropods in cotton. *Hilgardia*. **46**: (8) 257-282.
- CALTAGIRONE, L.E. (1981). Landmark examples in classical biological control. *Annual Review of Entomology*. **26**: 213-232
- CARVER, M. (1989). *Aphids, Their natural enemies and control*, Volume C, World Crop Pests A.K.Minks & P.Harrewijn.(eds.) Elsevier Science Publishers BV, Amsterdam 141-165 pp.
- CHAPMAN, R.N. (1928). The quantitative analysis of environmental factors. *Ecology*. **9**: 11-122.
- CHESSON, J. (1984). Effect of notonectids (hemiptera: Notonectidae) on mosquitoes (Diptera: Culicidae): predation or selective oviposition? *Environmental Entomology*. **13**: 531-538.
- CHESSON, J. (1989). The effect of alternative prey on the functional response of *Notonecta hoffmani*. *Ecology*. **70**: 1227-1235.
- CHOW, T., G.E. LONG & G. TAMAKI. (1983). Effects of temperature and hunger on the functional response of *Geocoris bullatus* (Say) (Hemiptera: Lygaeidae) to *Lygus* spp. (Hemiptera: Miridae) density. *Environmental Entomology*. **12**: 1332-1338.
- COX, J. C. (1981). The effects of endosulfan (Endosan®) and fenvalerate (Sumicidin®) on insects found in cotton crops. Hons. Thesis. University of New England. Armidale, NSW. Australia.
- CSIRO. (1991). *The Insects of Australia*, Melbourne University Press, 1137 pp.
- CULLEN, J.M. (1969). The reproduction and survival of *Heliothis punctigera* Wallengren in South Australia. PhD. thesis. University of Adelaide, Adelaide, Australia.
- DALY, J.C., J.H. FISK & N.W. FORRESTER. (1988). Selective mortality in field trials between strains of *Heliothis armigera* (Lepidoptera: Noctuidae) resistant and susceptible to pyrethroids: functional dominance of resistance and age class. *Journal of Economic Entomology* **85**: 1000-1007.
- DARWIN, C. (1859). The Origin of Species. J.W.Burrow (ed) (1968) *Penguin Books*. Ringwood, Australia.
- DEMPSTER, J.P. (1983). The natural control of populations of butterflies and moths. *Biological Reviews*. **58**: 461-481
- DENT, D. (1995). Integrated pest management. Chapman and Hall, London, 356 pp.

- DIETRICK, E.J. (1959). A new method for sampling arthropods using a suction collecting machine and modified Berlese funnel separator. *Journal of Economic Entomology*. **52**: 1085-1091.
- DIETRICK, E.J. (1961). An improved backpack motor fan for suction sampling of insect populations. *Journal of Economic Entomology*. **54**: 394-395.
- DILLON, G.E. & G.P. FITT. (1995). Reassessment of sampling relationships for *Helicoverpa* spp. (Lepidoptera: Noctuidae) in Australian cotton. *Bulletin of Entomological Research*. **85**: (3) 321-329.
- DINKINS, R.L., J.R. BRAZZEL & C.A. WILSON. (1970a). Seasonal incidence of major predaceous arthropods in Mississippi cotton fields. *Journal of Economic Entomology*. **63**: (3) 814-817.
- DINKINS, R.L., J.R. BRAZZEL & C.A. WILSON. (1970b). Species and relative abundance of *Chrysopa*, *Geocoris*, and *Nabis* in Mississippi cotton fields. *Journal of Economic Entomology*. **63**: (3) 660-661.
- DUFFIELD, S.J. & N.J. AEBISCHER. (1994). The effects of spatial scale of treatment with dimethoate on invertebrate population recovery in winter wheat. *Journal of Applied Ecology*. **31**: 263-281.
- ELLINGTON, J., K. KISER, G. FERGUSON & M. CARDENAS. (1984a). A comparison of sweepnet, absolute and insectvac sampling methods in cotton ecosystems. *Journal of Economic Entomology*. **77**: (3): 599-605.
- ELLINGTON, J., K. KISER, M. CARDENAS, J. DUTTLE and Y. LOPEZ. (1984b). The insectvac: a high clearance, high-volume arthropod vacuuming platform for agricultural ecosystems. *Environmental Entomology*. **13**: 259-265.
- EVANS, M.L. (1985). Arthropod species in soybean in southeast Queensland. *Journal of the Australian Entomological Society*. **24**: 169-177.
- FITT, G.P. (1989). The ecology of *Heliothis* species in relation to agrosystems. *Annual Review of Entomology* **34**: 17-52.
- FITT, G.P. (1994). Cotton pest management: Part 3. An Australian Perspective. *Annual Review of Entomology*. **39**: 543-62.
- FORRESTER, N.W. (1987). Insecticide Resistance Strategy for *Heliothis* *Agfact* A.E.43 first edition 1987, NSW. Department of Agriculture..
- FORRESTER, N.W., M. CAHILL, L.J. BIRD and J.K. LAYLAND. (1993). Management of pyrethroid and endosulfan resistance in *Helicoverpa armigera* (Lepidoptera: Noctuidae) in Australia. *Bulletin of Entomological Research*. Supplement series No. 1.
- FORRESTER, N.W. & WILSON. (1988). Insect pests of cotton. *Agfact P5.AE.1* first edition 1988, NSW. Department of Agriculture.
- FRANZ, J.M. (1972). Introductory review of the need for evaluation studies in relation to integrated control. in *Papers prepared at the symposium during the 14th international congress of entomology at Canberra in August 1972*. R.D. Hughes (Ed.). CSIRO, Canberra. 321-323 pp.
- FRAZER, B. D. and N. GILBERT. (1976). A quantitative study of the impact of ladybirds (Coleoptera: Coccinellidae) preying on field populations of pea aphids (Homoptera: Aphididae) *Journal of the Entomological Society of British Columbia*. **73**: 33-56.

- FRAZER, B. D. and B. GILL. (1981). Hunger, movement, and predation of *Coccinella californica* on pea aphids in the laboratory and in the field. *The Canadian Entomologist*. **113**: 1025-1033.
- FUCH, T.W. & J.A. HARDING. (1976). Seasonal abundance of arthropod predators in various habits in the lower Rio Grande Valley of Texas. *Environmental Entomology*. **5**: (2) 288-290.
- GARCIA, A. D. GONZALEZ & T.F. LEIGH. (1982). Three methods for sampling arthropod numbers on Californian cotton. *Environmental Entomology*. **11**: 565-572.
- GONZALEZ, D., D.A. RAMSEY, T.F. LEIGH, B.S. EKBOM & R. VAN DEN BOSCH. (1977). A comparison of vacuum and whole-plant methods for sampling predaceous arthropods on cotton. *Environmental Entomology*. **6**: (5) 750- 760.
- GRAVENA, S. & W.L. STERLING. (1983). Natural predation on the cotton leafworm (Lepidoptera: Noctuidae). *Journal of Economic Entomology*. **76**: (4) 779-784.
- GREENSTONE, M.H. (1996). Serological analysis of arthropod predation: past, present and future. In *The Ecology of Agricultural Pests: Biochemical Approaches* (eds.) W.O.C. Symondson and J.E. Liddell), Chapman and Hall, London, 265-300 pp.
- GREENSTONE, M.H. & J.H. HUNT. (1993). Determination of prey antigen half life in *Polistes metricus* using a monoclonal antibody-based immunodot assay. *Entomologia experimentalis et applicata*. **68**: 1-7.
- GREENSTONE, M.H. & C.E. MORGAN. (1989). Predation on *Heliothis zea* (Lepidoptera: Noctuidae): An instar specific ELISA assay for stomach analysis. *Annals of the Entomological Society of America*. **82**:(1) 45-49.
- GREENSTONE, M.H. & S.C. TROWELL. (1994). Arthropod predation: A simplified immunodot format for predator gut analysis. *Annals of the Entomological Society of America*. **87**: (2) 214-217
- GREGG, P.C. (1995). Migration of cotton pests: patterns and implications for management. in 'Challenging the Future' *Proceedings of the World Cotton Research Conference 1 (February 1994)* G. A. Constable and N. W. Forrester (eds.) CSIRO, Brisbane, Australia. 423-433 pp.
- HAGEN, K.S. (1962). Biology and ecology of predaceous Coccinellidae. *Annual Review of Entomology*. **7**: 289-326.
- HAGEN, K.S., P.GREANY, E.F. SAWALL (Jr.) & R.L. TASSAN. (1976). Tryptophan in artificial honeydews as a source of an attractant for adult *Chrysopa carnea*. *Environmental Entomology*. **5**: 458-468.
- HAGEN, K.S., E.F. SAWALL (Jr.) & R.L. TASSAN. (1971). The use of food sprays to increase effectiveness of entomophagous insects. *Proceedings of the Tall Timbers Conference on Ecology Animal Control and Habitat Management*. **2**: 59-81.
- HAGLER, J.R., A.C. COHEN, D. BRADLEY-DUNLOP & F.J. ENRIQUEZ. (1992). Field evaluation of predation on *Lygus hesperus* (Hemiptera: Miridae) using a species and stage-specific monoclonal antibody. *Environmental Entomology*. **21**: (4) 896-900.
- HAGLER, J.R., S.E. NARANJO, D. BRADLEY-DUNLOP, F.J. ENRIQUEZ & T.J. HENNEBERRY. (1994). A monoclonal antibody to pink bollworm (Lepidoptera: Gelechiidae) egg antigen: A tool for predator gut analysis. *Annals of the Entomological Society of America*. **87**: (1) 85-90.

- HASSELL, M.P. (1976). *The dynamics of competition and predation*. The Camelot Press Ltd. Southampton.
- HASSELL, M.P. (1985). Insect natural enemies as regulating factors. *Journal of Animal Ecology*. **54**: 323-334.
- HASSELL, M. P. (1987). Detecting regulation in patchily distributed animal populations. *Journal of Animal Ecology*. **56**: 705-713.
- HASSELL, M.P., J.H. LAWTON & J.R. BEDDINGTON. (1977). Sigmoid functional responses by invertebrate predators and parasitoids. *Journal of Animal Ecology*. **46**: 249-262
- HAWKES, R. B. (1972). A fluorescent dye technique for marking insect eggs in predation studies. *Journal of Economic Entomology*. **65**: (5) 1477-1478.
- HICKMAN, J.M., A. MACLEOD, D. TURNER, S.D. WRATTEN & M. LONGLEY. (1992). Prospects for compatible use of biological agents and insecticides in arable crops. *Pesticide Outlook*. **3**: 11-14.
- HOGG, D.B. & E.V. NORDHEIM. (1983). Age-specific survivorship analysis of *Heliothis* spp. populations on cotton. *Research of Population Ecology*. **25**: 280-297
- HOLLING, C.S. (1959a). The components of predation as revealed by a study of small-mammal predation of the European pine sawfly. *Canadian Entomologist*. **91**: 293-320.
- HOLLING, C.S. (1959b). Some characteristics of simple types of predation and parasitism. *Canadian Entomologist*. **91**: 385-398.
- HOUCK, M.A. & R.E. STRAUSS. (1985). The comparative study of functional responses: Experimental design and statistical interpretation. *Canadian Entomologist*. **117**: 617-629.
- HUTCHISON, W.D. & H.N. PITRE. (1983). Predation of *Heliothis virescens* (Lepidoptera: Noctuidae) eggs by *Geocoris punctipes* (Hemiptera: Lygaeidae) adults on cotton. *Environmental Entomology*. **12**: 1652-1656.
- IVES, P.M. (1981). Estimation of coccinellid numbers and movement in the field. *The Canadian Entomologist*. **113**: 981-997.
- JACKSON, R.D., E.G. KING & E.F. KNIPLING. (1989). *Heliothis*: Need for international collaborative research on biological control. in *Proceedings of the workshop on the biological control of Heliothis: Increasing the effectiveness of natural enemies New Dehli., Nov. 1985.* ed. E. G. King and R. D. Jackson. 15-19 pp.
- KABISSA, J.C.B. (1995). Comparative information on the local abundance of two pests, *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) and *Aphis gossypii* (Glover) (Homoptera: Aphididae), and selected predators on cotton in eastern Tanzania. in 'Challenging the Future' *Proceedings of the World Cotton Research Conference 1 (February 1994)* G. A. Constable and N. W. Forrester (eds.) CSIRO, Brisbane, Australia. 451-455 pp.
- KELLER M. A. & W. J. LEWIS. (1989). Behaviour modifying chemicals to increase the efficacy of predators and parasitoids of *Heliothis* spp. in, *Proceedings of the workshop on the biological control of Heliothis: Increasing the effectiveness of natural enemies New Dehli., Nov. 1985.* ed. E. G. King and R. D. Jackson. 449-467 pp.

- KEMENY, D. M. (1991). *A Practical Guide to ELISA*. Pergamon Press. Potts Point. Australia.
- KIMAN, Z.B. & K.V. YEARGAN. (1985). Development and reproduction of the predator *Orius insidiosus* (Hemiptera: Anthicoridae) reared on diets of selected plant material and arthropod prey. *Annals of the Entomological Society of America*. **78**: 464-467.
- KING, E.G. (1986). Insecticide use in cotton and the value of predators and parasites for managing *Heliothis*. *Proceedings of the beltwide cotton production research conference Nov. 1985* Memphis, Tenn. 155-162 pp.
- KING, E.G. & R.J. COLEMAN. (1989). Potential for biological control of *Heliothis* species. *Annual Review of Entomology*. **34**: 53-75.
- KIRITANI, K. & J.P. DEMPSTER. (1972). Different approaches to the quantitative evaluation of natural enemies. *14th International Congress of Entomology at Canberra in August 1972*. 323-330 pp.
- KNUTSON, A.E. & F. E. GILSTRAP. (1989). Direct evaluation of natural enemies of the southwestern cornborer (Lepidoptera: Pyralidae) in Texas corn. *Environmental Entomology*. **18**: (4) 732-739.
- KYI, A., M.P. ZALUCKI & I.J. TITMARSH. (1991). An experimental study of early stage survival of *Helicoverpa armigera* (Lepidoptera: Noctuidae) on cotton. *Bulletin of Entomological Research* **81**: 263-271.
- LEGASPI, J.C. and R.J. O'NEIL. (1993). Life history of *Podisus maculiventris* given low numbers of *Epilachna varivestis* as prey. *Environmental Entomology*. **22**: (5) 1192-1200.
- LEIGH, T.F. (1985). In, *Spider mites. Their biology, natural enemies and control*. W. Helle and M.W. Sabelis (eds.) Elsevier, New York. Vol. 1 349-358 pp.
- LEIGH, T.F., D.W. GRIMES, W.L. DICKENS & C.E. JACKSON. (1974). Planting pattern, plant population, irrigation and insect interactions in cotton. *Environmental Entomology*. **3**: 492-496.
- LEIGH, T.F., D. GONZALEZ & R. VAN DEN BOSCH. (1970). A sampling method for estimating absolute insect populations on cotton. *Journal of Economic Entomology*. **63**: (5) 1704-1706
- LINCOLN, C., J.R. PHILLIPS, W.H. WHITCOMB, G.C. DOWELL, W.P. BOYER, K.O. BELL (Jr.) G.L. DEAN, E.J. MATTHEWS, J.B. GRAVES, L.D. NEWSOM, D.F. CLOWER, J.R. BRADLEY (Jr.) & J.L. BAGENT. (1967). The bollworm-tobacco budworm problem in Arkansas and Louisiana. *Arkansas Agric. Exp. Stn. Bull.* 720. 66 pp.
- LINGREN, P.D. & R.L. RIDGWAY. (1969). Toxicity of five insecticides to several insect predators. *Journal of Economic Entomology*. **60**: (6) 1639-1641.
- LINGREN, P.D., R.L. RIDGWAY, C.B. COWAN (Jr.) J.W. DAVIS & W.C. WATKINS. (1968). Biological control of the bollworm and the tobacco budworm by arthropod predators affected by insecticides. *Journal of Economic Entomology*. **61**: (6) 1521-1525.
- LIVDAHL, T.P. & A.E. STIVEN. (1983). Statistical difficulties in the analysis of predator functional response data. *Canadian Entomologist*. **115**: 1365-1370.

- LOPEZ, J.D.(Jr.), R.L. RIDGWAY & R.E. PINNEL. (1976). Comparative efficiency of four insect predators of the bollworm and tobacco budworm. *Environmental Entomology*. **5**: (6) 1160-1164.
- LOVEI, G.L., P.I. SOPP & K.D. SUNDERLAND. (1990). Digestion rates in relation to alternative feeding in three species of polyphagous predators. *Ecological Entomology* **15**: 293-300
- LUCK, R.F., B. MERLE SHEPARD & P.E. KENMORE. (1988). Experimental methods for evaluating arthropod natural enemies. *Annual Review of Entomology*. **33**: 367-391.
- LUFF, M.L. (1983). The potential of predators for pest control. *Agriculture, Ecosystems and Environment*. **10**: 159-181.
- LUTTRELL, R.G., G.P. FITT, F.S. RAMALHO & E.S. SUGONYAEV. (1994). Cotton pest management: Part 1. A worldwide perspective. *Annual Review of Entomology*. **39**: 517-526.
- MACARTHUR, R.H. & E.O. WILSON. (1967). *The theory of island biogeography*. Princeton University Press, 203 pp.
- MATTHES, D. (1970). *Malachius bipustulatus* (Malachiidae) Balz und Kopulation. *Encyclopaedia Cinematographica* E1567/1969: 1-13 pp.
- McDANIEL, S.G., L.L. KEELEY & W.L. STERLING. (1978). Radiolabeling *Heliothis virescens* eggs by ³²P injection of adult females. *Annals of the Entomological Society of America*. **71**: (3) 432-434.
- McDANIEL, S.G. & W.L. STERLING. (1979). Predator determination and efficiency on *Heliothis virescens* eggs in cotton using ³²P. *Environmental Entomology*. **8**: 1083-1087.
- McDANIEL, S.G. & W.L. STERLING. (1981). Predators of tobacco budworm in Texas cotton. *The Southwestern Entomologist*. **6**: (2) 102-108.
- McDANIEL, S.G. & W.L. STERLING (1982). Predation of *Heliothis virescens* (F.) on cotton in east Texas. *Environmental Entomology*. **11**: 60-66.
- MENDAL, Z., D. BLUMBERG & I. ISHAAYA. (1994). Effects of some insect growth regulators on natural enemies of scale insects, (Hom: Coccidea). *Entomophaga*. **39**: (2) 199-209.
- MENSAH, R. K. (1996). Suppression of *Helicoverpa* spp.(Lepidoptera: Noctuidae) oviposition by use of the natural enemy food supplement Envirofeast®. *Australian Journal of Entomology*. **35**: (4) 323-329
- MENSAH, R. and W. HARRIS. (1995). Using Envirofeast (food) spray and refugia for cotton pest control. *Australian Cotton Grower*. **16**: 28-33.
- METCALF, R.L. & W.H. LUCKMANN. (1994). Introduction to Insect Pest Management. John Wiley and Sons, New York, 650 pp.
- MICHAEL, P.J. (1989). Importation and establishment of new natural enemies of *Heliothis* spp. (Lep.: Noctuidae) in Australia. in *Proceedings of the workshop on biological control of Heliothis: Increasing the Effectiveness of Natural Enemies. 11-15th Nov 1985*. Chanakya Puri, New Delhi, India. 363-373 pp.
- MICHAEL, P.J. and W. M. WOODS. (1980). An entomological review of cotton growing in the Ord River area of Western Australia. *Western Australian Department of Agriculture Technical Bulletin*. No. 48.

- MILES, M.M. (1995). Identification, pest status, ecology and management of the green mirid, *Creontiades dilutus* (Stal) (Hemiptera: Miridae), a pest of cotton in Australia. PhD. thesis, University of Queensland, 260 pp.
- MURDOCH, W.W. (1969). Switching in general predators: Experiments on predator specificity and stability of prey populations. *Ecological Monographs*. **39**: (4) 335-354.
- MURDOCH, W.W. (1973). The functional response of predators. *Journal of Applied Ecology*. **10**: 335-342.
- MURDOCH, W.W., J. CHESSON and P.L. CHESSON. (1985). Biological control in theory and practice. *The American Naturalist*. **125**: (3) 344-366.
- MURRAY, D. A., R. LLOYD. (1997). The effects of spinosad (Tracer®) on pests and beneficials. *Australian Cotton Grower* **18**: 62-64.
- MURRAY, D. A., B. SCHOLZ, R. LLOYD and K. RYNNE. (1994). Parasitoid releases on the Darling Downs. *Proceedings of the 7th Australian Cotton Conference*. Australian Cotton Growers Research Association, Brisbane, 65-68 pp.
- NAUMANN, I.D. (1993). *CSIRO Handbook of Australian Insect Names*. CSIRO, Melbourne, 200 pp.
- NICHOLSON, A. J. (1933). The balance of animal populations. *Journal of Animal Ecology*. **2**: 132-178
- NORDLUND, D.A. & R.K. MORRISON. (1990). Handling time, prey preference, and functional response for *Chrysoperla rufilabris* in laboratory. *Entomologia experimentalis et applicata*. **57**: 237-242.
- O'NEIL, R.J. (1989). Comparison of laboratory and field measurements of the functional response of *Podisus maculiventris* (Heteroptera: Pentatomidae). *Journal of the Kansas Entomological Society*. **62**: (2) 148-155.
- O'NEIL, R.J. & J.L. STIMAC. (1988). Measurement and analysis of arthropod predation on velvetbean caterpillar, *Anticarsia gemmatilis* (Lepidoptera: Noctuidae), in soybeans. *Environmental Entomology* **17**: 821-826.
- O'NEIL, R.J. & R.N. WIEDENMANN. (1987). Adaptations of arthropod predators to agricultural systems. *Florida Entomologist* **70**: (1) 40-48
- OSMAN, A.A., M. B. ATTIAH, A. EISA, & A. EL-NABAWI. (1985). Relative toxicity of pesticides to certain predators on cotton pests. *Indian Journal of Agricultural Science*. **55**: 536-538.
- PARRA, J.R.P., R.A. ZUCCHI, NETO SILVEIRA & M.L. HADDAD. (1990). Biology and thermal requirements of *Trichogramma galloi* Zucchi and *T. distinctum* Zucchi, on two alternative hosts. In *Trichogramma and other egg parasitoids*. 3rd international Symposium. E. Wajnberg and S.B. Vinson (eds.) September 23-27 1990. Institut National De La Recherche Agronomique, Paris. 81-84 pp.
- PEACOCK, W.J. & D.J. LLEWELLYN. (1996). Biotechnology in cotton - today and tomorrow. *Proceedings of the 8th Australian Cotton Conference*. Australian Cotton Growers Research Association, Brisbane, 61-67 pp.

- POTTINGER, R.P. (1967). An ecological approach to the study of insects. *New Zealand Agricultural Science*. 8-15 pp.
- PYKE, B.A. & E.H. BROWN. (1996). *The cotton pest and beneficial guide*. Cotton Research and Development Corporation, Narrabri, NSW.
- RIDGWAY, R.L. & S.L. JONES. (1968). Field- cage releases of *Chrysopa carnea* for suppression of populations of the Bollworm and the Tobacco budworm on cotton. *Journal of Economic Entomology*. **61**: (4) 892-898.
- RIDGWAY, R.L. & S.L. JONES. (1969). Inundative releases of *Chrysopa carnea* for control of *Heliothis* on cotton. *Journal of Economic Entomology*. **62**: (1) 177-1180.
- RIDGWAY, R.L., P.D. LINGREN, C.B. COWAN, Jr & J.W. DAVIS. (1967). Populations of arthropod predators and *Heliothis* spp. after applications of systemic insecticides to cotton. *Journal of Economic Entomology*. **60**: (1) 1012-1016.
- RIDGWAY, R.L. & R.K. MORRISON. (1985). Worldwide perspective on practical utilisation of *Trichogramma* with special reference to control of *Heliothis* on cotton. *The Southwestern Entomologist Supplement No. 8*. 190-198 pp.
- ROACH, S.T. (1984). Evaluation of selected vacuum sampling techniques in measuring predaceous arthropod populations in cotton. *Journal of the Georgia Entomological Society*. **19**: (2) 183-189.
- ROOM, P.M. (1977). ³²P labelling of immature stages of *Heliothis armigera* (Hubner) and *H. punctigera* (Wallengren). (Lepidoptera: Noctuidae): Relationships of dose to radioactivity, mortality and label half life. *Journal of the Australian Entomological Society* **16**: 245-251.
- ROOM, P.M. (1979a). Parasites and predators of *Heliothis* spp. (Lepidoptera: Noctuidae) in cotton on the Namoi Valley, New South Wales. *Journal of the Australian Entomological Society*. **18**: 223-228.
- ROOM, P.M. (1979b). *Insects and spiders of Australian cotton fields*. Field guide, NSW Department of Agriculture, Narrabri, N.S.W.
- ROOM, P.M. & K.G. WARDHOUGH. (1977). Seasonal occurrence of insects other than *Heliothis* spp. feeding on cotton in the Namoi valley of New South Wales. *Journal of the Australian Entomological Society*. **16**: 165-174.
- ROUSH, R.T. (1994). Managing pests and their resistance to *Bacillus thuringiensis*: can transgenic crops be better than sprays? *Biocontrol Science and Technology*. **4**: 501-516.
- RYAN, B.F., B.L. JOPINER & T.A. RYAN. (1992). MINITAB Handbook. 2nd ed., PWS-Kent, Boston, 409 pp.
- SCHICHA, E. (1974). Functional morphology of the mouth parts, and feeding habits of *Dicranolaius bellulus* (Guérin-Meneville) (Col.: Malachiidae). *Zool.Anz.,Jena*. **192**: 22-26
- SCHOLZ, B. (1990). Can egg parasites be used for *Heliothis* management in cotton? *Proceedings of the 5th Australian Cotton Conference*. Australian Cotton Growers Research Association, Brisbane, 357-359 pp.
- SCHOLZ, B. (1992). Foraging of egg parasitoids after release against *Heliothis* in dryland cotton. *Proceedings of the 6th Australian Cotton Conference*. Australian Cotton Growers Research Association, Brisbane, 257-259 pp.

- SCHOTZKO, D.J. & L.E.O'KEEFFE. (1989). Comparison of sweep net, D-vac and absolute sampling and diel variation of sweep net sampling estimates in lentils for pea aphid (Homoptera: Aphididae), nabids (Hemiptera: Nabidae), lady Beetles (Coleoptera: Coccinellidae), and lacewings (Neuroptera: Chrysopidae). *Journal of Economic Entomology*. **82**: (2) 491-506.
- SEYMOUR, J.E. & R.E. JONES. (1991). Evaluating natural enemy impact on *Heliothis*. In Zalucki M.P. (Ed). *Heliothis research methods and prospects*. Experimental Entomology Series: Springer Verlag, Sydney, 80-89 pp.
- SIDDIQUE, A.B. & R. B. CHAPMAN. (1987). Effect of prey type and quantity on the reproduction, development, and survival of pacific damsel bug, *Nabis kinbergii* Reuter (Hemiptera: Nabidae). *New Zealand Journal of Zoology*. **14**: 343-349
- SHEPARD, M., G.R. CARNER & S.G. TURNIPSEED. (1973). A comparison of three sampling methods for arthropods in soybean. *Environmental Entomology*. **3**: (2) 227-232.
- SHEPARD, M., G.R. CARNER & S.G. TURNIPSEED. (1974). Seasonal abundance of predacious arthropods in soybean. *Environmental Entomology*. **3**: (6) 985-988.
- SMITH, J.W., E.A. STADELBACHER & C.W. GANTT. (1976). A comparison of techniques for sampling beneficial arthropod populations associated with cotton. *Environmental Entomology*. **5**: (3) 435-444.
- SOLOMAN, M. E. (1949). The natural control of animal populations. **18**: (1) 1-35.
- SOPP, P.I. & K.D. SUNDERLAND. (1989). Some factors affecting the detection period of aphid remains in predators using ELISA. *Entomologia experimentalis et applicata*. **51**: 11-20.
- SOUSA, A.A., J.R. FRAZEE, M.H.J. WEIDEN & T.D.J. D'SILVA. (1977). UC 51762, a new carbamate insecticide. *Journal of Economic Entomology*. **70**: 803-807.
- SOUTHWOOD, T.R.E. (1966) Ecological methods with particular reference to the study of insect populations. Chapman and Hall Ltd. London.
- STARK, S.B. & F. WHITFORD. (1987). Functional response of *Chrysopa carnea* [Neur.: Chrysopidae] larvae feeding on *Heliothis virescens* [Lep.: Noctuidae] eggs on cotton in field cages. *Entomophaga*. **32**: (5) 521-527.
- STEPHENS, D.W. & J.R. KREBS. (1986). in *Foraging Theory. Monographs in Behaviour and Ecology*. J.R. Krebs and T. Clutton-Brock. (eds.) Princeton University Press, New Jersey.
- STERLING, W. (1989). Estimating the abundance and impact of predators and parasites on *Heliothis* populations. in *Proceedings of the workshop on the biological control of Heliothis: Increasing the effectiveness of natural enemies New Dehli., Nov. 1985.* (eds.) E. G. King and R. D. Jackson. 37-56 pp.
- STERLING, W. (1995). Pest models for decision support-development and applications. in 'Challenging the Future' *Proceedings of the World Cotton Research Conference 1 (February 1994)* G. A. Constable and N. W. Forrester (eds.) CSIRO, Brisbane, Australia. 490-493 pp.
- STONER, A. (1970). Plant feeding by a predaceous insect, *Geocoris punctipes*. *Journal of Economic Entomology*. **63**: 1911-1915.

- STONER, A. (1972). Plant feeding by *Nabis*, a predaceous genus. *Environmental Entomology*. **65**: 557-558.
- STUART, M.K. & M.H. GREENSTONE. (1990). Beyond ELISA: A rapid, sensitive, specific immunodot assay for identification of predator stomach contents. *Annals of the Entomological Society of America*. **83**: (6) 1101-1107.
- SUGONYAEV, E.S. (1994). Cotton pest management: Part 5. A Commonwealth of Independent States perspective. *Annual Review of Entomology*. **39**: 579-592.
- SUNDERLAND, K.D. (1988). Quantitative methods for detecting invertebrate predation occurring in the field. *Annals of Applied Biology*. **112**: 201-224.
- SUNDERLAND, K.D., N.E. CROOK, D.L. STACEY & B.J. FULLER. (1987). A study of feeding by polyphagous predators on cereal aphids using ELISA and gut dissection.
- SUNDERLAND, K.D. (1996). Progress in quantifying predation using antibody techniques. In *The Ecology of Agricultural Pests: Biochemical Approaches*. W.O.C. Symondson and J.E. Liddell (eds.). Chapman and Hall, London, 419-455 pp.
- TAKABAYASHI, J. & M. DICKE. (1996). Plant-carnivore mutualism through herbivore-induced carnivore attractants. *Trends in Plant Science* **1**: (4) 109-113.
- TAMIKI, G. & R.E. WEEKS. (1972). Biology and ecology of two predators, *Geocoris pallens* Stal and *G. bullatus* (Say). *Technical Bulletin No. 1446*. United States Department of Agriculture.
- TEAKLE, R.E. (1989). Estimating the abundance and impact of pathogens on *Heliothis* (Lep: Noctuidae) populations. in, *Proceedings of the workshop on the biological control of Heliothis: Increasing the effectiveness of natural enemies New Dehli., Nov. 1985*. ed. E. G. King and R. D. Jackson. 57-68 pp.
- TEAKLE, R.E. & J.M. JENSEN. (1985). *Heliothis punctigera*. in *Handbook of Rearing Insects*. (eds.) S. Pritam and R.F. Moore. Elsevier, Amsterdam. 313-322 pp.
- THEAD, L.G., H.N. PITRE & T.F. KELLOGG. (1987). Predation on eggs and larvae of *Heliothis virescens* [Lep.:Noctuidae] by an adult predator complex in cage studies on cotton. *Entomophaga*. **32**: 197-207.
- THOMPSON, N.J. & J.A. LEE. (1980). Insect resistance in cotton: A review and prospectus for Australia. *Journal of the Institute of Agricultural Science*. 75-86 pp.
- TITMARSH, I. J. (1992). Mortality of Immature Lepidoptera: A case study with *Heliothis* species (Lepidoptera: Noctuidae) in agricultural crops on the Darling Downs. PhD. Thesis. University of Queensland. Australia.
- TREACY, M.F., J.H. BENEDICT, J.D. LOPEZ and R.K. MORRISON (1987). Functional response of a predator (Neuroptera: Chrysopidae) to bollworm (Lepidoptera: Noctuidae) eggs on smoothleaf, hirsute and pilose cottons. *Journal of Economic Entomology*. **80**: 376-379.
- TREXLER, J.C., C.E. McCULLOCK and J. TRAVIS. (1988). How can the functional response best be determined? *Oecologia* **76**: 206-214.

- TROWELL, S., K. GARSIA, J. SKERRITT, A. HILL, N. FORRESTER and L. BIRD. (1994). The LepTon Test Kit (Pat. Pending) - The Heliothis ID project comes to fruition. Proceedings of the Seventh Australian Cotton Conference, Broadbeach, August 1994. Australian Cotton Growers Research Association, Brisbane, 17-20 pp.
- VAISSAYRE, M. (1995). Ecological attributes of major cotton pests: implications for management. In Constable, G.A. and Forrester, N.W. (eds.) *Challenging the future*. Proceedings of the World Cotton Research Conference I, CSIRO Australia, 569-578 pp.
- VAN DEN BERG, H. & J.W. COCK (1993). Exclusion cage studies on the impact of predation on *Helicoverpa armigera* in Cotton. *Biocontrol Science and Technology* **3**: 491-497.
- VAN DEN BERG, H. & J.W. COCK (1995). Natural control of *Helicoverpa armigera* in cotton: *Biocontrol Science and Technology* **5**: 453-463.
- VAN DEN BOSCH, R., T.F. LEIGH, D. GONZALEZ & R.E. STINNER. (1969). Cage studies on predators of the bollworm in cotton. *Journal of Economic Entomology*. **62**: (6) 1486-1489.
- VAN DRIESCH, R.G. and T.S. BELLOWS. (1996). *Biological Control*. Chapman and Hall. Melbourne. 302 pp.
- VARLEY, G.C., G.R. GRADWELL & M.P. HASSELL. (1973). *Insect Population Ecology: An Analytical Approach*. Blackwell Scientific Publications Oxford.
- VET, L.E.M. and M. DICKE. (1992). Ecology of infochemical use by natural enemies in a tritrophic context. *Annual Review of Entomology* **37**: 141-172.
- WARDHAUGH, K.G., P.M. ROOM & L.R. GREENUP. (1980). The incidence of *Heliothis armigera* (Hubner) and *H. Punctigera* (Wallingren) (Lepidoptera: Noctuidae) on cotton and other host-plants in the Namoi Valley of New South Wales. *Bulletin of Entomological Research* **70**: 113-131.
- WATKINSON, I.A. (1995). Novel biological and conventional insecticides - what's new? In Constable, G.A. and Forrester, N.W. (eds.) *Challenging the future*. Proceedings of the World Cotton Research Conference I, CSIRO Australia 569-578 pp.
- WHITCOMB, W.H. and K. BELL. (1964). Predaceous insects, spiders and mites of Arkansas cotton fields. *Arkansas Experimental Station Bulletin*. **690**: 6-84.
- WIEDENMANN, R.N. & R.J. O'NEIL. (1992). Searching strategy of the predator *Podisus maculiventris* (Say) (Heteroptera: Pentatomidae). *Environmental Entomology* **21**: (1) 1-9.
- WIEDENMANN, R.N. & J.W. SMITH Jr. (1993). Functional Response of the parasite *Cotesia flavipes* (Hymenoptera: Braconidae) at Low densities of the Host *Diatraea saccharalis* (Lepidoptera: Pyralidae). *Environmental Entomology*. **22**: (4) 849-858.
- WILLIAM, F.M. & S.A. JULIANO. (1985). Further difficulties in the analysis of functional-response experiments and a resolution. *Canadian Entomologist*. **117**: 631-640.
- WILSON, L. J. (1993). Impact of early season sprays on mites. *The Australian Cotton Grower*. **14**: (6) 26-28.
- WILSON, L. J. (1994). Resistance of Okra-leaf cotton genotypes to two spotted spider mites (Accri: Tetranychidae) *Journal of Economic Entomology*. **87**: 1726-1735.

WILSON, L. T. (1983). Abundance and mortality of overwintering *Heliothis* spp. *Journal of the Australian Entomological Society* **22**: 191-199.

WILSON, L.T. & A.P. GUTIERREZ. (1980). Within plant distribution of predators on cotton: comments on sampling efficiencies. *Hilgardia*. **48**: (2) 3-36.

WILSON, L. J. & R. MORTON. (1993). Seasonal abundance and distribution of *Tetranychus urticae* (Acari: Tetranychidae), the two spotted spider mite, on cotton in Australia and implications for management. *Bulletin of Entomological Research*. **83**: (2) 291-303.

WILSON, L.T. & P.M. ROOM. (1982). The relative efficiency and reliability of three methods for sampling arthropods in Australian cotton fields. *Journal of the Entomological Society of Australia*. **21**: 175-181.

WILSON, L.T. & G.K. WAITE. (1982). Feeding patterns of *Heliothis* on Cotton. *Environmental Entomology*. **11**: 297-300.

ZALUCKI, M.P., G. DALGLISH, S. FIREMPONG & P. TWINE. (1986). The biology and ecology of *Heliothis armigera* (Hubner) and *H. punctigera* Wallengren (Lepidoptera: Noctuidae) in Australia: What do we know? *Australian Journal of Zoology*. **34**: 779-824.
