

**A COMPARISON OF BUNK MANAGEMENT STRATEGIES
AND THEIR EFFECT ON
CATTLE PERFORMANCE AND HEALTH**

A thesis submitted as requirement for the
degree of Master of Rural Science

by

Robert John Lawrence

Faculty of the Sciences
University of New England
Armidale, NSW

December 1998

ACKNOWLEDGMENTS

This project would not have been possible without the support and approval of the Senior Japanese Management of Whyalla Feedlot, Mac Kimura and Norio Itazaki, domo arigato. I would like to thank my supervisors, Professor Jim Rowe and in particular Dr Simon Bird for his continual support and encouragement. Special thanks to Dr John Doyle for his constant enthusiasm and encouragement. Thanks also to all Whyalla staff members that were involved in the experimental stage, in particular Trevor Wells, Peter Burns, Jody Hodges and all members of the Whyalla feed team.

TABLE OF CONTENTS

	Page
PREFACE	i
ACKNOWLEDGMENTS	ii
TABLE OF CONTENTS	iii
LIST OF TABLES	vii
LIST OF FIGURES	ix
1.0 GENERAL INTRODUCTION	1
2.0 REVIEW OF LITERATURE	5
2.1 Scope of Review	5
2.2 DIGESTION IN THE RUMINANT	6
2.2.1 The Digestive Tract	6
2.2.2 Digestion of Forages	10
2.2.3 Digestion of Cereal Grains	11
2.2.3.1 Changes in the Rumen Environment with Grain Feeding	15
2.2.3.2 Acidosis and Physiological Dysfunction	18
2.2.4 Grain Feeding and Production	23
2.3 ENVIRONMENTAL EFFECTS ON ANIMAL PRODUCTION	24
2.3.1 Environmental Adaptation Mechanisms	24
2.3.2 Changes in Feeding Behaviour in Response to Heat Challenge	29
2.3.3 Heat Stress and Performance	31
2.3.3.1 Feed intake	31
2.3.3.2 Water consumption and electrolyte balance	32
2.3.3.3 Digestion and absorption	33
2.3.3.4 Energy availability and utilisation	33
2.3.4 Heat Stress and Animal Health	34

2.3.5 Management Strategies that Reduce the Effect of Heat	
Stress on Animal Performance and Health	35
2.3.5.1 Structural and physical	35
2.3.5.2 Genetic selection and environmental adaptation	35
2.3.5.3 Nutrition	36
2.3.5.4 Bunk management	36
2.4 ANIMAL BEHAVIOUR	38
2.4.1 Social Behaviour	38
2.4.1.1 Social order	38
2.4.1.2. Establishing social order	40
2.4.1.3 Maintaining social order	41
2.4.1.4 Disruption to social order	41
2.4.2 Buller Syndrome	43
2.4.3 Feeding Behaviour	45
2.4.3.1 Feed intake	45
2.4.3.2 Feeding behaviour and social interaction	46
2.4.3.3 Feeding behaviour and environmental interaction	48
2.4.4 Bunk Management, Animal Behaviour and Implications to Animal Performance	49
2.5 FEEDING MANAGEMENT OF GRAIN-FED CATTLE	50
2.5.1 The Allocator	50
2.5.2 Natural Feeding Behaviour and Bunk Management	53
2.5.3 Bunk Management Strategies	55
2.5.4 Bunk Management and Production	63
2.5.5 Conclusion	65
2.6 SUMMARY	66
3.0 ABSTRACT	67

4.0 A COMPARATIVE STUDY OF BUNK MANAGEMENT STRATEGIES	69
4.1 INTRODUCTION	69
4.2 METHODS AND MATERIALS	71
4.2.1 Experimental Design	78
4.3 RESULTS	82
4.3.1 Pen Results	82
4.3.1.1 Bunk management	82
4.3.1.2 Cattle performance	84
4.3.1.3 Carcass traits	84
4.3.1.4 Health	85
4.3.2 Feed Intake Variation	87
4.3.3 Individual Animal Results	90
4.3.4 Feeding Activity	94
4.4 DISCUSSION	97
4.4.1 Bunk Management and Feeding Costs	97
4.4.1.1 Feed wastage and labour costs	97
4.4.1.2 Machinery costs	97
4.4.1.3 Comparison of ad lib and clean bunk costs	98
4.4.2 Bunk Management and Health	99
4.4.3 Bunk Management and Carcass Composition	102
4.4.4 Bunk Management and Environment	103
4.4.5 Bunk Management and Feeding Activity	105
4.4.6 Bunk Management and Performance	109
4.4.7 Bunk Management Strategies	109

5.0 CONCLUSION	111
APPENDICES	112
APPENDIX 1 Environmental Effect on Feed Intake	114
APPENDIX 2 Feeding Activity	126
APPENDIX 3 Bunk Management Trial Period Environmental Record	142
REFERENCES	148

LIST OF TABLES

Table	Page
1.1 Grain fed beef market destinations.	1
1.2 AUS-MEAT carcass specifications and example of feedlot live animal specifications.	2
1.3 Price grid standard for short fed market.	2
2.1 Feed type and effect on saliva production rates.	6
2.2 Seasonal effects on performance of steers in a feedlot.	34
2.3 Scale used to define aggression score.	50
2.4 Definition of bunk conditions used to define bunk score.	50
2.5 Feed challenge guide, where the feed challenge (kg DM/hd/d) is the amount by which the feed offered can be increased on any one day.	51
2.6 Performance of restricted fed heifer calves compared to <i>ad lib</i> feeding.	56
2.7 Performance of restricted fed steers compared to <i>ad lib</i> feeding.	56
2.8 Performance of programme-fed steers compared to <i>ad lib</i> feeding.	57
2.9 Performance of plateau fed steer calves compared to <i>ad lib</i> feeding.	59
4.1 Nutritional analysis (theoretical values) of the ration fed during trial feeding period.	72
4.2 Average feeding times of replicates from each treatment.	73
4.3 Faecal assessment.	75
4.4 Japanese grain fed carcass specifications.	75
4.5 Summary of the various components of each treatment.	79
4.6 Allocation of treatments to pens, pen and replicate location.	79
4.7 Operation costs including feeding, feed wastage, labour required for cleaning, delivery and milling requirements associated with bunk management and the explanation of how each cost is determined.	81

4.8	Total feed wastage and costs associated with total feed offered, cleaning frequency, labour required for cleaning, delivery and milling required for feed wastage associated with bunk management strategies.	82
4.9	Treatment feed wastage (tonnes DM/pen/experimental period) associated with dry days and rain events (pen averages).	83
4.10	Bunk management strategy treatment results for feed intake, weight gain, daily gain and feed conversion efficiency over trial period.	84
4.11	Bunk management strategy treatment results for hot carcass weight, dressing percentage, bruising incidence and percentage of pen achieving Japanese grain fed carcass specifications.	85
4.12	Pen average health status of treatments in terms of total non treated, treated and returned, non returned, total pulls, bullers and deads.	85
4.13	Faecal treatment assessment scores for each faecal assessment category during trial feeding period (pen averages).	86
4.14	Number of animals diagnosed with health disorders during the trial period. Diagnosis divided into digestive, respiratory, physical, and foot/lame disorders (pen averages).	86
4.15	Intake variances for individual replicates during a 40 day (24th October 1996 to 2nd December 1996) and 36 day period (2nd January 1997 to 6th February 1997).	89
4.16	Intake variances for individual replicates during a 40 day (24th October 1996 to 2nd December 1996) and 36 day period (2nd January 1997 to 6th February 1997).	89
4.17	Number of animals and their distribution within the categories of breed, origin and age.	90
4.18	Number of animals and their distribution between scores within the categories of fat colour, marbling and meat colour.	91
4.19	Individual animal analysis of carcass measurements including carcass weight and fat depth in terms of treatment, breed, origin and age.	92

4.20 Individual animal analysis of scored measurements including marbling score, fat colour and meat colour in terms of treatment, breed, origin and age.	93
4.21 Average feeding activity of all treatments for all observations periods.	94

LIST OF FIGURES

Figure	Page
2.1 The ruminant digestive system.	7
2.2 Carbohydrate fermentation pathways of rumen microbes.	12
2.3 Ruminal fermentation products as influenced by pH	17
2.4 Sequence of events associated with induction of ruminal lactic acidosis.	17
2.5 TNZ concept.	25
2.6 Thermodynamic pathways of heat exchange between an animal and its environment.	25
2.7 Theoretical animal response to periodic climatic change.	27
2.8 Seasonal influence on diurnal eating activity of yearling steers in the feedlot	30
4.1 Average feeding activity and the relationship with radiation level.	96