

## REFERENCES

- Agee, H.R. (1969). Mating behaviour of bollworm moths. *Ann. Entomol. Soc. Am.* **62**, 1120-1122.
- Alford, A.R. & A.M. Hammond, Jr. (1982). Temperature modification of female sex pheromone release in *Trichoplusia ni* (Hübner) and *Pseudoplusia includens* (Walker) (Lepidoptera: Noctuidae). *Environ. Entomol.* **11**, 889-892.
- Armes, N.J. & R.J. Cooter (1991). Effects of age and mated status on flight potential of *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Physiol. Entomol.* **16**, 131-144.
- Babilis, N.A. & B.E. Mazomenos (1992). Mating behaviour of the corn stalk borer *Sesamia nonagrioides*. *Entomol. Exp. Appl.* **65**, 199-204.
- Baker, T.C. (1983). Variations in male oriental fruit moth courtship patterns due to male competition. *Experientia*. **39**, 112-114.
- Baker, T.C. (1989). Sex pheromone communication in the Lepidoptera: new research progress. *Experientia*. **45**, 248-262.
- Baker, T.C. & R.T. Cardé (1979). Endogenous and exogenous factors affecting periodicities of female calling and male sex pheromone response in *Grapholitha molesta* (Busck). *J. Insect Physiol.* **25**, 943-950.
- Baker, T.C. & W.L. Roelofs (1981). Initiation and termination of oriental fruit moth male response to pheromone concentrations in the field. *Environ. Entomol.* **10**, 211-218.
- Baker, T.C., W. Meyer & W.L. Roelofs (1981). Sex pheromone dosage and blend specificity of response by oriental fruit moth males. *Entomol. Exp. Appl.* **30**, 269-279.
- Barrer, P.M. & R.J. Hill (1977). Some relationships between the "calling" posture and sexual receptivity in unmated females of the moth, *Epeorus cautella*. *Physiol. Entomol.* **2**, 255-260.
- Bartell, R.J. & L.A. Lawrence (1973). Reduction in responsiveness of males of *Epiphyas postvittana* (Lepidoptera) to sex pheromone following previous brief pheromonal exposure. *J. Insect Physiol.* **19**, 845-855.
- Bartell, R.J. & W.L. Roelofs (1973). Inhibition of sexual response in males of the moth *Argyrotaenia velutinana* by brief exposure to synthetic pheromone or its geometrical isomer. *J. Insect Physiol.* **19**, 655-661.

- Cusson, M. & J.N. McNeil (1989a). Involvement of juvenile hormone in the regulation of pheromone release activities in a moth. *Science*. **243**, 210-212.
- Cusson, M. & J.N. McNeil (1989b). Ovarian development in female armyworm moths, *Pseudaletia unipuncta*: its relationship with pheromone release activities. *Can. J. Zool.* **67**, 1380-1385.
- Cusson, M., J.N. McNeil & S.S. Tobe (1990). *In vitro* biosynthesis of juvenile hormone by corpora allata of *Pseudaletia unipuncta* virgin females as a function of age, environmental conditions, calling behaviour and ovarian development. *J. Insect Physiol.* **36**, 139-146.
- Cusson, M., K.J. Yagi, S.S. Tobe & J.N. McNeil (1993). Identification of release products of corpora allata of male and female armyworm moths, *Pseudaletia unipuncta*. *J. Insect Physiol.* **39**, 775-783.
- Delisle, J. (1992). Age related changes in the calling behaviour and the attractiveness of obliquebanded leafroller virgin females, *Choristoneura rosaceana*, under different constant and fluctuating temperature conditions. *Entomol. Exp. Appl.* **63**, 55-62.
- Delisle, J. & J.N. McNeil (1986). The effect of photoperiod on the calling behaviour of virgin females of the true armyworm, *Pseudaletia unipuncta* (Haw.) (Lepidoptera: Noctuidae). *J. Insect Physiol.* **32**, 199-206.
- Delisle, J. & J.N. McNeil (1987a). The combined effect of photoperiod and temperature on the calling behaviour of the true armyworm, *Pseudaletia unipuncta*. *Physiol. Entomol.* **12**, 157-164.
- Delisle, J. & J.N. McNeil (1987b). Calling behaviour and pheromone titre of the true armyworm *Pseudaletia unipuncta* (Haw.) (Lepidoptera: Noctuidae) under different temperature and photoperiodic conditions. *J. Insect Physiol.* **33**, 315-324.
- Del Socorro, A.P. (1991). *Ecology of armyworms, particularly the common armyworm, in north-eastern New South Wales*. M.R.U. Sc. thesis, University of New England, 196 pp.
- Del Socorro, A.P. & P.C. Gregg (1991). Nocturnal behaviour of female *Mythimna convecta*. *Entomol. Exp. Appl.* **60**, 271-282.
- Devore, J. & R. Peck (1993). *Statistics: The exploration and analysis of data*. 2nd ed. Duxbury Press, California, 881 pp.
- Dingle, H. (1982). Function of migration in the seasonal synchronization of insects. *Entomol. Exp. Appl.* **31**, 36-48.
- Doane, C.C. (1968). Aspects of mating behaviour of the gypsy moth. *Ann. Entomol. Soc. Am.* **61**, 768-773.

- Drake, V.A. (1982). Insects in the sea breeze front at Canberra, a radar study. *Weather.* **37**, 134-143.
- Drake, V.A. & R.A. Farrow (1988). The influence of atmospheric structure and motions on insect migration. *Annu. Rev. Entomol.* **33**, 183-210.
- Dumont, S. & J. N. McNeil (1992). Responsiveness of *Pseudaletia unipuncta* (Lepidoptera: Noctuidae) males, maintained as adults under different temperature and photoperiodic conditions, to female sex pheromone. *J. Chem. Ecol.* **18**, 1797-1807.
- Dunkelblum, E. & M. Kehat (1992). Sexual behaviour of *Heliothis peltigera* (Lepidoptera: Noctuidae). *Bull. Entomol. Res.* **82**, 13-17.
- Dunkelblum, E., M. Kehat, M. Harel & D. Gordon (1987). Sexual behaviour and pheromone titre of the *Spodoptera littoralis* female moth. *Entomol. Exp. Appl.* **44**, 241-247.
- Ewing, A. W. (1984). Acoustic signals in insect sexual behaviour In Lewis, T.(ed.) *Insect communication*. Academic Press, London, pp. 223-240.
- Farkas, S. R. & H. H. Shorey (1974). Mechanisms of orientation to a distant pheromone source In Birch, M.C. (ed.) *Pheromones*. Elsevier North-Holland, Amsterdam, pp. 81-95.
- Farkas, S.R., H.H. Shorey & L.K. Gaston (1974). Sex pheromones of Lepidoptera. Influence of pheromone concentration and visual cues on aerial odor-trail following by males of *Pectinophora gossypiella*. *Ann. Entomol. Soc. Am.* **67**, 633-638.
- Farrow, R.A. & G. McDonald (1987). Migration strategies and outbreaks of noctuid pests in Australia. *Insect Sci. Applic.* **8**, 531-542.
- Fitzpatrick, S. M., J. N. McNeil, & S. Dumont (1988). Does male pheromone effectively inhibit competition among courting true armyworm males (Lepidoptera: Noctuidae)? *Anim. Behav.* **36**, 1831-135.
- Foster, S.P. (1993). Neural inactivation of sex pheromone production in mated lightbrown apple moths, *Epiphyas postvittana* (Walker). *J. Insect Physiol.* **39**, 267-273.
- Gadenne, C. (1993). Effects of fenoxy carb, juvenile hormone mimetic, on female sexual behaviour of the black cutworm, *Agrotis ipsilon* (Lepidoptera: Noctuidae). *J. Insect Physiol.* **39**, 25-29.
- Gatehouse, A. G. & K. P. Woodrow (1987). Simultaneous monitoring of flight and oviposition of individual velvetbean caterpillar moths (by Wales, Barfield and Leppla, 1985); a critique. *Physiol. Entomol.* **12**, 117-121.

- Gerber, G.H. & M.A. Howlader (1987). The effects of photoperiod and temperature on calling behaviour and egg development of the bertha armyworm, *Mamestra configurata* (Lepidoptera: Noctuidae). *J. Insect Physiol.* **33**, 429-436.
- Giebultowicz, J.M., R.E. Webb, A.K. Raina & R.L. Ridgway (1992). Effects of temperature and age on daily changes in pheromone titer in laboratory-reared and wild gypsy moth (Lepidoptera: Lymantriidae). *Environ. Entomol.* **21**, 822-826.
- Gould, J.L. (1980). The case for magnetic sensitivity in birds and bees (such as it is). *Amer. Sci.* **68**, 256-267.
- Grant, G.G., U.E. Brady & J.M. Brand (1972). Male armyworm scent brush secretion: identification and electroantennogram study of major components. *Ann. Entomol. Soc. Am.* **65**, 1224-1227.
- Gregg, P.C., G.P. Fitt, M. Coombs & G.S. Henderson (1993). Migrating moths collected in tower-mounted light traps in northern New South Wales, Australia: species composition and seasonal abundance. *Bull. Entomol. Res.* **83**, 563-578.
- Griffith, I.P. & A.M. Smith (1977). A convenient method for rearing lepidopteran larvae in isolation. *J. Aust. Entomol. Soc.* **16**, 366.
- Guppy, J. C. (1961). Life history and behaviour of the armyworm, *Pseudaletia unipuncta* (Haw.) (Lepidoptera: Noctuidae) in eastern Ontario. *Can. Entomol.* **93**, 1141-1153.
- Haggis, M.J. (1986). Distribution of the African armyworm, *Spodoptera exempta* (Walker) (Lepidoptera: Noctuidae), and frequency of larval outbreaks in Africa and Arabia. *Bull. Entomol. Res.* **77**, 651-660.
- Han, E. & A.G. Gatehouse (1991a). Genetics of precalling period in the oriental armyworm, *Mythimna separata* (Walker) (Lepidoptera: Noctuidae), and implications for migration. *Evolution*. **45**, 1502-1510.
- Han, E. & A.G. Gatehouse (1991b). Effect of temperature and photoperiod on the calling behaviour of a migratory insect, the oriental armyworm *Mythimna separata*. *Physiol. Entomol.* **16**, 419-427.
- Han, E. & A. G. Gatehouse (1993). Flight capacity: genetic determination and physiological constraints in a migratory moth *Mythimna separata*. *Physiol. Entomol.* **18**, 183-188.
- Harker, J.E. (1961). Diurnal rhythms. *Annu. Rev. Entomol.* **6**, 131-146.

- Haynes, K.F. & M.C. Birch (1984). The periodicity of pheromone release and male responsiveness in the artichoke plume moth, *Platytilia carduidactyla*. *Physiol. Entomol.* **9**, 287-295.
- Haynes, K.F., L.K. Gaston, M.M. Pope & Y.C. Baker (1983). Rate and periodicity of pheromone release from individual female artichoke plume moths, *Platytilia carduidactyla* (Lepidoptera:Pterophoridae). *Environ. Entomol.* **12**, 1597-1600.
- Hendrikse, A. & Vos-Bunnemeyer, E. (1987). Role of host-plant stimuli in sexual behaviour of small ermine moths (*Yponomeuta*). *Ecol. Entomol.* **12**, 363-371.
- Hill, J.K. & A.G. Gatehouse (1992). Effects of temperature and photoperiod on development and pre-reproductive period of the silver Y moth *Autographa gamma* (Lepidoptera: Noctuidae). *Bull. Entomol. Res.* **82**, 335-341.
- Hill, J.K. & A.G. Gatehouse (1993). Phenotypic plasticity and geographical variation in the pre-reproductive period of *Autographa gamma* (Lepidoptera: Noctuidae) and its implications for migration in this species. *Ecol. Entomol.* **18**, 39-46.
- Hirai, K. (1980). Male scent emitted by armyworms, *Pseudaletia unipuncta* and *P. separata* (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.* **15**, 310-315.
- Hirai, K., H.H. Shorey & L.K. Gaston (1978). Competition among courting male moths: male-to-male inhibitory pheromone. *Science*. **202**, 644-645.
- Hollander, A.L. & C. M. Yin (1982). Neurological influences on pheromone release and calling behaviour in the gypsy moths, *Lymantria dispar*. *Physiol. Entom.* **7**, 163-166.
- Howlader, M.A. & G.H. Gerber (1986a). Calling behaviour of the bertha armyworm, *Mamestra configurata* Walker (Lepidoptera: Noctuidae). *Can. Entomol.* **118**, 735-743.
- Howlader, M.A. & G.H. Gerber (1986b). Effects of age, egg development and mating on calling behaviour of the bertha armyworm, *Mamestra configurata* Walker (Lepidoptera: Noctuidae). *Can. Entomol.* **118**, 1221-1230.
- Itagaki, H. & W.E. Conner (1986). Physiological control of pheromone release behaviour in *Manduca sexta* (L.). *J. Insect Physiol.* **32**, 657-664.
- Itagaki, H. & W.E. Conner (1988). Calling behaviour of *Manduca sexta* (L.) (Lepidoptera: Sphingidae) with notes on the morphology of the female sex pheromone gland. *Ann. Entomol. Soc. Am.* **81**, 798-807.

- Jefferson, R.N., H.H. Shorey & L.K. Gaston (1966). Sex pheromones of noctuid moths. X. The morphology and histology of the female sex pheromone gland of *Trichoplusia ni* (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.* **59**, 1166-1169.
- Jefferson, R.N., H.H. Shorey & R.E. Rubin (1968). Sex pheromones of noctuid moths. XVI. The morphology of the female sex pheromone glands of eight species. *Ann. Entomol. Soc. Am.* **61**, 861-864.
- Johnson, C.G. (1969). *Migration and dispersal of insects by flight*. Methuen, London, 763 pp.
- Kaae, R.S. & H.H. Shorey (1972). Sex pheromone of noctuid moths. XXVII. Influence of wind velocity on sex pheromone releasing behaviour of *Trichoplusia ni* females. *Ann. Entomol. Soc. Am.* **65**, 436-440.
- Kanda, K.I. & A. Naito (1979). Behaviour of oriental armyworm moth, *Leucania separata* from emergence to oviposition. *Jap. J. Appl. Entomol. Zool.* **23**, 69-77.
- Kanno, H. (1979). Effects of age on calling behaviour of the rice stem borer, *Chilo suppressalis* Walk. (Lepidoptera: Pyralidae). *Bull. Entomol. Res.* **69**, 331-335.
- Kanno, H. (1981a). Mating behaviour of the rice stem borer, *Chilo suppressalis* Walker (Lepidoptera: Pyralidae) V. Critical illumination intensity for female calling and male sexual response under various temperatures. *Appl. Entomol. Zool.* **16**, 179-185.
- Kanno, H. (1981b). Mating behaviour of the rice stem borer moth, *Chilo suppressalis* Walker (Lepidoptera: Pyralidae) VI. Effects of photoperiod on the diel rhythms of mating behaviours. *Appl. Entomol. Zool.* **16**, 406-411.
- Kanno, H. & Sato, A. (1979). Mating behaviour of the rice stem borer moth, *Chilo suppressalis* Walker (Lepidoptera: Pyralidae) II. Effects of temperature and relative humidity on mating activity. *Appl. Entomol. Zool.* **14**, 419-427.
- Kawasaki, K. (1985). Factors affecting the orientation flight of *Spodoptera litura* to its sex pheromone in a wind tunnel. *Appl. Entomol. Zool.* **20**, 352-353.
- Konno, Y. (1986). Time-lag between sex pheromone content and the calling behaviour in the yellow peach moth, *Conogethes punctiferalis* (Guenée) (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.* **21**, 622-624.
- Kou, R. & Y. Chow (1987). Calling behaviour of the cotton bollworm, *Heliothis armigera* (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.* **80**, 490-493.
- Kuenen, L.P.S. & T.C. Baker (1981). Habituation versus sensory adaptation as the cause of reduced attraction following pulsed and constant sex pheromone pre-exposure in *Trichoplusia ni*. *J. Insect Physiol.* **27**, 721-726.

- Landolt, P.J. & R.R. Heath (1989). Attraction of female cabbage looper moths (Lepidoptera: Noctuidae) to male - produced sex pheromone. *Ann. Entomol. Soc. Am.* **82**, 520-525.
- Lawrence, L.A. & R.J. Bartell (1972). The effect of age on calling behaviour of virgin females of *Epiphyas postvittana* (Lepidoptera) and on their pheromone content and ovarian development. *Entomol. Exp. Appl.* **15**, 455-464.
- Lenczewski, B. & P.J. Landolt (1991). Temporal partitioning of dual sexual attraction strategies in *Trichoplusia ni* (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.* **84**, 124-130.
- Leppla, N.C. (1972). Calling behavior during pheromone release in the female pink bollworm moth. *Ann. Entomol. Soc. Am.* **65**, 281-282.
- Lingren, P.D., G.L. Greene, D.R. Davis, A.H. Baumhover & T.J. Henneberry (1977). Nocturnal behaviour of four Lepidopteran pests that attack tobacco and other crops. *Ann. Entomol. Soc. Am.* **70**, 161-167.
- Lingren, P.D., A.N. Sparks, J.R. Raulston & W.W. Wolf (1978). Applications for nocturnal studies of insects. *Bull. Entomol. Soc. Am.* **24**, 206-212.
- Linn, C.E. Jr. & L.K. Gaston (1981). Behavioral responses of male *Trichoplusia ni* in a sustained - flight tunnel to two sex pheromone components. *Environ. Entomol.* **10**, 379-385.
- Linn, C.E. Jr., M.G. Campbell & W.L. Roelofs (1986). Male moth sensitivity to multi-component pheromones: critical role of female-released blend in determining the functional role of components and active space of the pheromone. *J. Chem. Ecol.* **12**, 659-668.
- Linn, C.E. Jr., A. Hammond, J. Du & W. L. Roelofs (1988a). Specificity of male response to multi - component pheromones in noctuid moths *Trichoplusia ni* and *Pseudoplusia includens*. *J. Chem. Ecol.* **14**, 47-57.
- Linn, C.E., M.G. Campbell & W.L. Roelofs (1988b). Temperature modulation of behavioural thresholds controlling male moth sex pheromone response specificity. *Physiol. Entomol.* **13**, 59-67.
- Liu, Y. & K.F. Haynes (1993). Pheromone - mediated responses of male cabbage looper moths, *Trichoplusia ni* following various exposures to sex pheromone or (Z)-7-dodecenol. *J. Chem. Ecol.* **19**, 503-515.
- Löfstedt, C., C.E. Linn, Jr. & J. Lofqvist (1988). Behavioral responses of male turnip moths, *Agrotis segetum*, to sex pheromone in a flight tunnel and in the field. *J. Chem. Ecol.* **11**, 1209-1221.

- Marsden, J.S., G.E. Martin, D.J. Parham, T.J. Ridsdill-Smith & B.G. Johnston (1980). *Returns on Australian Agricultural Research*. CSIRO, Melbourne.
- Mbata, G.N. & S.B. Ramaswamy (1990). Rhythmicity of sex pheromone content in female *Heliothis virescens*: impact of mating. *Physiol. Entomol.* **15**, 423-432.
- McDonald, G. (1989). Colonisation of grasslands and crops of eastern Australia by the common armyworm, *Mythimna convecta* (Walker). *Proc. 5th Australasian Conf. Grassl. Invert. Ecol.* (P.P. Stahle, ed.). August, 1988, Melbourne, pp. 169-176.
- McDonald, G. (1990). A fermentation trap for selectively monitoring populations of the common armyworm moth, *Mythimna convecta* (Lepidoptera: Noctuidae). *J. Aust. Entomol. Soc.* **29**, 107-108.
- McDonald, G. (1991). The phenology, ecology and movement of the common armyworm, *Mythimna convecta* (Walker) (Lepidoptera: Noctuidae), in eastern Australia, Ph.D. Thesis, La Trobe University, 432 pp.
- McDonald, G. (1994). Insect migration in an arid continent. I. The common armyworm (*Mythimna convecta*) in eastern Australia. In Drake, V.A. & A.G. Gatehouse (eds.) *Insect migration: physical factors and physiological mechanisms*. Cambridge University Press, in press.
- McDonald, G. & P.G. Cole (1991). Factors influencing oocyte development in *Mythimna convecta* (Walker) (Lepidoptera: Noctuidae) and their possible impact on migration in eastern Australia. *Bull. Entomol. Res.* **18**, 175-184.
- McDonald, G. & A.M. Smith (1986). The incidence and distribution of the armyworms *Mythimna convecta* (Walker) and *Persectania* spp. (Lepidoptera: Notuidae) and their parasitoids in major agricultural districts of Victoria, south-eastern Australia. *Bull. Entomol. Res.* **76**, 199-210.
- McNeil, J.N. (1986). Calling behavior: can it be used to identify migratory species of moths? *Fla. Entomol.* **69**, 78-83.
- McNeil, J.N. (1987). The true armyworm *Psuedaletia unipuncta*. A victim of the pied piper or a seasonal migrant? *Insect Sci. Applic.* **8**, 591-597.
- McNeil, J.N. (1991). Behavioral ecology of pheromone-mediated communication in moths and its importance in the use of pheromone traps. *Annu. Rev. Entomol.* **36**, 407-430.
- McNeil, J.N. & J. Delisle (1989a). Are host plants important in pheromone-mediated mating systems of Lepidoptera? *Experientia*. **45**, 236-240.
- McNeil, J.N. & J. Delisle (1989b). Host plant pollen influences calling behaviour and ovarian development of the sunflower moth, *Homoeosoma electellum*. *Oecologia*. **80**, 201-205.

- Miller, J.R. & W.L. Roelofs (1977). Sex pheromone titer correlated with pheromone gland developments and age in the redbanded leafroller moth, *Argyrotaenia velutinana*. *Ann. Entomol. Soc. Am.* **70**, 136-139.
- Miller, J.R. & W.L. Roelofs (1978). Sustained - flight tunnel for measuring insect responses to windborne sex pheromones. *J. Chem. Ecol.* **4**, 142-149.
- Miyashita, K. & M. Fuwa (1972). The occurrence time, reiterative ability, and duration of mating in *Spodoptera litura* F. (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.* **7**, 171-173.
- Murlis, J., .S. Elkinton & R.T. Cardé (1992). Odor plumes and how insects use them. *Annu. Rev. Entomol.* **37**, 505-532.
- Noguchi, H. & Y. Tamaki (1985). Conspecific female-sex pheromone delays calling behaviour of *Adoxophyes* sp. and *Homona magnanima* (Lepidoptera: Tortricidae). *Jap. J. Appl. Entomol. Zool.* **29**, 113-118.
- Noldus, L.P.J.J. & R.P.J. Potting (1990). Calling behaviour of *Mamestra brassicae*: effect of age and photoperiod. *Entomol. Exp. Appl.* **56**, 23-30.
- Nordlund, D.A. & U.E. Brady (1974). The calling behaviour of female *Plodia interpunctella* (Hübner) under two light regimes. *Environ. Entomol.* **3**, 793-796.
- Ohbayashi, N., T. Yushima, H. Noguchi & Y. Tamaki (1973). Time of mating and sex pheromone production and release of *Spodoptera litura* (F.) (Lepidoptera: Noctuidae). *Kontyû*, **41**, 389-395.
- Ono, T. & M. Ito (1989). Pattern of pheromone-oriented flight in male potato tuberworm moths. *J. Chem. Ecol.* **15**, 2357-2368.
- Ono, T., R.E. Charlton & R.T. Cardé (1990). Variability in pheromone composition and periodicity of pheromone titer in potato tuberworm moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae). *J. Chem. Ecol.* **16**, 531-542.
- Ostaff, D.P., R.F. Shepherd & J.H. Borden (1974). Sex attraction and courtship behavior in *Lambdina fiscellaria lugubrosa* (Lepidoptera: Geometridae). *Can. Entomol.* **106**, 493-501.
- Palaniswamy, P. & W.D. Seabrook (1978). Behavioral responses of the female eastern spruce budworm *Choristoneura fumiferana* (Lepidoptera: Tortricidae) to the sex pheromone of her own species. *J. Chem. Ecol.* **4**, 649-655.
- Palaniswamy, P. & W.D. Seabrook (1985). The alteration of calling behaviour by female *Choristoneura fumiferana* when exposed to synthetic sex pheromone. *Entomol. Exp. Appl.* **37**, 13-16.

- Payne, C.D. (1985). *The generalised linear interactive model system release 3.77 manual*. Royal Statistical Society, NAG, Oxford, UK, 305 pp.
- Payne, T.L., M.C. Birch & C.J. Kennedy (eds.). (1986). *Mechanisms in insect olfaction*. Clarendon Press, Oxford, 364 pp.
- Pittendrigh, B.R. & K.A. Pivnick (1993). Effects of a host plant, *Brassica juncea*, on calling behaviour and egg maturation in the diamondback moth, *Plutella xylostella*. *Entomol. Exp. Appl.* **68**, 117-126.
- Preiss, R. & E. Priesner (1988). Responses of male codling moths (*Laspeyresia pomonella*) to codlemone and other alcohols in a wind tunnel. *J. Chem. Ecol.* **14**, 797-813.
- Rafaeli, A. & V. Soroker (1989). Influence of diel rhythm and brain hormone on pheromone production in two Lepidopteran species. *J. Chem. Ecol.* **15**, 447-455.
- Raina, A.K. & J.A. Klun (1984). Brain factor control of sex pheromone production in the female corn earworm moth. *Science*. **225**, 531-533.
- Raina, A.K., J.A. Klun, & E.A. Stadelbacher (1986a). Diel periodicity and effect of age and mating on female sex pheromone titer in *Heliothis zea* (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.* **79**, 128-131.
- Raina, A.K., J.A. Klun, J.D. Lopez & B.A. Leonhardt (1986b) Female sex pheromone of *Heliothis phloxiphaga* (Lepidoptera: Noctuidae): chemical identification, male behavioral response in the flight tunnel, and field tests. *Environ. Entomol.* **15**, 931-935.
- Raina, A.K., E.A. Stadelbacher & R.L. Ridgway (1989). Comparison of sex pheromone composition and pheromone-mediated male behavior of laboratory-reared and wild *Heliothis zea* (Lepidoptera: Noctuidae). *J. Chem. Ecol.* **15**, 1259-1265.
- Raina, A.K., J.C. Davis & E.A. Stadelbacher (1991). Sex pheromone production and calling in *Helicoverpa zea* (Lepidoptera: Noctuidae): effect of temperature and light. *Environ. Entomol.* **20**, 1451-1456.
- Raina, A.K., T.G. Kingan & A.K. Mattoo (1992). Chemical signals from host plant and sexual behavior in a moth. *Science*. **255**, 592-594.
- Ramaswamy, S.B. (1990). Periodicity of oviposition, feeding, and calling by mated female *Heliothis virescens* in a field cage. *J. Insect Behav.* **3**, 417-427.
- Ramaswamy, S.B., P.W.K. Ma & H.N. Pitre (1988). Calling rhythm and pheromone titers in *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) from Mississippi and Honduras. *J. Appl. Entomol.* **106**, 90-96.

- Raulston, J.R., J.W. Snow, H.M. Graham & P.D. Lingren (1975). Tobacco budworm: effect of prior mating and sperm content on the mating behaviour of females. *Ann. Entomol. Soc. Am.* **68**, 701-704.
- Roelofs, W.L. & R.T. Cardé (1974a). Sex pheromones in the reproductive isolation of lepidopterous species In Birch, M.C. (ed.) *Pheromones, Frontiers of Biology Vol. 32*. North-Holland Pub. Co., Amsterdam, pp 96-114.
- Roelofs, W.L. & R.T. Cardé (1974b). Responses of Lepidoptera to synthetic sex pheromone chemicals and their analogues. *Annu. Rev. Entomol.* **22**, 377-405.
- Ross, G.J.S. (1987). *MLP User Manual*. NAG, Rothamsted Experimental Station, 218 pp.
- Rothschild, G.H.L. & A.K. Minks (1974). Time of activity of male oriental fruit moths to pheromone sources in the field. *Environ. Entomol.* **3**, 1003-1007.
- Rothschild, G.H.L., A.G.L. Wilson & K.W. Malafant (1982). Preliminary studies on the female sex pheromones of *Heliothis* species and their possible use in control program in Australia. Proceedings of the International Workshop on *Heliothis* Management, Patancheru, A.P., India, November 1981, ICRISAT, pp. 319-327.
- Royer, L. & J.N. McNeil 1991). Changes in calling behaviour and mating success in the European corn borer (*Ostrinia nubilalis*), caused by relative humidity. *Entomol. Exp. Appl.* **61**, 131-138.
- Royer, L. & J. N. McNeil (1993). Effect of relative humidity conditions on responsiveness of European corn borer (*Ostrinia nubilalis*) males to female sex pheromone in a wind tunnel. *J. Chem. Ecol.* **19**, 61-69.
- Ryan, B.F., B.L. Joiner & T.A. Ryan, Jr. (1992). *MINITAB handbook*. 2nd ed., PWS-Kent, Boston, 409 pp.
- Sanders, C.J. (1969). Extrusion of the female sex pheromone gland in the eastern spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Can. Entomol.* **101**, 760-762.
- Sanders, C.J. (1987). Flight and copulation of female spruce budworm in pheromone-permeated air. *J. Chem. Ecol.* **13**, 1749-1758.
- Sanders, C.J. & G.S. Lucuik (1972). Factors affecting calling by female eastern spruce budworm, *Choristoneura fumiferana* (Lepidoptera:Tortricidae).*Can. Entomol.* **104**, 1751-1762.
- Sappington, T.W. & W.B. Showers (1992). Reproductive maturity, mating status, and long - duration flight behavior of *Agrotis ipsilon* (Lepidoptera: Noctuidae) and the conceptual misuse of the oogenesis-flight syndrome by entomologists. *Environ. Entomol.* **21**, 677-688.

- Sasaki, M. & L.M. Riddiford (1984). Regulation of reproductive behaviours and egg maturation in the tobacco hawk moth, *Manduca sexta*. *Physiol. Entomol.* **9**, 315-327.
- Schal, C. & R.T. Cardé (1986). Effects of temperature and light on calling in the tiger moth *Holomelina lamae* (Freeman) (Lepidoptera: Arctiidae). *Physiol. Entomol.* **11**, 75-87.
- Schal, C., R.E. Charlton & R.T. Cardé (1987). Temporal patterns of sex pheromone titers and release rates in *Holomelina lamae* (Lepidoptera: Arctiidae). *J. Chem. Ecol.* **13**, 1115-1129.
- Seabrook, W.D. (1978). Neurobiological contributions to understanding insect pheromone systems. *Annu. Rev. Entomol.* **23**, 471-485.
- Sharma, H.C. & J.C. Davies (1983). The oriental armyworm, *Mythimna separata* (Walk.), distribution, biology and control: A literature review. *Tropical Development and Research Institute, Miscellaneous Report No. 59*. T.D.R.I. London.
- Schmidt, J.O. & W.D. Seabrook (1981). Tethered moths as a measure of mating disruption in sex pheromone atmospheric permeation studies. *J. Georgia Entomol. Soc.* **16**, 60-64.
- Shorey, H.H. (1964). Sex pheromones of noctuid moths II. Mating behaviour of *Trichoplusia ni* (Lepidoptera: Noctuidae) with special reference to the role of the sex pheromone. *Ann. Entomol. Soc. Am.* **57**, 371-377.
- Shorey, H.H. (1974). Environmental and physiological control of insect sex pheromone behavior In Birch, M. C. (ed.) *Pheromones*. American Elsevier, New York, pp. 62-80.
- Shorey, H.H. (1976). *Animal communication by pheromones*. Academic Press, New York, 167 pp.
- Shorey, H.H., S.U. McFarland & L.K. Gaston (1968). Sex pheromones of noctuid moths. XXIII. Changes in pheromone quantity as related to reproductive age and mating history, in females of seven species of Noctuidae (Lepidoptera). *Ann. Entomol. Soc. Am.* **61**, 372-376.
- Simmons, A.M. & O.G. Marti, Jr. (1992). Mating by the fall armyworm (Lepidoptera: Noctuidae): frequency, duration, and effect of temperature. *Environ. Entomol.* **21**, 371-375.
- Sinchaisri, N. & K. Sogawa (1969). The influence of temperature on the development of the armyworm, *Leucania separata* Walker (Lepidoptera: Noctuidae). *Jap. J. Appl. Entomol. Zool.* **4**, 102-103.

- Smith, A.M. (1984). Larval instar determination and temperature-development studies of immature stages of the common armyworm, *Mythimna convecta* (Walker) (Lepidoptera: Noctuidae). *J. Aust. Entomol. Soc.* **23**, 91-97.
- Smith, A.M. (1986). Fecundity and survival of the common armyworm, *Mythimna convecta*: effects of temperature and larval nutrition. *Entomol. Exp. Appl.* **42**, 31-37.
- Snir, R., E. Dunkelblum, S. Gothilf & I. Harpaz (1986). Sexual behaviour and pheromone titre in the tomato looper, *Plusia chalcites* (Esp.) (Lepidoptera: Noctuidae). *J. Insect Physiol.* **32**, 735-739.
- Sohal, R. S. (1985). Aging in insects In Kerkut, G.A. & L.I. Gilbert (eds.) *Comprehensive insect physiology, biochemistry and pharmacology, Vol. 10, Biochemistry*. Pergamon Press, Oxford, pp. 596-631.
- Sower, L.L., H.H. Shorey & L.K. Gaston (1970). Sex pheromones of noctuid moths. XXI. Light:dark cycle regulation and light inhibition of sex pheromone release by females of *Trichoplusia ni*. *Ann. Entomol. Soc. Am.* **63**, 1090-1092.
- Sower, L.L., H.H. Shorey & L.K. Gaston (1971). Sex pheromones of noctuid moths. XXV. Effects of temperature and photoperiod on circadian rhythms of sex pheromone release by females of *Trichoplusia ni*. *Ann. Entomol. Soc. Am.* **64**, 488-492.
- Steel, R.G.D. & J.H. Torrie (1981). *Principles and procedures of statistics, a biometrical approach*. 2nd ed., McGraw-Hill, 633 pp.
- Strickland, G. R. & G. D. Rimes (1976). New control for Esperance armyworms. *J. Agric. West. Aust.* **17**, 14-15.
- Swaby, J.A., G.E. Daterman & L.L. Sower (1987). Mating behaviour of Douglas-fir tussock moth, *Orgyia pseudotsugata* (Lepidoptera: Lymantriidae) with special reference to effects of female age. *Ann. Entomol. Soc. Am.* **80**, 47-50.
- Swier, S.R., R.W. Rings & G.J. Musick (1976). Reproductive behaviour of the black cutworm, *Agrotis ipsilon*. *Ann. Entomol. Soc. Am.* **69**, 546-550.
- Swier, S.R., R.W. Rings & G.J. Musick (1977). Age-related calling behaviour of the black cutworm, *Agrotis ipsilon*. *Ann. Entomol. Soc. Am.* **70**, 919-924.
- Tauber, M.J., C.A. Tauber & S. Masaki (1986). *Seasonal adaptations of insects*. Oxford University Press, 411 pp.
- Teal, P.E.A., J.R. McLaughlin & J.H. Tunlinson (1981). Analysis of the reproductive behaviour of *Heliothis virescens* (F.) under laboratory conditions. *Ann. Entomol. Soc. Am.* **74**, 324-330.

- Traynier, R.M.M. (1968). Sex attraction in the Mediterranean flour moth, *Anagasta kühniella*, location of the female by the male. *Can. Entomol.* **100**, 5-10.
- Traynier, R. M. M. (1970). Sexual behaviour of the Mediterranean flour moth, *Anagasta kühniella*: some influences of age, photoperiod and light intensity. *Can. Entomol.*, **102**, 534-540.
- Turgeon, J. & J.N. McNeil (1982). Calling behaviour of the armyworm, *Pseudaletia unipuncta*. *Entomol. Exp. Appl.* **31**, 402-408.
- Turgeon, J. & J.N. McNeil (1983). Modifications in the calling behaviour of *Pseudaletia unipuncta* (Lepidoptera: Noctuidae), induced by temperature conditions during pupal and adult development. *Can. Entomol.* **115**, 1015-1022.
- Turgeon, J.J., J.N. McNeil & W.L. Roelofs (1983). Responsiveness of *Pseudaletia unipuncta* males to the female sex pheromone. *Physiol. Entomol.* **8**, 339-344.
- Vetter, R.S. & T.C. Baker (1983). Behavioral responses of male *Heliothis virescens* in a sustained flight tunnel to combinations of seven compounds identified from female sex pheromone glands. *J. Chem. Ecol.* **9**, 747-759.
- Wall, C. & J. N. Perry (1987). Range of action of moth sex-attractant sources. *Entomol. Exp. Appl.* **44**, 5-14.
- Webster, R.P. (1988). Modulation of the expression of calling by temperature in the omnivorous leafroller moth, *Platynota stultana* (Lepidoptera: Tortricidae) and other moths: a hypothesis. *Ann. Entomol. Soc. Am.* **81**, 138-151.
- Webster, R.P. & R. T. Cardé (1982). Relationships among pheromone titre, calling and age in the omnivorous leafroller moth (*Platynota stultana*). *J. Insect Physiol.* **28**, 925-933.
- Webster, R.P. & R.T. Cardé (1984). The effects of mating, exogenous juvenile hormone and a juvenile hormone analogue on pheromone titre, calling and oviposition in the omnivorous leafroller moth (*Platynota stultana*). *J. Insect Physiol.* **30**, 113-118.
- Webster, R.P. & W.E. Conner (1986). Effects of temperature, photoperiod, and light intensity on the calling rhythm in arctiid moths. *Entomol. Exp. Appl.* **40**, 239-245.
- West, R.J., P.E.A. Teal, J.E. Laing & G.M. Grant (1984). Calling behaviour of the potato stem borer, *Hydraecia micacea* Esper (Lepidoptera: Noctuidae) in the laboratory and the field. *Environ. Entomol.* **13**, 1399-1404.
- Wilson, K. & A.G. Gatehouse (1993). Seasonal and geographical variation in the migratory potential of outbreak populations of the African armyworm moth, *Spodoptera exempta*. *J. Anim. Ecol.* **62**, 169-181.

Woods, W., P.J. Lawrence & P. Booth (1980). Armyworm-a damaging pest of coarse grains on the south coast. *J. Agric. West. Aust.* **21**, 22-25.

Yushima, T., H. Noguchi, Y. Tamaki, N. Fukazawa & T. Sugino (1973). Mating and sex pheromone of *Spodoptera litura* F. (Lepidoptera: Noctuidae): an introductory report. *Appl. Entomol. Zool.* **8**, 18-26.

## APPENDIX

Reprint of "Del Socorro, A.P. & P.C. Gregg (1991). Nocturnal behaviour of female *Mythimna convecta*. *Entomologia Experimentalis et Applicata*. **60**, 271-282."