

## REFERENCES

- Anderson, D. J. (1982). The home range: a new nonparametric estimation technique. *Ecology*, 63: 103-12.
- Anon. (1977). Individual/simultaneous determination of nitrogen and/or phosphorous in BD acid digests. Technicon Autoanalyser Manual, Technicon Industrial Systems, N. Y.
- Anon. (1990). Statistix Users Guide. Analytical Software, Minnesota.
- Anon. (1995). Soils report: Walcha/Nundle and Styx River Management Areas. Environmental Impact Statement supporting document No. 5. Report to State Forests of NSW, Australia. Mitchell McCotter Pty. Ltd.
- Archer, M. (1981). A review of the origins and radiations of Australian mammals. In *Ecological Biogeography of Australia*. (ed. A. Keast), 1436-1488. Junk, The Hague.
- Archer, M. (1984). The Australian marsupial radiation. In *Vertebrate Zoogeography and Evolution in Australia*. (Eds. M. Archer and G. Clayton), 633-808. Hesperian Press, Carlisle, Western Australia.
- Archer, M., Hand, S. J. and Godthelp, H. (1991). *Riversleigh*. Reed Books, Balgowlah, Australia.
- Arnould, J. P. Y. (1995). Indices of body condition and body composition in Antarctic fur seals (*Arctocephalus gazella*). *Marine Mammal Science*, 11: 301-313).

- Bakker, H. R. and Main, A. R. (1980). Condition, body composition and total body water estimation in the Quokka, *Setonix brachyurus* (Macropodidae). *Australian Journal of Zoology*, 28: 395-406.
- Barboza, P. S. (1989). The nutritional physiology of the Vombatidae. PhD. Thesis, University of New England, Armidale, Australia.
- Barboza, P. S. (1993). Digestive strategies of the wombats: feed intake, fibre digestion, and digesta passage in two grazing marsupials with hindgut fermentation. *Physiological Zoology*, 66: 983-999.
- Barboza, P. S. and Hume, I. D. (1992). Digestive tract morphology and digestion in the wombats (Marsupialia: Vombatidae). *Journal of Comparative Physiology* 162: 552-560.
- Barboza, P. S., Hume, I. D. and Nolan, J. V. (1993). Nitrogen metabolism and requirements of nitrogen and energy in the wombats (Marsupialia: Vombatidae). *Physiological Zoology*, 66: 807-828.
- Barnett, J. L. How, R. A. and Humphries, W. F. (1979). Blood parameters in natural populations of *Trichosurus* species (Marsupialia: Phalangeridae). I. Age, sex and seasonal variation in *T. caninus* and *T. vulpecula*. *Australian Journal of Zoology*, 27: 913-26.
- Batzli, G. O. (1981). Populations and energetics of small mammals in the tundra ecosystem. In *Tundra ecosystems: a comparative analysis*. (Eds L. C. Bliss, O. W. Heal and J. J. More). 377-396. Cambridge University Press, Cambridge.
- Bedell, T. E. 1973. Botanical composition of subclover-grass pastures as affected by single and dual grazing by cattle and sheep. *Agronomy Journal*, 65: 502-4.

- Bell, R. H. V. (1970). The use of the herb layer by grazing ungulates in the Serengeti. In *Animal population in relation to their food resources*. (ed A. Watson). 111-123. Blackwell Scientific, New York.
- Bell, R. H. V. (1971). A grazing system in the Serengeti. *Scientific American*, 225: 86-93.
- Berman, D. M. and Jarman, P. J. 1988. Feral horses in the Northern Territory. Volume I. Ecology of feral horses in central Australia and their interaction with cattle. Unpublished report to the Conservation Commission of the Northern Territory.
- Black, J. L. (1971). A theoretical consideration of the effect of preventing rumen fermentation on the efficiency of utilization of dietary energy and protein in lambs. *British Journal of Nutrition*, 25: 31-55.
- Blaxter, K. L. (1962). *The energy metabolism of ruminants*. Hutchinson Scientific and Technical, London
- Bradley, W. G. and Yousef, M. K. (1972). Small mammals in the desert. In *Physiological Adaptations*. (eds. M. K. Yousef, S. M. Horvath and R. W. Bullard) 127-142. Academic Press, New York.
- Brochu, L., Caron, L. and Bergeron, J. (1988). Diet quality and body condition of dispersing and resident voles (*Microtus pennsylvanicus*). *Journal of Mammalogy*, 69: 704-710.
- Brown, G. D., and Taylor, L. S. (1984). Radio-telemetry transmitters for use in studies of the thermoregulation of unrestrained common wombats, *Vombatus ursinus*. *Australian Wildlife Research*, 11: 289-98.
- Bryan, W. W. and Evans, T. R. 1973. Effects of soils, fertilizers and stocking rates on pastures and beef production on the wallum of south-eastern Queensland. I.

- Botanical composition and chemical effects on plants and soils. *Australian Journal of Experimental Agriculture and Animal Husbandry*, 13: 516-29.
- Burt, W. H. (1943). Territoriality and home range concepts as applied to mammals. *Journal of Mammalogy*, 24: 346-52.
- Catchpoole, S. (1988). The vegetation of Epping Forest National Park. Hons thesis. University of Queensland, Brisbane.
- Cid, M. S. and Brizuela, M. A. (1990). Grass blade and sheath quantification by microhistological analysis. *Journal of Wildlife Management*, 54: 349-52.
- Clarke, J. L., Jones, M. E. and Jarman, P. J. (1989). A day in the life of a Kangaroo: Activities and movements of eastern grey Kangaroos *Macropus giganteus* at Wallaby Creek. In *Kangaroos, wallabies and rat-kangaroos*. (Eds G. Grigg, P. Jarman and I. Hume), 611-618. Surrey Beatty and Sons, Sydney.
- Cochran, W. W. and Lord, R. D. (1963). A radio-tracking system for wild animals. *Journal of Wildlife Management* 27: 9-24.
- Colwell, R. K. and Futuyma, D. J. (1971). On the measurement of niche breadth and overlap. *Ecology*, 52: 567-76.
- Cooke, B. D. (1982). Reduction of food intake and other physiological responses to a restriction of drinking water in captive wild rabbits, *Oryctolagus cuniculus*. *Wildlife Research*, 9: 247-252.
- Cooke, B. D. (1998). Did introduced European Rabbits *Oryctolagus cuniculus* (L.) displace common wombats *Vombatus ursinus* (Shaw) from part of the range in South Australia? In *Wombats*. (Eds. R. Wells and P. Pridmore), 262-70. Surrey Beatty and Sons, Chipping Norton.

- Coulson, G., Norbury, G. and Walters, B. 1989. Forage biomass and kangaroo populations (Marsupialia: Macropodidae) in summer and autumn at Hattah-Kulkyne National Park, Victoria. *Australian Mammalogy*, 13: 219-221.
- Crossman, D. G. (1988). Population and diet studies of the northern hairy-nosed wombat (*Lasiorhinus kreftii* (Owen)). Unpublished report to the Queensland National Parks and Wildlife Service. Queensland.
- Darlington, J. M. and Hershberger, T. V. (1968). Effect of forage maturity on digestibility, intake and nutritive value of alfalfa, timothy and orchardgrass by equine. *Journal of Animal Science*, 27: 1572-1576.
- Dawson, T. J., Denny, M. J. S., Russell, E. M. and Ellis, B. (1975). Water usage and diet preferences of free-ranging kangaroos, sheep and feral goats in the Australian arid zone during summer. *Journal of Zoology*, 177: 1-23.
- Dawson, T. J. and Ellis, B. A. (1979). Comparison of the diets of yellow-footed rock wallabies and sympatric herbivores in western New South Wales. *Australian Wildlife Research*, 6: 245-54.
- Degen, A. A. and Kam, M. (1995). Scaling of field metabolic rate to basal metabolic rate ratio in homeotherms. *Ecoscience*, 2: 48-54.
- Demment, M. W. and Van Soest, P. J. (1985). A Nutritional explanation for body-size patterns of ruminant and nonruminant herbivores. *American Naturalist*, 125:641-672.
- Dixon, K. R. and Chapman, J. A. (1980). Harmonic mean measure of animal activity areas. *Ecology*, 61: 1040-44.
- Duncan, P., Foose, T. J., Gordon, I. J., Gakahu, C. G. and Lloyd, M. (1990). Comparative nutrient extraction from forages by grazing bovids and equids: a test of the

- nutritional model of equid/bovid competition and coexistence. *Oecologia*, 84: 411-418.
- Dunnet, G. M., Harvie, A. E. and Smit, T. J. (1973). Estimating the proportions of various leaves in the diet of the opossum, *Trichosurus vulpecula*, Kerr, by faecal analysis. *Journal of Applied Ecology*, 10: 737-745.
- Evans, M. C. (1996). Home ranges and movement schedules of sympatric bridled nailtail and black-striped wallabies. *Wildlife Research*, 23: 547-56.
- Evans, M. C. (1997). A collar design for wombats. *Australian Mammalogy*, 20: 119-120.
- Evans, M. C. and Green, S. A. (1997). A simple blow-dart for use on trapped animals. *Australian Mammalogy*, 20: 115-117.
- Evans, M. C., Atkinson, S. and Horsup, A. (1998). Combination of tiletamine and zolazepam as a sedative and anaesthetic for wombats. *Australian Veterinary Journal*, 5: 24-25.
- Evans, M. C. and Jarman, P. J. (1999). Diets and feeding selectivity of sympatric bridled nailtail wallabies and black-striped wallabies. *Wildlife Research*, 26: 1-19.
- Fancy, S. G., Blanchard, J. M., Holleman, D. F., Kokjer, K. J. and White, R. G. (1986). Validation of the doubly-labelled water method using a ruminant. *American Journal of Physiology*, 251: 143-149.
- Feinsinger, P., Spears, E. E. and Poole, R. W. (1981). A simple measure of niche breadth. *Ecology*, 62: 27-32.
- Foley, W. J. and Hume, I. D. (1987). Digestion and metabolism of high-tannin *Eucalyptus* foliage by the brushtail possum (*Trichosurus vulpecula*) (Marsupialia: Phalangeridae). *Journal of Comparative Physiology*, 157B: 67-76.

Foley, W. J. and Cork, S. J. (1992). Use of fiberous diets by small herbivores: how far can the rules be bent? *Trends in Evolution and Ecology*, 7 (5): 159-162.

Feeose, T. J. (1982). Trophic strategies of ruminant vs non-ruminant ungulates. PhD thesis, University of Chicago.

Freudenberger, D. O. and Hume, I. D. (1992). Ingestive and digestive responses to dietary fibre and nitrogen by two macropodid marsupials (*Macropus robustus erubescens* and *M. r. robustus*) and a ruminant (*Capra hircus*). *Australian Journal of Zoology*, 40:191-194.

Friedel, M. H., Chewings, V. H. and Bastin, G. N. (1988). The use of comparative yield and dry-weight-rank techniques for monitoring arid rangeland. *Journal of Range Management*, 41: 430-35.

Gales, R. P. (1989). Validation of the use of tritiated water, doubly-labelled water, and  $^{22}\text{Na}$  for estimating food, energy and water intake in little penguins, *Eudyptula minor*. *Physiological Zoology*. 62: 147-169.

Garrot, R. A., White, G. C., Bartmann, R. M. and Weybright, D. L. (1986). Reflected signal bias in biotelemetry triangulation systems. *Journal of Wildlife Management*, 50: 747-52.

Gaughwin, M. D., Judson, G. D., Macfarlane, W. V. and Siebert, B. D. (1984). Effect of drought on the health of wild hairy-nosed wombats *Lasiorhinus latifrons*. *Australian Wildlife Research*, 11: 455-63.

Gipps, J. M. and Sanson, G. D. (1984). Mastication and digestion in *Pseudochirus*. In *Possoms and Gliders*. (Eds. A. P. Smith and I. D. Hume.), 234-46. Australian Mammal Society, Sydney.

Goering, H. K. and Van Soest, P. J. (1970). Forage fibre analyses. Agriculture Handbook No 379. U. S. Department of Agriculture.

Golly, F. B. (1960). Energy dynamics of a food chain of an old-field community. *Ecological Monographs*, 30: 187-206.

Gordon, G., Riney, T., Toop, J., Lawrie, B., and Godwin M. D. (1985). Observations on the Queensland Hairy-nosed Wombat *Lasiorhinus kreftii* (Owen). *Biological Conservation*, 33:165-195.

Green, B. (1997). Field energetics and water fluxes in marsupials. In *Marsupial Biology: Recent Research, New perspectives*. (Eds N.R. Saunders and L.A. Hinds.), 143-162. University of New South Wales Press.

Green, B. and Eberhard, I. D. (1983). Water and sodium intake, and estimated food consumption, in free-living quolls, *Dasyurus viverrinus*. *Australian Journal of Zoology*, 31: 871-880.

Hammond, K. A. and Diamond, J. (1997). Maximal sustained energy budgets in humans and animals. *Nature*, 386: 457-462.

Harris, S., Cresswell, P. G. Forde, P. G., Trewhella, W. J., Woollard, T. and Wray, S. (1990). Home-range analysis using radio-tracking data - a review of problems and techniques particularly as applied to the study of mammals. *Mammal Review*, 20:97-123.

Harvey, P. H., Pagel, M. D. and Rees, J. A. (1991). Mammalian metabolism and life histories. *The American Naturalist*, 137: 556-566.

Haydock, K. P. and Shaw, H.H. (1975). The comparative yield method for estimating dry matter yield of pasture. *Australian Journal of Experimental Agriculture and Animal Husbandry*, 15: 663-70.

- Heezen, K. H. and Tester, J. R. (1967). Evaluation of radio-tracking by triangulation with special reference to deer movements. *Journal of Wildlife Management*, 31: 124-141.
- Heyward, R. (1996). Survey for the common wombat, *Vombatus ursinus*, in Butterleaf State Forest, Glen Innes Management Area. Unpublished report by State Forests of New South Wales to the Department of Urban Affairs and Planning.
- Hill, G. J. E. (1982). Seasonal movement patterns of eastern grey kangaroo in southern Queensland. *Australian Wildlife Research*, 9: 373-387.
- Hope, J. (1984). The Australian Quarternary. In *Vertebrate Zoogeography and Evolution in Australia*. (Eds. M. Archer and G. Clayton), 633-808. Hesperian Press, Carlisle, Western Australia.
- Hoppe, P. P. (1977). Rumen fermentation and body weight in African ruminants. In *Proc. 13th International Congress of Game Biologists*. (ed T. J. Peterle). 141-150. Wildlife Society, Washington, D. C.
- Horsup, A and Davidson, C. (1994). *Recovery Plan for the Northern Hairy-nosed Wombat*. Unpublished report to the Australian Nature Conservation Agency, Canberra.
- Hulbert, S. H. (1978). The measurement of niche overlap and some relatives. *Ecology*, 59: 67-77.
- Hume, I. D. (1982). *Digestive physiology and nutrition of marsupials*. Cambridge University Press, Cambridge.
- Hume, I. D. (1989). Optimal digestive strategies in mammalian herbivores. *Physiological Zoology*, 62(6): 1145-1163.

Illius, A. W. and Gordon, I. J. (1992). Modelling the nutritional ecology of the ungulate herbivores: evolution of body size and competitive interactions. *Oecologia*, 89: 428-434.

IUCN (International Union for the Conservation of Nature and natural resources). (1990). *Red List of Threatened Animals*. IUCN, Gland Switzerland and Cambridge U.K.

Ivlev, V. S. (1961). *Experimental ecology of the feeding of fishes*. Yale University Press, Connecticut, USA.

Janis, C. (1976). The evolutionary strategy of the Equidae and the origins of the rumen and caecal digestion. *Evolution*, 30: 757-774.

Janis, C. M. (1989). A climatic explanation for the patterns of evolutionary diversity in ungulate mammals. *Palaeontology*, 32: 463-81.

Janis, C. M. (1993). Tertiary mammal evolution in the context of changing climates, vegetation, and tectonic events. *Annual Review of Ecological Systematics*, 24: 467-500.

Jaremovic, R. V. and Croft, D. B. (1987). Comparison of techniques to determine eastern grey kangaroo home range. *Journal of Wildlife Management*, 51: 921-30.

Jarman, P. J. (1973). The free water intake of impala in relation to the water content of their food. *East African Agricultural and Forestry Journal*, 4: 343-51.

Jarman, P. J. (1974). The social organisation of antelope in relation to their ecology. *Behaviour*, 48: 215-266.

Jarman, P. J. (1994). The eating of seedheads by species of Macropodidae. *Australian Mammalogy*, 17:51-63.

Jarman, P. J. and Sinclair, A.R. E. (1979). Feeding strategy and the pattern of resource partitioning in ungulates. In *Serengeti. Dynamics of an ecosystem*. (eds A.R.E. Sinclair and M. Norton-Griffiths). 130-163. University of Chicago Press, Chicago.

Jarman, P. J. and Phillips, C. M. (1989). Diets in a Community of Macropod Species. In *Kangaroos, wallabies and rat-kangaroos*. (Eds G. Grigg, P. Jarman and I. Hume), 143-149. Surrey Beatty and Sons, Sydney.

Jeffery, E. (1996). Diets of macropods and wombats at Riamukka State Forest, New South Wales. Grad. Dip. Thesis. University of New England, Armidale.

Jenrich, R. I. and Turner, F. B. (1969). Measurement of non-circular home range. *Journal of Theoretical Biology*, 22: 227-37.

Johnson, C. N. (1991a). Utilization of habitat by the northern hairy-nosed wombat *Lasiorhinus krefftii*. *Journal of the Zoological Society of London*, 225: 495-507.

Johnson, C. N. (1991b). Behaviour and ecology of the northern hairy-nosed wombat *Lasiorhinus krefftii*. Unpublished report to the Queensland National Parks and Wildlife Service. Queensland.

Johnson, C. N. (1998). The evolutionary ecology of wombats. In *Wombats*. (Eds. R. Wells and P. Pridmore), 34-41. Surrey Beatty and Sons, Chipping Norton.

Johnson, C. N. and Gordon, G. (1995). Northern hairy-nosed wombat. In *The Mammals of Australia*. (Ed R. Strahan.) 200-201. Reed Books, Chatswood Australia.

Karasov, W. H. (1991). Daily energy expenditure and the cost of activity in a free-living mammal. *Oecologia*, 51:253-259.

Kenward, R. (1987). *Wildlife radio-tagging: equipment, field techniques and data analysis*. Academic Press, California.

Kinnear, J. E., Cockson, A., Christensen, P. and Main, A. R. (1979). The nutritional biology of the ruminants and ruminant-like mammals - a new approach. *Comp. Biochem. Physiol.*, 64: 357-365.

Kirsch, J. (1984). Marsupial origins: taxonomic and biological considerations. In *Vertebrate Zoogeography and Evolution in Australia*. (Eds. M. Archer and G. Clayton), 633-808. Hesperian Press, Carlisle, Western Australia.

Kleiber, M. (1975). *The fire of life: an introduction to animal energetics*. Krieger, Huntington, N.Y.

Klein, P. D., James, W. P. T., Wong, W. W. Irving, C. S., Murgatroyd, P. R., Cabrera, M., Dallosso, H. M., Klein, E. R. and Nichols, B. L. (1984). Calorimetric validation of the doubly-labelled water method for determination of energy expenditure in man. *Human Nutrition: Clinical Nutrition*, 38C: 95-106.

Koeppel, J. W., Slade, N. A. and Hoffmann, R. S. (1975). A bivariate home range model with possible application fo ethological data analysis. *Journal of Mammalogy*, 56: 81-90.

Koeppel, J. W., Slade, N. A., Harris, K. S. and Hoffmann, R. S. (1977). A three dimensional home range model. *Journal of Mammalogy*, 58: 213-20.

Koteja, P. (1991). On the relation between basal and field metabolic rates in birds and mammals. *Functional Ecology*, 5: 56-64.

Krebs, C. J. and Singleton, G. R. (1993). Indicies of condition for small mammals. *Australian Journal of Zoology*, 41: 317-323.

- Kunz, T. H. and Nagy, K. A. (1989). Methods of energy budget analysis. In *Ecological and Behavioural Methods for the study of Bats*. (ed T. H. Kunz), 277-302. Smithsonian Institute Press, Washington D. C.
- Langer, P. (1984). Anatomical and nutritional adaptations in wild herbivores. In *Herbivore nutrition in the tropics and subtropics*. (eds. F. M. C. Gilchrist and R. I. Mackie), 185-203. Science Press. Craighall, South Africa.
- Langer, P. (1988). *The mammalian herbivore stomach*. Stuttgart, New York.
- Lechowicz, M. J. (1982). The sampling characteristics of Electivity Indicies. *Oecologia*, 52: 22-30.
- Lee, J. E., White, G. C., Garrot, R. A., Bartmann, R. M. and Alldredge, A. W. (1985). Assessing accuracy of a radiotelemetry system for estimating animal locations. *Journal of Wildlife Management*, 43: 658-63.
- Lehmann, P. A Myoporum-Stipa Community Grazed by wombats - with particular reference to herbage production. MSc Thesis, University of Adelaide. Australia.
- Lenth, R. V. (1981). On finding the source of a signal. *Technometrics*, 23: 149-154.
- Levins, R. (1968). *Evolution in changing environments*. Princeton University Press, New Jersey, USA.
- Lifson, N., and McClintock, R. (1966). Theory of use of the turnover rates of body water for measuring energy and material balance. *Journal of Theoretical Biology*, 12: 46-74.

- MacDonald, D. W. (1978). Radio-tracking: some applications and limitations. In *Animal marking: Recognition marking of animals in research*. (Ed B. Stonehouse), 192-204. MacMillan, London.
- Macdonald, D. W., Ball, F. C. and Hough, N. G. (1980). The evaluation of home range size and configuration using radio tracking data. In *A handbook on biotelemetry and radio tracking*. (Eds. C. J. Amlaner and D. W. Macdonald). Pergamon Press. Oxford.
- Mallett, K. J. and Cooke B. D. (1986). *The ecology of the common wombat in South Australia*. Nature Conservation Society of South Australia, Adelaide.
- Martin, H. (1984). Australian Phytogeography. In *Vertebrate Zoogeography and Evolution in Australia*. (Eds. M. Archer and G. Clayton), 633-808. Hesperian Press, Carlisle, Western Australia.
- McBee, R. H. (1971). Significance of intestinal microflora in herbivory. *Annual Review of Ecological Systematics*, 2: 165-176.
- McIlroy, J. C. (1976). Aspects of the ecology of the common wombat, *Vombatus ursinus*. 1. Capture, handling, marking and radio-tracking techniques. *Australian Wildlife Research*, 3(2): 105-16.
- McIlroy, J. C. (1995). Common Wombat. In *The Mammals of Australia*. (Ed R. Strahan.) 204-205. Reed Books, Chatswood Australia.
- McNaughton, S. J., Tarrants, J. L., McNaughton, M. M. and Davis, R. H. (1985). Silica as a defense against herbivory and a growth promoter in African grasses. *Ecology*, 66: 528-535.

- Minson, D. J. and McLeod, M. N. (1970). The digestibility of temperate and tropical grasses. In Proceedings of the 11th Grassland Congress. (Ed M. J. T. Norman), 719-732. University of Queensland Press, St Lucia, Australia.
- Mohr, C. O. (1947). Table of equivalent populations of the North American small mammals. *American Midlands Naturalist*, 37: 223-49.
- Morris, P. A. (1988). A study of home-range and movements in the hedgehog (*Erinaceus europaeus*). *Journal of Zoology*, 214:433-49.
- Munks, S. A. (1990). Ecological energetics and reproduction in the common ringtail possum, *Pseudocheirus peregrinus* (Marsupialia: Phalangeroidea). PhD thesis. University of Tasmania.
- Murray, P. (1998). Palaeontology and palaeobiology of wombats. In *Wombats*. (Eds. R. Wells and P. Pridmore), 1-33. Surrey Beatty and Sons, Chipping Norton.
- Nagy, K. A. (1980). CO<sub>2</sub> production in animals: analysis of potential errors in the doubly labelled water method. *American Journal of Physiology*, 238: 466-73.
- Nagy, K. A. (1983). *The doubly labelled water (<sup>3</sup>HH<sup>18</sup>O)method: a guide to its use*. UCLA Publication No. 12-1417. UCLA, California.
- Nagy, K. A. (1987). Field metabolic rate and food requirement scaling in mammals and birds. *Ecological Monographs*, 52: 111-128.
- Nagy, K. A. (1989). Doubly-labelled water studies of vertebrate physiological ecology. In *Stable isotopes in ecological research*. (Eds P. W. Rundel, J. R. Ehleringer and K. A. Nagy), 270-287. Springer-Verlag, New York.
- Nagy, K. A. (1994a). Seasonal water, energy and food use by free-living, arid adapted mammals. *Australian Journal of Zoology*, 42: 55-63.

- Nagy, K. A. (1994b). Field bioenergetics of mammals: what determines field metabolic rates? *Australian Journal of Zoology*, 42: 43-53.
- Nagy, K. A. and Milton, K. (1979). Energy metabolism and food consumption by wild howler monkeys (*Alouatta palliata*). *Ecology*, 60: 475-80.
- Nagy, K. A. and Costa, D. P. (1980). Water flux in animals: analysis of potential errors in the tritiated water method. *American Journal of Physiology*, 238: 454-65.
- Nagy, K. A. and Martin, R. W. (1985). Field metabolic rate, water flux, food composition and time budget of koalas, *Phascolarctos cinereus* (Marsupialia: Phascolarctidae) in Victoria. *Australian Journal of Zoology*, 33: 655-65.
- Nagy, K. A. and Peterson, C. P. (1988). Scaling of water flux in animals. *Uni. Calif. Publ. Zool.*, 120.
- Nagy, K. A., Foley, W. J., Kaplan, I. R., Meredith, D. and Minagawa, M. (1990). Doubly-labelled water validation in the marsupial *Petauroides volans*. *Australian Journal of Zoology*, 38: 469-77.
- Nagy, K. A., Sanson, G. D. and Jacobsen, N. K. (1990). Comparative field energetics of two macropod marsupials and a ruminant. *Wildlife Research*, 17: 591-9.
- Nams, V. O. and Boutin, S. (1991). What is wrong with error polygons? *Journal of Wildlife Management*, 55: 172-76.
- Newsome, A. (1995). Red Kangaroo. In *The Mammals of Australia*. (Ed R. Strahan.) 353-355. Reed Books, Chatswood Australia.
- Norbury, G. L., Sanson, G. D. (1992). Problems with measuring diet selection of terrestrial mammalian herbivores. *Australian Journal of Ecology*, 17: 1-7.

Owen, R. (1877). Researches on the fossil remains of the extinct mammals of Australia; with a notice of the extinct marsupials of England. *J. Erxleben*, Vol 1.

Owen-Smith, R. N. (1982). Factors influencing the consumption of plant products by large herbivores. In *Ecology of tropical savannas*. (Ed B. . Huntley and B. H. Walker), 359-404. Springer-Verlag, Berlin.

Owen-Smith, R. N. (1992). *Megaherbivores. The influence of very large body size on ecology*. Cambridge University Press, Cambridge.

Pace, N and Rathburn, E. N. (1945). Studies on body composition. III. The body water and chemically combined nitrogen in relation to fat content. *Journal of Biological Chemistry*, 158: 685-691.

Parra, R. (1978). Comparison of foregut and hindgut fermentation in herbivores. In *The ecology of arboreal folivores*. (ed G. Montgomery), 205-229. Smithsonian Institute, Washington D.C.

Pearson, D. J. (1983). The diet of the rufous hare-wallaby, *Lagorchestes hirsutus*, in the Tanami Desert. B. Nat. Res. thesis, University of New England.

Peters, D. G. and Rose, R. W. (1979). The oestrus cycle and basal body temperature in the common wombat (*Vombatus ursinus*). *Journal of Reproduction and Fertility*, 57: 453-460.

Petraitis, P. S. (1979). Likelihood measures of niche breadth and overlap. *Ecology*, 60: 703-10.

Pledge, N. (1992). The Curramulka Local Fauna: a new late Tertiary fossil assemblage from York Peninsula, S. Australia. *The Beagle, Records of the Northern Territory Museum of Arts and Science*. 9(1): 115-42.

Poole (1995). Eastern Grey Kangaroo. In *The Mammals of Australia*. (Ed R. Strahan.) 335-338. Reed Books, Chatswood Australia.

Priddel, D. (1983). The movement of red and western grey kangaroos in western New South Wales. PhD. thesis, University of Sydney.

Pyke, G. H. and O'Connor, P. J. (1990). The accuracy of a radio-tracking system for monitoring honeyeater movements. *Australian Wildlife Research*, 17: 501-9.

Raymond, W. F. D., Minson, J. and Harris, C. E. (1959). Studies in the digestibility of herbage. 7. Further evidence on the effect of level of intake on the digestive efficiency of sheep. *Journal of the British Grasslands Society*, 14: 75-77.

Rees, M. C., Jones, R. M. and Roe, R. 1976. Evaluation of pasture grasses and legumes grown in mixtures in south-east Queensland. *Tropical Grasslands*, 10: 65-78.

Rishworth, C., McIlroy, J. C. and Tanton, M. T. (1995). Diet of the common wombat, *Vombatus ursinus*, in plantations of *Pinus radiata*. *Wildlife Research*, 22: 333-339.

Roajevic, I., Simpson, R. J., St.John, J. A. and Humphreys, M. O. (1994). Chemical composition and in vitro digestibility of lines of *Lolium perenne* selected for high concentrations of water-soluble carbohydrate. *Australian Journal of Agricultural Research*, 45: 901-912.

Robbins, C. T. (1983). *Wildlife feeding and nutrition*. Academic Press, London.

Robertson, G. G. and Gepp, B. (1982). Capture of kangaroos by 'stunning'. *Australian Wildlife Research*, 9: 393-6.

Rothwell, N. J. and Stock, M. J. (1979). *In vivo* determination of body composition by tritium dilution in the rat. *British Journal of Nutrition*, 4: 625-628.

- Saltz, D. and Alkon, P. U. (1985). A simple computer-aided method for estimating radio-location error. *Journal of Wildlife Management*, 49: 664-681.
- Samual, M. D. and Garton, E. O. (1985). Home range: a weighted normal estimate and tests of underlying assumptions. *Journal of Wildlife Management*, 49: 664-68.
- Sanson, G. D. (1989). Morphological adaptations of teeth to diets and feeding in the Macropodidae. In *Kangaroos, wallabies and rat-kangaroos*. (Eds G. Grigg, P. Jarman and I. Hume). 151-168. Surrey Beatty and Sons, Sydney.
- Schoener, T. W. (1974). Some methods for calculating competition coefficients from resource utilization spectra. *American Naturalist*, 108: 332-40.
- Sheng, H. P. and Huggins, R. A. (1979). A review of body composition with emphasis on total body water and fat. *American Journal of Clinical Nutrition*, 32: 630-647.
- Short, J. 1986. The effect of pasture availability on food intake, species selection and grazing behaviour of kangaroos. *Journal of Applied Ecology*, 23: 559-571.
- Sinclair, A. R. E. (1977). *The African buffalo: a study of resource limitation of populations*. University of Chicago Press, Chicago.
- Sinclair, A. R. E., Krebs, C. J., Smith, J. N. M. (1982). Diet quality and food limitation in herbivores: the case of the snowshoe hare. *Can. J. Zool.*, 60: 889-897.
- Slayter, J. and Jones, R. J. (1971). Estimation of diets selected by grazing animals from microscopic analysis of the faeces - a warning. *Journal of the Australian Institute of Agricultural Science*, 37: 238-240.

Smith, A. P., Moore, D. M. and Andrews, S. (1992). Proposed operations in the Glen Innes Forest Management Area. Impacts on Fauna. Report to the Forestry Commission of New South Wales.

Snedecor, G. W. and Cochran, W. G. (1980). *Statistical methods*. Iowa State University Press, Ames, Iowa.

Southwell, C. (1981). Sociobiology of the eastern grey kangaroo, *Macropus giganteus*. Ph.D. Thesis, University of New England, Armidale.

Southwell, C. (1987). Macropod Studies at Wallaby Creek II. Density and Distribution of Macropod Species in Relation to Environmental Variables. *Australian Wildlife Research*, 14: 15-33.

Southwood, T. R. E. (1966). *Ecological methods: with particular reference to the study of insect populations*. Chapman and Hall, London.

Springer, J. T. (1979). Some source of bias and sampling error in radio triangulation. *Journal of Wildlife Management*, 43: 926-35.

Taylor, R. J. (1993). Observations on the behaviour and ecology of the common wombat *Vombatus ursinus* in northeast Tasmania. *Australian Mammalogy*, 16: 1-7.

Thonney, M. L., Touchberry, R. W., Goodrich, R. D. and Meiske, J. C. (1976). Intraspecies relationship between fasting heat production and body weight: a reevaluation of  $W^{0.75}$ . *Journal of Animal Science*, 43: 692-704.

Tideman, C. R., Priddel, D., Nelson, J. E. and Pettigrew, J. D. (1985). Foraging behaviour of the Australian ghost bat, *Macroderma gigas* (Microchiroptera: Megadermatidae). *Australian Journal of Zoology*, 33: 705-13.

- Tixier, H., Duncan, P., Scehovic, J., Yani, A., Gleizes, M. and Lila, M. (1997). Food selection by European roe deer (*Capreolus capreolus*): effects of plant chemistry and consequences for the nutritional value of their diets. *Journal of Zoology*, 242: 229-245.
- 't Mannetje, L. and Haydock, K. P. (1963). The dry-weight-rank method for botanical analysis of pasture. *Journal of the British Grassland Society*, 18: 268-275.
- Tothill, J. C. Hargreaves, J. N. G. and Jones, R. M. 1978. BOTANAL - a comprehensive sampling and computational procedure for estimating pasture yield and composition. *CSIRO Tropical Agronomy Technical Memorandum*, 8.
- Triggs, B. (1996). *The Wombat*. Revised Edition, UNSW Press, Sydney.
- Troughton, E. (1948). *Furred animals of Australia*. 8th edn. Angus and Robertson, Sydney.
- Trumble, H. C. (1948). Rainfall, vegetation and drought frequency in South Australia. *South Australian Journal of Agriculture*, 52: 55-64.
- Tyrrell, H. F. and Moe, P. W. (1975). Effect of intake on digestive efficiency. *Journal of Dairy Science*, 58: 602-610.
- Van Soest, P. J. (1964). Symposium on nutrition and forage of pastures: new chemical procedures for evaluating forages. *Journal of Animal Science*, 123: 838-845.
- Van Soest, P. J. (1982). *Nutritional ecology of the ruminant*. O and B Books, Corvallis, Ore.
- Vanderploeg, H. A. and Scavia, D. (1979). Two Electivity Indices for feeding with special reference to zooplankton grazing. *Journal of the Fisheries Research Board of Canada*, 36: 362-5.

- Vaughn, B. E. and Boling, E. A. (1961). Rapid assay procedures for tritium-labelled water in body fluids. *Journal of Laboratory and Clinical Medicine*, 57: 159-164.
- Walker, B. H. 1976. An approach to the monitoring of changes in the composition and utilisation of woodland and savanna vegetation. *South African Journal of Wildlife Research*, 6: 1-32.
- Wells, R. T. (1973). Physiological and behavioural adaptations of the hairy-nosed wombat (*Lasiorhinus latifrons* Owen) to its arid environment. PhD Thesis, University of Adelaide, Australia.
- Wells, R. T. (1978). Thermoregulation and activity in the hairy-nosed wombat, *Lasiorhinus latifrons* (Owen) (Vombatidae). *Australian Journal of Zoology*, 26: 639-51.
- Wells, R. T. (1989). Wombats. In *Fauna of Australia. Vol 1B Mammalia*. 755-768. Australian Biological Resources Study, Canberra.
- Wells, R. T. (1995). Southern hairy-nosed wombat. In *The Mammals of Australia*. (Ed R. Strahan.) 202-203. Reed Books, Chatswood Australia.
- Westoby, M. Rost, R. R. and Weis, J. A. (1976). Problems with estimating herbivore diets by microscopically identifying plant fragments from stomachs. *Journal of Mammalogy*, 57: 167-72.
- White, G. C. (1985). Optimal locations of towers for triangulation studies using biotelemetry. *Journal of Wildlife Management*, 49: 190-96.
- White, G. C. and Garrot, R. A. (1986). Effects of biotelemetry triangulation error on detecting habitat selection. *Journal of Wildlife Management*, 50: 509-13.

White, G. C. and Garrot, R. A. (1990). *Analysis of wildlife radio-tracking data*. Academic Press, California.

Wood, R. A., Nagy, K. A., MacDonald, N. S., Wakakuwa, S. T., Beckman, R. J. and Kaaz, H. (1975). Determination of Oxygen-18 in water contained in biological samples by charged particle activation. *Analytical Chemistry*, 47: 646-650.

Woolnough, A. P. (1998). The feeding ecology of the northern hairy-nosed wombat, *Lasiorhinus krefftii* (Marsupialia: Vombatidae). PhD thesis, James Cook University, Australia.

Woolnough, A. P., Foley, W. J., Johnson, C. N. and Evans, M. C. (1997). Evaluation of techniques for indirect measurement of body composition in a free-ranging large herbivore, the southern hairy-nosed wombat. *Wildlife Research*, 24: 649-660.