

## OPEN ACCESS

The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at [www.threatenedtaxa.org](http://www.threatenedtaxa.org). All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.



# Journal of Threatened Taxa

Building evidence for conservation globally

[www.threatenedtaxa.org](http://www.threatenedtaxa.org)

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

## NOTE

**ANIMAL-FUNGAL INTERACTIONS 2: FIRST REPORT OF MYCOPHAGY BY THE EASTERN EUROPEAN HEDGEHOG *ERINACEUS CONCOLOR* MARTIN, 1837 (MAMMALIA: EULIPOTYPHILA: ERINACEIDAE)**

Todd F. Elliott, James M. Trappe & Aziz Türkoğlu

26 August 2018 | Vol. 10 | No. 9 | Pages: 12277–12279  
10.11609/jott.4350.10.9.12277-12279



For Focus, Scope, Aims, Policies and Guidelines visit <http://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0>  
For Article Submission Guidelines visit <http://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions>  
For Policies against Scientific Misconduct visit <http://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2>  
For reprints contact [info@threatenedtaxa.org](mailto:info@threatenedtaxa.org)

## Partners



صندوق محمد بن زايد  
للمحافظة على  
الكائنات الحية  
The Mohamed bin Zayed  
SPECIES CONSERVATION FUND



zooh!  
ZÜRICH

Member



Publisher & Host



Threatened Taxa



## ANIMAL-FUNGAL INTERACTIONS 2: FIRST REPORT OF MYCOPHAGY BY THE EASTERN EUROPEAN HEDGEHOG *ERINACEUS CONCOLOR* MARTIN, 1837 (MAMMALIA: EULIPOTYPHILA: ERINACEIDAE)

Todd F. Elliott<sup>1</sup> , James M. Trappe<sup>2</sup>  & Aziz Türkoğlu<sup>3</sup> 

<sup>1</sup>Ecosystem Management, University of New England, Armidale, NSW 2351, Australia

<sup>2</sup>Department of Forest Ecosystems and Society, Oregon State University, Corvallis, Oregon 97331, USA

<sup>3</sup>U.S. Forest Service, Pacific Northwest Research Station, Forestry Sciences Laboratory Corvallis, Oregon 97331, USA

<sup>3</sup>School of Environmental and Forest Sciences, Box 35 2100, University of Washington Seattle, WA 98195-2100, USA

<sup>1</sup>toddfelliott@gmail.com (corresponding author),

<sup>2</sup>trappej@gmail.com, <sup>3</sup>azizturk@uw.edu

The fungal species frequently eaten by small mammals often fruit below ground (hypogeous fungi) and have evolved a dependence on animals for their spore dispersal (Maser et al. 1978, 2008; Cazares & Trappe 1994; Trappe & Claridge 2005; Vernes & Dunn 2009; Elliott & Trappe 2018). The digging required to excavate these fungi enhances soil aeration and allows for better water penetration and soil hydration (Flemming et al. 2013). Most fungi consumed by animals are mycorrhizal and serve important functions as symbionts in soil nutrient acquisition and exchanges with plants (Maser et al. 1978; Colgan & Claridge 2002; Schickmann et al. 2012). Many small mammals depend on hypogeous fungi as a staple food, and numerous larger animals opportunistically

feed on them (Fogel & Trappe 1978; Claridge & Trappe 2005).

Fungi contain important dietary components such as water, essential amino acids, protein, fat, carbohydrates, and crude fiber (Hussain & Al-Ruqaie 1999; Claridge & Trappe 2005; Wallis et al. 2012). Fungi appear in a wide array of animals' diets around the world (Fogel & Trappe 1978; Blaschke & Bäumler 1989; Claridge & May 1994; Hanson et al. 2003; Schickmann et al. 2012). Despite the prevalence and importance of fungi as animal food, mycophagy has been understudied/unreported even among otherwise well-researched animal species. For example, we provide the first confirmed documentation of mycophagy by *Erinaceus concolor* (Martin), the Eastern European Hedgehog. This small mammal has been frequently reported to eat insects, snakes, and plant matter (Özen 2006). Naem et al. (2015) note that "...mushrooms may supplement the diet" of the West European Hedgehog, *E. europaeus* (L.); but their evidence for this statement is unclear and the species of mushrooms are not indicated. New Zealand has a diversity of hypogeous fungi and a large population of invasive hedgehogs, *E. europaeus*, that likely consume local fungi, but this has yet to be confirmed (Wood et al. 2015). As part of this study we conducted microscopic examinations of 30 scats collected from *E. europaeus* in New Zealand and we found no fungal spores. These scats were collected by collaborators and we were unable to determine if many fungi were fruiting during



ISSN 0974-7907 (Online)  
ISSN 0974-7893 (Print)

OPEN ACCESS



DOI: <https://doi.org/10.11609/jott.4350.10.9.12277-12279>

Editor: Jamie Wood, Landcare Research, Lincoln, New Zealand.

Date of publication: 26 August 2018 (online & print)

Manuscript details: Ms # 4350 | Received 21 June 2018 | Finally accepted 25 July 2018

Citation: Elliott, T.F., J.M. Trappe & A. Turkoglu (2018). Animal-fungal interactions 2: first report of mycophagy by the Eastern European Hedgehog *Erinaceus concolor* Martin, 1837 (Mammalia: Eulipotyphla: Erinaceidae). *Journal of Threatened Taxa* 10(9): 12277–12279; <https://doi.org/10.11609/jott.4350.10.9.12277-12279>

Copyright: © Elliott et al. 2018. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use of this article in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

Funding: The Scientific and Technological Research Council of Turkey, project number T-BAG-111T530.

Competing interests: The authors declare no competing interests.

Acknowledgements: Funding from The Scientific and Technological Research Council of Turkey, project number T-BAG-111T530, enabled the fieldwork. We are grateful to Pouri Rakete-Stones for collecting scats from New Zealand populations of hedgehogs. We appreciate input from Allein Stanley of the Schiele Museum, Bob and Babs Strickland of Walnut Creek Preserve, Andrew Claridge of New South Wales Office of Environment and Heritage, and Karl Vernes, University of New England, New South Wales. We appreciate the help of Şerife Çaka, Mehrican Yaratankul Güngör, and Cansu Korkmaz in specimen processing and operation of the microscope lab.




the sampling period.

Santana et al. (2010) mention African Four-toed Hedgehogs, *Atelerix albiventris* (Wagner), eating fungi, but it was not recorded how this or the species of fungus consumed was determined. Skinner & Chimimba (2005) mention fungi in the diet of the southern African Hedgehog *A. frontalis* (A. Smith), but we were unable to determine the original source of their report, and they do not indicate the identity of the fungi consumed. Members of the hedgehog genera *Hemiechinus*, *Mesechinus*, and *Paraechinus* have either had inadequate seasonal dietary studies or insufficient microscopic examination of fecal or stomach samples to determine their mycophagist status.

During mycological fieldwork in March and April 2014 in Muğla-Ula, Turkey, we encountered an Eastern European Hedgehog. *Erinaceus concolor* (Image 1a) crossing the road in a plantation of Umbrella Pine *Pinus pinea* (L.). The understory was predominantly Kermes Oak *Quercus coccifera* (L.), and *Cistus* spp. accompanied by a diverse ectomycorrhizal fungal community. Our interest in mycophagy and hypogeous fungal spore dispersal led us to wait and collect a fecal sample from

the animal. The fecal sample we collected was dried and once at the laboratory, several small pieces were placed with forceps into drops of ethanol on microscope slides. After several minutes the ethanol evaporated and the structures softened. Then water and a cover slip were added and the slide studied with a binocular compound microscope at  $\times 100$ ,  $\times 400$ , and  $\times 1000$  magnification. Additional slides were mounted in Meltzer's reagent to test for taxonomically diagnostic staining reactions of fungal structures; all tests were nonresponsive. Micrographs were taken in water. Percentage volume of fungal material in the slide mounts was estimated by visual scanning. To increase sample size we attempted to find additional individuals to collect scats by spotlighting, but the thick understory and brief time at the site prevented success.

Pieces of fungal tissue and masses of spores composed at least 90% of the Turkish fecal sample. The fungal tissues were characteristic of the false truffle genus *Rhizopogon* (Image 1b). Fungal surveys in the area where the *E. concolor* was found revealed prolific fruitings of *Rhizopogon vulgaris* (Image 1b).



**Image 1.** a - The Eastern European Hedgehog that was the source of the fecal sample examined; b - Fruiting bodies of the false truffle *Rhizopogon vulgaris* found near where the hedgehog was foraging, surface and cross-sectional views; c - Piece of *R. vulgaris* peridium extracted from hedgehog fecal sample; d - Spores of *R. vulgaris* that comprised the majority of the hedgehog fecal sample.

This observation provides strong evidence of gaps in our understanding of the dietary behavior of this and other species of hedgehogs. Historically, the use and importance of fungi has been frequently overlooked in animal dietary studies, and that is likely true for hedgehogs. We had one fecal sample from one individual, insufficient for broad conclusions on behavior and dietary preferences of *E. concolor* or related species, but the large volumes of spores and chunks of fungal tissue in the sample examined (Image 1 c,d) indicate preferential or opportunistic consumption of *Rhizopogon vulgaris* over other food sources. Our estimate of 90% fungus by volume in the scat sampled shows that Eastern European Hedgehogs are opportunistic or possibly preferential mycophagists and we encourage researchers working on this and other species of hedgehogs to apply similar methods to test that hypothesis.

## References

- Blaschke, H. & W. Bäumler (1989). Mycophagy and spore dispersal by small mammals in Bavarian forests. *Forest Ecology and Management* 26(4): 237–245; [http://doi.org/10.1016/0378-1127\(89\)90084-4](http://doi.org/10.1016/0378-1127(89)90084-4)
- Cázares, E. & J.M. Trappe (1994). Spore dispersal of ectomycorrhizal fungi on a glacier forefront by mammal mycophagy. *Mycologia* 86(4): 507–510; <http://doi.org/10.2307/3760743>
- Claridge, A.W. & J.M. Trappe (2005). Sporocarp mycophagy: nutritional, behavioral, evolutionary and physiological aspects, pp. 599–611. In: Dighton, J., J.F. White & P. Oudemans (eds.). *The Fungal Community - Its Organization and Role in the Ecosystem*. Taylor and Francis, Boca Raton, FL, USA; <http://doi.org/10.1201/9781420027891.ch29>
- Claridge, A.W. & T.W. May (1994). Mycophagy among Australian mammals. *Austral Ecology* 19(3): 251–275; <http://doi.org/10.1111/j.1442-9993.1994.tb00489.x>
- Colgan, W. III & A.W. Claridge (2002). Mycorrhizal effectiveness of *Rhizopogon* spores recovered from fecal pellets of small forest-dwelling mammals. *Mycological Research* 106(3): 314–320.
- Elliott, T.F. & J.M. Trappe (2018). A worldwide nomenclature revision of sequestrate *Russula* species. *Fungal Systematics and Evolution* 1(1): 229–242; <http://doi.org/10.3114/fuse.2018.01.10>
- Flemming, P.A., H. Anderson, A.S. Prendergast, M.R. Bretz, L.E. Valentine & G.E. Hardy (2013). Is the loss of Australian digging mammals contributing to a deterioration in ecosystem function? *Mammal Review* 44(2): 94–108.
- Fogel, R. & J.M. Trappe (1978). Fungus consumption (mycophagy) by small mammals. *Northwest Science* 52(1): 1–31.
- Hanson, A.M., K.T. Hodge & L.M. Porter (2003). Mycophagy among primates. *Mycologist* 17(1): 6–10; <http://doi.org/10.1017/S0269915X0300106X>
- Hussain, G. & I.M. Al-Ruqaie (1999). Occurrence, chemical composition, and nutritional value of truffles: An overview, *Pakistan Journal Biological Science* 2(2): 510–514.
- Maser, C., A.W. Claridge & J.M. Trappe (2008). *Trees, Truffles and Beasts - How Forests Function*. Rutgers University Press, New Brunswick, NJ, USA.
- Maser, C., J.M. Trappe & R.A. Nussbaum (1978). Fungal-small mammal inter-relationships with emphasis on Oregon coniferous forests. *Ecology* 59(4): 799–809.
- Naem, S., B. Pourreza & T. Gorgani-Firouzjaee (2015). The European Hedgehog (*Erinaceus europaeus*) as a reservoir for helminth parasites in Iran. *Vet Res Forum* 6(2): 149–153.
- Özen, A. S. (2006). Some biological, ecological and behavioural features of *Erinaceus concolor* Martin, 1838 (Mammalia: Insectivora) in Turkey. *Gazi University Journal of Science* 19(2): 91–97.
- Santana, E.M., H.E. Jantz & T.L. Best (2010). *Atelerix albiventris* (Erinaceomorpha: Erinaceidae). *Mammalian Species* 42(1): 99–110; <http://doi.org/10.1644/857.1>
- Schickmann, S., A. Urban, K. Kräutler, U. Nopp-Mayr & K. Hackländer (2012). The interrelationship of mycophagous small mammals and ectomycorrhizal fungi in primeval, disturbed and managed Central European mountainous forests. *Oecologia* 170(2): 395–409; <http://doi.org/10.1007/s00442-012-2303-2>
- Skinner, J.D. & C.T. Chimimba (2005). *The Mammals of the Southern African Sub-region*. Cambridge University Press, Cambridge, UK; <http://doi.org/10.1017/CBO9781107340992>
- Trappe, J.M. & A.W. Claridge (2005). Hypogeous fungi: evolution of reproductive and dispersal strategies through interactions with animals and plants, pp. 613–622. In: Dighton, J., J.F. White & P. Oudemans (eds.). *The Fungal Community - Its Organization and Role in the Ecosystem*. Taylor and Francis, Boca Raton, FL, USA.
- Vernes, K. & L. Dunn (2009). Mammal mycophagy and fungal spore dispersal across a steep environmental gradient in eastern Australia. *Austral Ecology* 34(1): 69–76; <http://doi.org/10.1111/j.1442-9993.2008.01883.x>
- Wallis, I., A.W. Claridge & J.M. Trappe (2012). Nitrogen content, amino acid composition and digestibility of fungi from a nutritional perspective in animal mycophagy. *Fungal Biology* 116(5): 590–602; <http://doi.org/10.1016/j.funbio.2012.02.007>
- Wood, J.R., I.A. Dickie, H.V. Moeller, D.A. Peltzer, K.I. Bonner, G. Rattray & J.M. Wilmshurst (2015). Novel interactions between non-native mammals and fungi facilitate establishment of invasive pines. *Journal of Ecology* 103(1): 121–129; <http://doi.org/10.1111/1365-2745.12345>





OPEN ACCESS



The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at [www.threatenedtaxa.org](http://www.threatenedtaxa.org). All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.

ISSN 0974-7907 (Online); ISSN 0974-7893 (Print)

August 2018 | Vol. 10 | No. 9 | Pages: 12147–12298

Date of Publication: 26 August 2018 (Online & Print)

DOI: 10.11609/jott.2018.10.9.12147-12298

[www.threatenedtaxa.org](http://www.threatenedtaxa.org)

### Article

**Appearances are deceptive: molecular phylogeny recovers the Scaly Gecko *Hemidactylus scabriceps* (Reptilia: Squamata: Gekkonidae) as a member of a scansorial and rupicolous clade**

-- Achyuthan N. Srikanthan, Gandla Chethan Kumar, Aishwarya J. Urs & Sumaithangi Rajagopalan Ganesh, Pp. 12147–12162

### Communications

**Foraging and roosting ecology of the Lesser Dog-faced Fruit Bat *Cynopterus brachyotis* (Mammalia: Chiroptera: Pteropodidae) in southern India**

-- T. Karuppudurai & K. Sripathi, Pp. 12163–12172

**Diversity and status of avifauna in man-made sacred ponds of Kurukshetra, India**

-- Parmesh Kumar & Archana Sharma, Pp. 12173–12193

**Diversity and distribution of freshwater turtles (Reptilia: Testudines) in Goa, India**

-- Trupti D. Jadhav, Nitin S. Sawant & Soorambail K. Shyama, Pp. 12194–12202

**Breeding behaviour of the Coromandel Damselfly *Ceriatrigon coromandelianum* (Fabricius) (Zygoptera: Coenagrionidae) in central India: copulation**

-- Nilesh R. Thaokar, Payal R. Verma & Raymond J. Andrew, Pp. 12203–12209

**The status assessment of *Corynandra viscosa* subsp. *nagarjunakondensis* (Magnoliopsida: Cleomaceae), endemic to Nagarjunakonda, Andhra Pradesh, India**

-- Veeravarapu Hanumantha Rao, Vaidyula Vasudeva Rao, Anuti Baleeshwar Reddy & Vatsavaya Satyanarayana Raju, Pp. 12210–12217

### Short Communications

**New records of termites (Blattodea: Termitidae: Syntermitinae) from Colombia**

-- Olga Patricia Pinzón & Daniel Castro, Pp. 12218–12225

**New reports of thrips (Thysanoptera: Terebrantia: Thripidae) from India**

-- R.R. Rachana & R. Varatharajan, Pp. 12226–12229

**New records of earthworm fauna (Oligochaeta: Glossoscolecidae and Megascolecidae) collected from Satkosia-Baisipalli Wildlife Sanctuary of Odisha, India**

-- Rinku Goswami, Pp. 12230–12234

**Diversity and endemism of butterflies of montane forests of Eravikulam National Park in the Western Ghats, India**

-- E.R. Sreekumar, S. Nikhil, K.G. Ajay & P.O. Nameer, Pp. 12235–12246

### Partners



### Angiosperm diversity of Sonbhadra District, Uttar Pradesh: a checklist

-- Arun Kumar Kushwaha, Lalit Mohan Tewari & Lal Babu Chaudhary, Pp. 12247–12269

### Contribution to the Macromycetes of West Bengal, India: 23–27

-- Meghma Bera, Soumitra Paloi, Arun Kumar Dutta, Prakash Pradhan, Anirban Roy & Krishnendu Acharya, Pp. 12270–12276

### Notes

**Animal-fungal interactions 2: first report of mycophagy by the Eastern European Hedgehog *Erinaceus concolor* Martin, 1837 (Mammalia: Eulipotyphla: Erinaceidae)**

-- Todd F. Elliott, James M. Trappe & Aziz Türkoğlu, Pp. 12277–12279

**Rostral anomaly in a juvenile Spiny Butterfly Ray *Gymnura altavela* (Linnaeus, 1758) (Elasmobranchii: Myliobatiformes: Gymnuridae) from the Canary Islands**

-- Filip Osaer & Krupskaya Narváez, Pp. 12280–12281

**A record after 52 years, and additional description of the emesine assassin bug *Emesopsis nubila* (Hemiptera: Reduviidae: Emesinae) from western India**

-- Balasaheb V. Sarode, Nikhil U. Joshi, Pratik P. Pansare & Hemant V. Ghatge, Pp. 12282–12285

***Gentiana aperta* (Gentianaceae) - a new record to India from Ladakh Himalaya**

-- Mohd Shabir, Priyanka Agnihotri, Jay Krishan Tiwari & Tariq Husain, Pp. 12286–12289

**Notes on *Cinnamomum travancoricum* Gamble (Lauraceae) - a Critically Endangered species from the southern Western Ghats, India**

-- A.J. Robi, P. Sujanalal & P.S. Udayan, Pp. 12290–12293

**A reassessment and lectotypification of the name *Striga masuria* (Buch.-Ham. ex Benth.) Benth. (Orobanchaceae) and its collection from the Western Ghats of India**

-- M. Omalsree & V.K. Sreenivas, Pp. 12294–12297

### Miscellaneous

National Biodiversity Authority

Member



Publisher & Host

