



ELSEVIER

Contents lists available at ScienceDirect

MethodsX

journal homepage: [www.elsevier.com/locate/mex](http://www.elsevier.com/locate/mex)

## Method Article

# A molecular based method for rapid detection of *Salmonella* spp. in poultry dust samples



Md Ahaduzzaman<sup>a,b,\*</sup>, Peter J Groves<sup>c</sup>, Stephen W Walkden-Brown<sup>a</sup>, Priscilla F Gerber<sup>a,\*</sup>

<sup>a</sup> Animal Science, School of Environmental and Rural Science, University of New England, Armidale, NSW 2351, Australia

<sup>b</sup> Faculty of Veterinary Medicine, Chattogram Veterinary and Animal Sciences University, Chattogram 4225, Bangladesh

<sup>c</sup> Faculty of Veterinary Science, The University of Sydney, Camden, NSW 2570, Australia

## A B S T R A C T

Salmonellosis, caused by *Salmonella* spp., is a widely reported foodborne zoonosis frequently associated with ingestion of poultry products. *Salmonella* vaccination of chickens can be used to reduce bacterial shedding and risk of human infection. To determine *Salmonella* burden in chicken farms, culture methods of environmental samples that require a turn-around time of 5–7 days are usually used. Rapid screening using molecular assays such as PCR of pre-enriched broth has been reported for *Salmonella* spp. detection in feed, floor dust, and drag swabs within 2–3 days. Here we report an adaptation of the method for detection of *Salmonella* in poultry dust samples collected using a settle plate method under experimental conditions. Key features:

- Passive dust sample collection using dry settle plates without media suspended from dropper lines of drinkers.
- Small amount of sample required for the pre-enrichment process.
- Quantification of *Salmonella* DNA with high sensitivity using an inexpensive extraction protocol.

© 2021 The Author(s). Published by Elsevier B.V.  
This is an open access article under the CC BY-NC-ND license  
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

## A R T I C L E I N F O

**Method name:** Detection of *Salmonella* in poultry dust

**Keywords:** *Salmonella*, Chicken, Disease monitoring, Poultry dust, qPCR

**Article history:** Received 19 January 2021; Accepted 12 April 2021; Available online 21 April 2021

DOI of original article: [10.1016/j.vaccine.2020.12.044](https://doi.org/10.1016/j.vaccine.2020.12.044)

\* Corresponding authors at: Animal Science, School of Environmental and Rural Science, University of New England, Armidale, NSW 2351, Australia.

E-mail addresses: [zaman.cvasu@gmail.com](mailto:zaman.cvasu@gmail.com) (M. Ahaduzzaman), [pgerber2@une.edu.au](mailto:pgerber2@une.edu.au) (P.F. Gerber).

<https://doi.org/10.1016/j.mex.2021.101356>

2215-0161/© 2021 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license  
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

## Specifications Table

Subject Area:	Veterinary Science and Veterinary Medicine
More specific subject area:	Environmental microbiology
Method name:	Detection of <i>Salmonella</i> in poultry dust
Name and reference of original method:	Gole et al., Shedding of <i>Salmonella</i> in single age caged commercial layer flock at an early stage of lay. International Journal of Food Microbiology, 2014 [5]. De Medici et al., Evaluation of DNA extraction methods for use in combination with SYBR green I real-time PCR to detect <i>Salmonella enterica</i> serotype enteritidis in poultry. Applied and Environmental Microbiology, 2003 [11].
Resource availability:	All consumables and equipment required are described in the manuscript.

## Method details

*Salmonella* spp. cause fowl typhoid and pullorum diseases in poultry and are responsible for non-typhoidal *Salmonella* infection in humans. Prevention of *Salmonella* infection in chicken flocks is an important step in reducing salmonellosis outbreaks in humans. Vaccination with live attenuated vaccine of certain strains such as e.g. *S. Enteritidis*, *S. Typhimurium* and *S. Gallinarum* has been commonly used in the field to prevent the disease, and proven to be useful to reduce bacterial shedding [1,2]. To monitor *Salmonella* shedding patterns in vaccinated and unvaccinated flocks, a combination of litter, drag swab, dust swab, feed and egg belt swab are usually used [3,4]. However, extraction of genomic DNA from these various sample types is expensive and time-consuming due to the requirement for faecal or soil DNA extraction kits [4] or overnight processing of extracted DNA samples [5], restricting the success of routine monitoring of *Salmonella* in poultry flocks.

Dust sample collection using settle plates has been a useful tool for routine monitoring of pathogens in poultry flocks [6]. The use of media-free, dry settle plates enables easy collection of a stable material, dry dust, from which nucleic acid extraction can effectively be done using inexpensive extraction kits [7]. Even less costly methods such as the boiling method of DNA extraction have been effective for *Salmonella* detection from different food specimens [8]. However, it is unknown if this method would be effective for *Salmonella* detection in dust samples.

This study reports the longitudinal tracking of *Salmonella* using dust samples collected on dry settle plates from an experimental flock exposed to *Salmonella enterica* serovar Typhimurium DT 135 at 1 day of age, based on PCR detection of *Salmonella invA* gene in dust with and without enrichment and with two methods of DNA extraction: a commercial kit and a method based on boiling of samples enriched over-night in buffered peptone water (BPW). The *invA* gene-based PCR was chosen for its ability to detect multiple *Salmonella* spp. as a screening tool. Furthermore, the samples used for this protocol development were derived from an experimental flock challenged with *S. Typhimurium* PT 135, and no other serotypes were identified by culture and biochemical tests from monitoring of cloacal swabs and drag swabs during the experiment [9]. In samples positive for the *invA* gene DNA, multiplex PCRs would be required for the differentiation of specific *Salmonella* serovars according to the research and diagnostic needs.

## Materials

Autoclave (Hiclve™, HV-85L, Hirayama)  
 Biosafety cabinet class II or Bunsen burner  
 Buffered Peptone Water (BPW, Oxoid, Cat. no. CM0509B, Australia)  
 Colony counter (SCC100, Selby®)  
 Corbett Robot  
 Digital microbalance  
 Distilled water  
 Erlenmeyer flask (conical flask)  
 Glass pipette (10 ml)  
 Heat block  
 Incubator (Labmaster®)

ISOLATE II Genomic DNA Kit (Bioline, Cat. No. BIO-52067, Australia)  
Microcentrifuge tubes  
PCR tubes  
Petri dishes  
Pipette filter tips  
Pipettes (10, 100 and 1000 µl)  
Polypropylene L-shaped spreaders (Wiltronics, Cat. No. TL1350, Australia)  
PYREX® screw cap culture tubes with phenolic caps (Thomas Scientific, Cat. no. 9212C21, USA)  
RNase-free water  
Rotor-gene qPCR system  
*Salmonella enterica* serovar Typhimurium (strain: ACM 3598)  
Screw cap bottle/media bottle (Thomas Scientific, Cat. no.1743G56, USA)  
Settle plates  
Test tube rack  
Test tubes  
Vortex  
Xylose Lysine Deoxycholate (XLD) Agar (Oxoid, Cat. no. CM0469B, Australia)

## Protocol


### Sample collection

Dust samples ( $n = 40$ ) were collected in a curtain sided conventional experimental chicken house divided into 16 pens of which 7 pens containing 210 Hy-Line Brown layer chickens (30 birds/pen) and 9 pens remaining empty as previously described [9]. Briefly, the birds of 6 pens were exposed to a field isolate of *S. Typhimurium* DT 135 using a seeder bird technique at 0 day of age and birds of remaining pen were unchallenged control (Supplementary Table 1). At weeks 10 and 16 of age, the birds were vaccinated with a live *Aro-A* deletion mutant vaccine against *S. Typhimurium* strain STM1 (VaxsafeST, Bioproperties Australia, Ringwood, Victoria) in 4 pens of the 6 challenged pens by intramuscular injection of approximately  $2.3 \times 10^7$  colony forming unit (CFU)/bird.

Dust samples were collected at 8.5, 9, 9.5, 12, 14, 15, 16, 18 and 30 weeks of age using a dry plastic settle plate with a surface area of 520 cm<sup>2</sup>. Settle plates were installed in five sites within the house near pens 13, 14, 17, 19 and 23. Samples from each site were collected into separate zip-lock bags by shaking and scraping the dust deposited on the surface of the settle plate into the bag then sealing it. After collection, samples were stored at -20 °C and then transported to the University of New England for further storage at -80 °C prior to use. Dust samples were homogenised by vortexing and divided into two aliquots before further testing. Additionally, drag swabs (1 swab/pen) of individual bird pens were collected from 6 challenged pens using two tampons per pen soaked in sterile BPW at 8, 12, 16 and 18 weeks of age and were cultured for the presence of *Salmonellae* using the Australian Standard method (AS 5013.10-2009) [10]. This involved incubation and inoculation into modified semisolid Rappaport-Vassiliadis medium, Hektoen and xylose-lysine-deoxycholate media and chromogenic agar. Presumptive *Salmonella* colonies were confirmed serologically with poly-O and poly-H antisera by the slide agglutination technique after sub-culture on a nutrient agar slope [9]. The experiment was approved by the University of Sydney Animal Ethics Committee (Approval no. 2017/1207).

### Dust sample processing and extraction

1. Samples enriched for *Salmonella*: Approximately 100 mg of dust was resuspended in 10 ml of BPW and mixed by vortexing for 20–30 s. For samples for which less than 100 mg was available, 10 to 50 mg of dust was used. The samples were incubated at 37 °C for 18 h. Enriched samples were extracted by two methods.
  - (i) Boiling method: 1 ml of enriched sample was pipetted from the middle of tube (Fig. 1) and was transferred to 1.5 ml microcentrifuge tubes and centrifuged for 10 min at  $13,000 \times g$ . The supernatant was discarded carefully. The pellet was resuspended using



**Fig 1.** Buffered peptone water broth media after incubation of 100 mg of dust samples in 10 ml of BPW at 37 °C for 18 h. Dust formed a sediment at the bottom of the broth culture tubes and turbidity was observed mostly in the middle of the tube. The arrow mark indicates the pipetting site for transfer of the pre-enriched sample for DNA extraction.

300  $\mu$ l of molecular water by vortexing followed again by centrifugation at 13,000  $\times$  g for 5 min and discarding of the supernatant. The pellet was resuspended using 200  $\mu$ l of molecular grade water by vortexing. The microcentrifuge tube was incubated at 100 °C for 15 min using a heat block then immediately chilled on ice for 15 min. The sample was then centrifuged for 13,000  $\times$  g for 5 min. The supernatant was carefully transferred to a new microcentrifuge tube and incubated again at 100 °C for 10 min then chilled immediately on ice for 10 min. The sample was then mixed with a pipette prior to use in the PCR reaction. This method was described previously by De Medici et al. [11], but the centrifugation was performed at 13,000  $\times$  g in this experiment instead of 14,000  $\times$  g.

- (ii) Extraction kit: 1 ml of the pre-enriched sample was extracted using the ISOLATE II Genomic DNA kit according to the manufacturer's instructions. Extracted DNA was eluted in a volume of 100  $\mu$ l.
2. Non-enriched dust samples: Five mg of dust was extracted using the ISOLATE II Genomic DNA kit and DNA eluted in a final volume of 100  $\mu$ l as described by Ahaduzzaman et al. [7]. The boiling method was not used to extract DNA from non-enriched dust samples because of the reduced sensitivity of this method and the insufficient amount of dust samples to carry on this test.
3. DNA concentration: Extracted DNA quality and quantity were measured using a spectrophotometer (NanoDropND-1000 UV-Vis) and stored at -20 °C until use.

#### Real-time qPCR for *Salmonella*

1. Development of *Salmonella* standards: A suspension of *S. Typhimurium* was 10-fold diluted from  $10^{-1}$  to  $10^{-8}$  in molecular grade water. For the preparation of standards, DNA was extracted from each dilution by both the boiling method and the commercial extraction kit as described above. To quantify the *Salmonella* counts in each standard, 100  $\mu$ l of the same bacterial suspension was plated by using spread plate technique as described by Sanders [12] on XLD agar plates with 6 replicates per dilution and incubated for 24 h at 37 °C. Colony counting was performed using a colony counter. Based on the number of enumerated colonies on agar plates, the concentration of *Salmonella* in the original suspension was calculated in CFU/ml. CFU was estimated using the standard formula of  $CFU = (\text{Number of colony} \times \text{dilution factor})/\text{amount plated}$ .

The sensitivity and amplification efficiency for both real-time PCR assays were tested by amplification of 10-fold serial dilutions of *S. Typhimurium* DNA samples. The threshold cycles were plotted against the  $\log_{10}$  values of the initial number of *S. Typhimurium* DNA copies in the PCR to construct a standard curve for each assay. Linearity was observed with the following linear regression

**Table 1**

Proportion of extracted DNA quality extracted from poultry dust samples with and without overnight pre-enrichment in pre-enrichment in BPW using extraction kit and boiling methods. A 260/280 ratio of 1.8–2.0 is indicative of high DNA quality.

Spectrophotometer DNA quality assessment parameter	Non-enriched KitN (%)		Enriched KitN (%)		Enriched BoilN (%)	
	260/280	260/230	260/280	260/230	260/280	260/230
< 1	40 (100)	39 (97.5)	3 (7.5)	4 (10)	12 (30)	12 (30)
1.0–1.49	0 (0)	1 (2.5)	4 (10)	4 (10)	3 (7.5)	27 (67.50)
1.5–1.79	0 (0)	0 (0)	5 (12.50)	6 (15)	4 (10)	1 (2.50)
≥ 1.8–2.0	0 (0)	0 (0)	28 (70)	26 (65)	21 (52.50)	0 (0)

curves:  $y = -3.77x + 30.23$  and  $y = -4.37x + 35.70$  for the kit and boiling methods of extraction respectively, where  $y$  is the CT value and  $x$  is the  $\log_{10}$  concentration of *Salmonella*. A high coefficient of determination ( $R^2 = 0.999$ ) was attained for both methods, and a high amplification efficiency (87% by kit and 70% by boiling methods) was achieved. Therefore, it is considered that these assays produced an accurate quantification of *Salmonella* when the initial amount of *Salmonella* DNA in the PCR fell in the range of  $1.00 \times 10^0$  to  $1.09 \times 10^5$  copies/reaction for the kit extraction and  $4.85 \times 10^0$  to  $4.85 \times 10^5$  copies/reaction for the boiling method of extraction.

- Real-time qPCR conditions: Extracted DNA was tested for *Salmonella* by a qPCR using primers (forward: 5'-AACGTGTTCCGTGCGTAAT-3' and reverse: 5'-TCCATCAAATTAGCGGAGGC-3') and TaqMan probe (5'-FAM-TGGAAGCGCTCGCATTGTGG-BHQ-1-3') targeting the *invA* gene [13]. Each 25  $\mu$ l of real-time PCR reaction contained 0.5  $\mu$ l of each primer (0.5 mM), 0.5  $\mu$ l of probe (10 mM), 5  $\mu$ l of template DNA (1:10 dilution in molecular water), 12.5  $\mu$ l of 2  $\times$  master mix, and 6  $\mu$ l of nuclease-free water. PCR conditions were 95 °C for 3 min, was followed by 40 cycles of 15 s at 95 °C and 30 s at 60 °C. The standards developed following DNA extraction by each extraction method were used to quantify samples extracted by the same method. The results were analysed by the Rotor-Gene Q version 2.3.1.49 software and reported as *Salmonella*  $\log_{10}$  CFU per gram of dust.

### Data analysis

Statistical analyses were performed using JMP v.14 software (SAS Institute, Cary, NC, USA). *Salmonella* CFU per g of dust (with or without enrichment) was analysed and expressed on the  $\log_{10}$  scale. The proportion of samples positive for *Salmonella* are presented as number of positive samples by the total number of samples sampled (%) in each category (non-enriched kit, enriched kit, enriched boil, and drag swab) and McNemar's test was used for pairwise comparison to determine whether the proportions of PCR-positive samples were significantly different. Prevalence of *Salmonella* in drag swab and enriched dust at given sampling day linear association was explored. Analysis of variance was used to test the effects of sampling time and treatment effects with means and standard errors reported. Association between the methods of extraction of *Salmonella* CFU was analysed using an intraclass correlation coefficient (ICC) test.

### Method validation

Quality assessment results of extracted DNA are presented in Table 1. The 260/280 and 260/230 ratio were <1 in case of 97.5 to 100% of samples when DNA extraction was done from non-enriched dust samples using the extraction kit, and between 1.8 and 2 on 52.5 and 70% of samples when extraction was done from enriched dust using boiling method and extraction kit, respectively. This indicates that enrichment of samples prior to DNA extraction significantly improves the quality of the obtained DNA.

The proportion of dust samples positive for *Salmonella* increased with overnight enrichment (Table 2) particularly when bacterium concentrations were low. Overall, 62.5% (25/40) of samples were positive after enrichment while 40% (16/40) of non-enriched samples were positive ( $P = 0.003$ ).

**Table 2**

Ratio of *Salmonella* qPCR positive samples in dust with and without overnight pre-enrichment in pre-enrichment in BPW, compared to culture method of drag swabs.


Age (weeks)	Salmonella positive/n (%)		P-value <sup>1</sup>	Drag swab	P-value <sup>2</sup>
	Dust with enrichment	Dust without enrichment			
8–8.5	4/4 (100)	4/4 (100)	1.00	6/6 (100)	1.00
9	4/4 (100)	2/4 (50)	0.10	-	-
9.5	2/2 (100)	2/2 (100)	1.00	-	-
12	5/5 (100)	5/5 (100)	1.00	6/6 (100)	1.00
14	4/5 (80)	0/5 (00)	0.009	-	-
15	1/5 (20)	1/5 (20)	1.00	-	-
16	4/5 (80)	2/5 (40)	0.16	4/6 (66.7)	1.00
18	1/5 (20)	0/5 (00)	0.29	4/6 (66.7)	0.08
30	0/5 (00)	0/5 (00)	1.00	-	-
Total	25/40 <sup>a</sup>	16/40 <sup>b</sup>	0.003	20/24 <sup>a</sup>	0.08

Different superscript letters (<sup>ab</sup>) indicate statistical significance

<sup>1</sup> Comparison between dust with enrichment and dust without enrichment.

<sup>2</sup> Comparison between dust with enrichment and drag swab.


-Sample not tested.



**Fig. 2.** *Salmonella* load in dust samples enriched or not by incubation in BPW with DNA extraction by a commercial kit or a simple sample boiling method. Dust samples were collected at the indicated time points from experimental flock challenged with a field isolate of *S. Typhimurium* DT 135 using a seeder bird technique at 1 day of age and vaccinated with a non-propagative strain of live *Salmonella* vaccine at 10 and 16 weeks of age.

When *Salmonella* levels were high, from weeks 8 to 12 of bird age (Fig. 2) all samples were positive irrespective of enrichment. However, for the samplings at 14–18 weeks, when *Salmonella* levels were 2–6 logs lower, only 20–40% samples were positive without enrichment while 20–80% of samples were positive after enrichment. No difference was observed in the detection pattern of *Salmonella* between kit and boiling method of extraction after enrichment ( $P = 1.00$ ).

Compared to detection using drag swabs, there was no significant difference in the detection rates of enriched dust and drag swab samples on the same sampling days ( $P = 0.08$ ) (Table 2). Linear



**Fig. 3.** A. Intraclass correlation coefficient of  $\text{Log}_{10}$  *Salmonella* CFU/g of BPW enriched dust extracted by Kit and boiling method. Here, the intraclass correlation coefficient (ICC) = 0.99, degrees of freedom (df) = 39 and  $P < 0.0001$ . B. intraclass correlation coefficient of  $\text{Log}_{10}$  *Salmonella* CFU/g of non-enriched dust extracted by kit and BPW enriched dust extracted by kit. Here, the ICC value is 0.72, degrees of freedom (df) = 39 and  $P < 0.0001$ .

association between *Salmonella* prevalence (%) in dust and drag swabs showed a strong positive relationship ( $r = 0.76$ ), suggesting that *Salmonella* shed actively in faeces and found in litter can be detected in settled dust in the environment.

The level of *Salmonella* detected in dust (CFU/g) also depended on overnight enrichment of dust ( $P = 0.004$ ) and bird age/day post challenge infection ( $P < 0.0001$ ) but not by the extraction method used on enriched samples (Fig. 2). The level was similar ( $P = 0.94$ ) for enriched samples extracted using a commercial kit ( $3.70 \pm 0.31$ ) or the boiling method ( $3.76 \pm 0.31$ ) and was lower ( $1.60 \pm 0.31$ ) for non-enriched samples extracted using the commercial kit ( $P < 0.001$ ). Levels declined gradually to low levels at week 15, followed by a small increase at week 16, then a decline to low levels at week 18 with all samples negative at week 30.

There was an almost perfect agreement between the real-time PCR results from enriched samples following DNA extraction by the boiling and commercial kit methods (ICC = 0.99,  $P < 0.0001$ , Fig. 3A). On the other hand, there was a moderate agreement between PCR values of enriched and non-enriched dust extracted by the commercial kit method (ICC = 0.72,  $P < 0.0001$ , Fig. 3B).

Longitudinal tracking of *Salmonella* in dust samples was possible after birds were challenged with *Salmonella*. Enrichment of dust samples in BPW improved sensitivity and *Salmonella* detection pattern using qPCR. No difference was observed between the boiling method and extraction kit method. Collection of dust samples using settle plates followed by enrichment in BPW, extraction of DNA using the boiling method and qPCR analysis may offer a cost effective and practical method of *Salmonella* monitoring in poultry populations. Dust is a stable, easily collected population level sample and qPCR of dust for pathogen detection offers advantages in speed, simplicity and potentially sensitivity over methods based on culture of the organism.

### Additional information

Classical methods for detection of *Salmonella* involve analysis of faeces or cloacal swabs, or both. However, as classical methods are impractical for large surveys, gauze pad or tampon based drag swabs being the most widely used for flock level detection often require different diluents (e.g. BPW) to collect samples and considerable amount of time (20–30 min./shed) [14,15]. Dust sampling is comparatively easier than drag swabbing and previously been used in a nationwide survey in Japan [16]. In this study *Salmonella* in enriched dust samples collected using settle plates could-be detected throughout the 22 weeks observation period following experimental infection and

vaccination. Another study detected *Salmonella* in dust samples collected by drag swab samples or scraped dust samples from a caged-layer shed floor and found that *Salmonella* was constantly detectable in dust over the 26 weeks observation period following oral *S. Typhimurium* challenge and 3 doses of live *Salmonella* vaccine [4]. In another study, randomly collected litter materials or soil has been used for the detection of *Salmonella* in broiler farms [17]. Collection of such samples may under or over-estimate the presence of *Salmonella* due to the localised nature of these samples. Dust samples represent material dispersed and mixed in the air, and studies with other pathogens have demonstrated that location of settle plates within poultry houses does not influence the level of pathogen detected [18]. In fact, Groves et al. [9] describes that dust samples of the untreated unchallenged group separated by 6 m from the nearest challenged pen were positive at 8.5 and 12 weeks of age when the dust samples of challenged groups were also positive. In that study, drag swab samples were only positive in the control pen when birds were 18-week old, suggesting that testing of settled dust samples provides a useful complementary tool for *Salmonella* monitoring.

Compared to culture methods, PCR analysis of dust samples can reduce the turn-around time from 4–5 days to 1–2 days [19]. Detection of *Salmonella* from any given sample by culture prior to PCR analysis may reduce sensitivity [20]. In the present study enrichment of dust sample in BPW for 18 h increased the sensitivity of detection of *Salmonella* with 25/40 and 16/40 of samples positive for enriched and non-enriched samples respectively. This is consistent with other studies indicating that enrichment in BPW for 18–24 h followed by PCR detection can improve sensitivity of detection in faeces [19], chicken meat [20], feed and water [21], clinical samples [22] and environmental samples [22]. Thomason et al. [23] reported that BPW increased the detection of *Salmonella* by about 25% in absolute percentage terms from environmental samples. In this study, pre-enrichment of dust in BPW increased the recovery of *Salmonella* from 40% to 62.5%.

In this study, no difference was observed between pre-enriched samples extracted by boiling method and samples extracted by DNA extraction kit in the *Salmonella* detection pattern ( $P=1.00$ ) and CFU count ( $P=0.94$ ) using real-time PCR. Overall detection rates for *Salmonella* in dust was 62.5% by either method. The lower limits of detection of *Salmonella* in this study were  $1.00 \times 10^0$  copies/reaction for the extraction kit and  $4.85 \times 10^0$  for the boiling method. These were similar to previous reported detection rates in foodstuff [24,25]. De Medici et al. [11] compared four methods of DNA extraction from poultry samples pre-enriched in a broth media and reported boiling method as the preferred extraction method because of simplicity. Similarly, Sweeney et al. [26] compared three methods of DNA extraction from an automated broth culture system and found that boiling extraction is the most suitable method for real-time qPCR. Moreover, the boiling method was also found suitable in several studies for human faecal and oral microbiome analysis [27,28]. The present study indicates that the boiling method could be used to monitor *Salmonella* in poultry shed dust using PCR, however, when the quality of extracted DNA is paramount such as for high-throughput sequencing applications, the use of an extraction kit would be more appropriate.

## Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgments

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors. We thank Sithara Ralapanawe, Alison Pankhurst, Michael Raue, Chake Keerqin and Md Zohorul Islam for their support at different stages of this work.

## Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:[10.1016/j.mex.2021.101356](https://doi.org/10.1016/j.mex.2021.101356).



## References

- [1] M.C. Cerquetti, M.M. Gherardi, Vaccination of chickens with a temperature-sensitive mutant of *Salmonella enteritidis*, *Vaccine* 18 (11-12) (2000) 1140–1145.
- [2] T.S. Desin, W. Köster, A.A. Potter, *Salmonella* vaccines in poultry: past, present and future, *Expert Rev. Vaccines* 12 (1) (2013) 87–96.
- [3] F.C. Dórea, et al., Effect of *Salmonella* vaccination of breeder chickens on contamination of broiler chicken carcasses in integrated poultry operations, *Appl. Environ. Microbiol.* 76 (23) (2010) 7820–7825.
- [4] P. Sharma, et al., Shedding of *Salmonella* Typhimurium in vaccinated and unvaccinated hens during early lay in field conditions: a randomised controlled trial, *BMC Microbiol.* 18 (1) (2018) 78.
- [5] V.C. Gole, et al., Shedding of *Salmonella* in single age caged commercial layer flock at an early stage of lay, *Int. J. Food Microbiol.* 189 (2014) 61–66.
- [6] S.W. Walkden-Brown, et al., Development, application, and results of routine monitoring of Marek's disease virus in broiler house dust using real-time quantitative PCR, *Avian Dis.* 57 (2s1) (2013) 544–554.
- [7] M. Ahaduzzaman, et al., A practical method for assessing infectious laryngotracheitis vaccine take in broilers following mass administration in water: spatial and temporal variation in viral genome content of poultry dust after vaccination, *Vet. Microbiol.* 241 (2019) 108545.
- [8] C. Löfström, et al., Detection of *Salmonella* in meat: comparative and interlaboratory validation of a noncomplex and cost-effective pre-PCR protocol, *J. AOAC Int.* 95 (1) (2012) 100–104.
- [9] P.J. Groves, et al., Can a combination of vaccination, probiotic and organic acid treatment in layer hens protect against early life exposure to *Salmonella* Typhimurium and challenge at sexual maturity? *Vaccine* 39 (5) (2020) 815–824.
- [10] AS 5013.10-2009, Microbiology of food and animal feeding stuffs: horizontal method for the detection of *Salmonella* spp. AS 5013.10-2009, DAWE Australian Govt. Dep. Agric. Water Environ. 4 (6) (2020) 1–2.
- [11] D. De Medici, et al., Evaluation of DNA extraction methods for use in combination with SYBR green I real-time PCR to detect *Salmonella enterica* serotype enteritidis in poultry, *Appl. Environ. Microbiol.* 69 (6) (2003) 3456–3461.
- [12] E.R. Sanders, Aseptic laboratory techniques: plating methods, *J. Vis. Exp.* (63) (2012).
- [13] C.M. Cheng, et al., Rapid detection of *Salmonella* in foods using real-time PCR, *J. Food Prot.* 71 (12) (2008) 2436–2441.
- [14] A. Pavic, P.J. Groves, J.M. Cox, Development and validation of a drag swab method using tampons and different diluents for the detection of members of *Salmonella* in broiler houses, *Avian Pathol.* 40 (6) (2011) 651–656.
- [15] E. Mallinson, et al., Monitoring poultry farms for *Salmonella* by drag-swab sampling and antigen-capture immunoassay, *Avian Dis.* (1989) 684–690.
- [16] E. Iwabuchi, et al., Nationwide survey of *Salmonella* prevalence in environmental dust from layer farms in Japan, *J. Food Prot.* 73 (11) (2010) 1993–2000.
- [17] J. Brooks, et al., Cultivation and qPCR detection of pathogenic and antibiotic-resistant bacterial establishment in naïve broiler houses, *J. Environ. Qual.* 45 (3) (2016) 958–966.
- [18] T.V. Nguyen, et al., Spatial and temporal variation of Marek's disease virus and infectious laryngotracheitis virus genome in dust samples following live vaccination of layer flocks, *Vet. Microbiol.* (2019) 108393.
- [19] D. Sommer, et al., *Salmonella* detection in poultry samples, *Tierärztl. Prax. Ausg. G Großtiere Nutztiere* 40 (06) (2012) 383–389.
- [20] M. Myint, et al., The effect of pre-enrichment protocol on the sensitivity and specificity of PCR for detection of naturally contaminated *Salmonella* in raw poultry compared to conventional culture, *Food Microbiol.* 23 (6) (2006) 599–604.
- [21] H.M. Nam, et al., Application of SYBR green real-time PCR assay for specific detection of *Salmonella* spp. in dairy farm environmental samples, *Int. J. Food Microbiol.* 102 (2) (2005) 161–171.
- [22] I. Schrank, et al., Influence of enrichment media and application of a PCR based method to detect *Salmonella* in poultry industry products and clinical samples, *Vet. Microbiol.* 82 (1) (2001) 45–53.
- [23] B.M. Thomason, D.J. Dodd, W.B. Cherry, Increased recovery of salmonellae from environmental samples enriched with buffered peptone water, *Appl. Environ. Microbiol.* 34 (3) (1977) 270–273.
- [24] A. Oliveira, et al., Validating the Efficiency of a Simplex PCR and Quantitative SYBR Green qPCR for the Identification of *Salmonella* spp., *J. Food Microbiol. Saf. Hyg.* 3 (130) (2018) 2476–2059.1000130.
- [25] C. Techathuvanan, F.A. Draughon, D.H. D'Souza, Real-time reverse transcriptase PCR for the rapid and sensitive detection of *Salmonella typhimurium* from pork, *J. Food Prot.* 73 (3) (2010) 507–514.
- [26] R.W. Sweeney, R.H. Whitlock, S.C. McAdams, Comparison of three DNA preparation methods for real-time polymerase chain reaction confirmation of *Mycobacterium avium* subsp. *paratuberculosis* growth in an automated broth culture system, *J. Vet. Diagn. Invest.* 18 (6) (2006) 587–590.
- [27] X. Peng, et al., Comparison of direct boiling method with commercial kits for extracting fecal microbiome DNA by Illumina sequencing of 16S rRNA tags, *J. Microbiol. Methods* 95 (3) (2013) 455–462.
- [28] J. Yamagishi, et al., Comparison of boiling and robotics automation method in DNA extraction for metagenomic sequencing of human oral microbes, *PLoS One* 11 (4) (2016) e0154389.