

Methane and Nitrous Oxide Emissions from Cattle Excreta on an East African Grassland

D. E. Pelster,* B. Gisore, J. K. Koske, J. Goopy, D. Korir, M. C. Rufino, and K. Butterbach-Bahl

Abstract

Greenhouse gas (GHG) emission measurements from livestock excreta in Africa are limited. We measured CH₄ and N₂O emissions from excreta of six Boran (*Bos indicus*) and six Friesian (*Bos taurus*) steers near Nairobi, Kenya. The steers were fed one of three diets (T1 [chaffed wheat straw], T2 [T1 + *Calliandra calothyrsus* Meissner – 0.2% live weight per day], and T3 [T1 + calliandra – 0.4% live weight every 2 d]). The T1 diet is similar in quality to typical diets in the region. Calliandra is a leguminous fodder tree promoted as a feed supplement. Fresh feces and urine were applied to grasslands and emissions measured using static chambers. Cumulative 28-d fecal emissions were 302 ± 52.4 and 95 ± 13.8 mg CH₄-C kg⁻¹ dry matter for Friesian and Boran steers, respectively, and 11.5 ± 4.26 and 24.7 ± 8.32 mg N₂O-N kg⁻¹ dry matter for Friesian and Boran steers, respectively. For urine from Friesian steers, the N₂O emissions were 2.8 ± 0.64 mg N₂O-N 100 mL urine⁻¹. The CH₄ emission factors (EFs) (246 ± 49.5 and 87 ± 12.7 g CH₄-C yr⁻¹ animal⁻¹ for Friesian and Boran, respectively) were lower than the International Panel on Climate Change EFs (750 g CH₄-C animal⁻¹ yr⁻¹), whereas the N₂O EFs (0.1 and 0.2% for the Friesian and Boran feces, respectively, and 1.2% for urine) were also lower than International Panel on Climate Change estimates. The low N content of the excreta likely caused the low emissions and indicates that current models probably overestimate CH₄ and N₂O emissions from African livestock manure.

Core Ideas

- GHG emissions from African livestock excreta is lower than IPCC tier 1 emission factors.
- Low-quality feeds with low protein content result in low N content of excreta.
- Supplementation of cattle diet with calliandra reduced the methane emissions from cattle feces.
- The species of cattle causes differences in GHG emissions from feces.

INCREASED ATMOSPHERIC CONCENTRATIONS of greenhouse gases (GHG) such as carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O) over the last century are strongly correlated with global warming, and these gas concentrations continue to increase at an annual rate of approximately 0.4, 0.6, and 0.25%, respectively (Myhre et al., 2013). Although agricultural GHG sources account for approximately 12% of total anthropogenic emissions globally (Tubiello et al., 2015), in Africa GHG emissions from agriculture, forestry, and other land use accounts for approximately 61% of emissions (Valentini et al., 2014). Between 1990 and 2000, the agricultural emissions in Africa increased by about 99 MtCO₂e (18%) (Vergé et al., 2007) and continued to increase by about 2% per year between 2001 and 2011 (Tubiello et al., 2014).

In sub-Saharan Africa, livestock comprise a large proportion of total agricultural emissions, most of which is from enteric CH₄ production in ruminants (Valentini et al., 2014). However, between 7 and 15% of agricultural GHG emissions are associated with livestock manure (Smith et al., 2014; Tubiello et al., 2014). However, these emission rates from livestock manure in Africa are estimated using emission factors (EFs) from the International Panel on Climate Change (IPCC) that have been derived using measurements primarily from states within the Organization for Economic Cooperation and Development. These regions have livestock species, breeds, diets, management systems, and climatic conditions that often differ from those in tropical Africa (IPCC, 2006).

In tropical and subtropical agricultural production systems, the climate is generally warmer than temperate systems, which could result in greater N₂O and CH₄ emissions from excreta because emissions are often positively correlated with temperature (González-Avalos and Ruiz-Suárez, 2001; Rochette et al., 2014). However, the types of management systems used, the quality of the feeds, and the species of cattle raised may also affect emissions. The majority of African ruminants graze for much of their life (Schlecht et al., 2006), which results in over 40% of excreta

Copyright © American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America. 5585 Guilford Rd., Madison, WI 53711 USA. All rights reserved.

J. Environ. Qual. 45:1531–1539 (2016)

doi:10.2134/jeq2016.02.0050

This is an open access article distributed under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Received 10 Feb. 2016.

Accepted 18 Apr. 2016.

*Corresponding author (D.Pelster@cgiar.org).

D.E. Pelster, B. Gisore, J. Goopy, and D. Korir, International Livestock Research Institute, PO Box 30709, Nairobi, Kenya; J.K. Koske, School of Environmental Studies, Kenyatta Univ., PO Box 43844, Nairobi, Kenya; M.C. Rufino, Centre for International Forestry Research, PO Box 30677-00100, UN Avenue, Nairobi Kenya; K. Butterbach-Bahl, International Livestock Research Institute, PO Box 30709, Nairobi, Kenya and Karlsruhe Institute of Technology–Institute of Meteorology and Climate Research, Atmospheric Environmental Kreuzteckbahnstr. 19-82467 Garmisch-Partenkirchen, Germany. Assigned to Associate Editor Heidi Waldrip.

Abbreviations: DM, dry matter; GHG, greenhouse gas; IPCC, International Panel on Climate Change; LW, live weight.

being deposited on rangelands and pastures (Rufino et al., 2006), much more than in temperate regions. Thus, approximately 80% of the emissions associated with excreta deposited on rangelands and pasture occurs in developing countries (Smith et al., 2014).

The diet of African ruminants tends to be based on grasses and crop residues that are more fibrous than their counterparts from temperate regions, with lower digestibility and protein content (Schlecht et al., 2006). Dietary protein content, feed digestibility, and sugar content are known to influence the amounts and types of N and C voided in cattle excreta (Dijkstra et al., 2011; Merry et al., 2006; Rotz, 2004). Therefore, the lower-quality feeds likely result in excreta with reduced N concentrations and higher C/N ratios. To compensate for the low-protein diet, many agencies across eastern Africa promote the use of the legume fodder tree calliandra (*Calliandra calothyrsus* Meissner) as a feed supplement (Dawson et al., 2014). However, calliandra has high condensed tannin concentrations that cause higher recalcitrant N concentrations in cattle excreta (Delve et al., 2001). The increased C/N ratio and higher concentrations of recalcitrant N could result in lower N₂O emissions from the feces (Chantigny et al., 2013) compared with temperate systems. Also, native African cattle (*Bos indicus*) tend to have better water-scavenging abilities than their counterparts from temperate regions (*Bos taurus*), resulting in different fecal dry matter (DM) content (Quarterman et al., 1957). This physical difference in the fresh feces may also affect N₂O and CH₄ emissions from the excreta.

Agriculture in Kenya accounts for about 33% of the country's GHG emissions, and 90% of those emissions are from livestock (Stiebert, 2012). However, unlike temperate systems, much (roughly 60%) of the ruminant livestock in Kenya are kept by pastoralists in Kenya's arid and semiarid lands (Government of Kenya, 2010). Like most sub-Saharan countries, emission estimates in Kenya are estimated using the IPCC's Tier 1 approach. For development of proper emission reduction and mitigation strategies, actual measurements are necessary (Rosenstock et al., 2013; Rufino et al., 2014). Therefore, our objective was to estimate CH₄ and N₂O emissions from the excreta of Boran and Friesian steers. Boran cattle are a common native breed, originally from the Borana plateau (Felius, 1995), whereas the Friesian is a European breed that is commonly promoted in eastern Africa to increase dairy production. We hypothesized that (i) there would be no significant differences in emissions from feces and urine between the species, (ii) the protein supplementation with calliandra would increase N₂O emissions, and (iii) feeding patterns (i.e., oscillating protein supplementation; daily vs. bi-daily) of calliandra would not affect emissions.

Materials and Methods

The study was performed on grassland at the campus of the International Livestock Research Institute (1°16'14" S; 36°43'28" E) having a mixture of kikuyu grass (*Pennisetum clandestinum* Hochst. ex Chiov.) and Rhodes grass (*Chloris gayana* Kunth), with grazing being simulated by cutting the grass approximately every 14 d. The mean annual precipitation for the area is 980 mm, and the mean annual temperature is 17.5°C. The driest month is July (mean precipitation, 15 mm), and the wettest month is April (mean, 219 mm). The warmest month is March (mean temperature, 19.1°C), and the coolest month is

July (mean temperature, 15.1°C) (AmbiWeb GmbH, 2015). The soils are well drained, deep humic nitisols (Jaetzold et al., 2006), with clay-textured (16% sand, 65.6% clay) topsoils (0–20 cm), a pH of 7.0, and a bulk density of 0.8 g cm⁻³.

Excreta samples were obtained from Friesian ($n = 6$; live weight [LW], 183 ± 5.8 kg) and Boran ($n = 6$; LW, 126 ± 3.4 kg) yearling steers forming the experimental cohort for a feeding trial. Experimental design and sampling protocol has been described in detail elsewhere (Korir et al., 2015). Briefly, steers were assigned initially to one of three dietary treatments: T1, ad libitum wheat (*Triticum aestivum* L.) straw; T2, T1 + daily calliandra supplementation (0.2% of LW); and T3, T1 + bi-daily calliandra supplementation (0.4% LW). The basal diet was similar to feed typically given to east-African cattle during the dry season. Each steer received all treatments in a 3 × 3 × 2 factorial crossover design (3 treatments by 3 periods by 2 breeds). Animals were housed individually with free access to water. Each treatment period lasted 4 wk. For additional information on the feed, feces, and urine collection and analysis, see Korir et al. (2015). On the final evening of each period, the stalls were cleaned, and the urine collection buckets were emptied and rinsed to remove any residual acid. The following morning (~12 h later), the fresh urine and feces were collected for application.

Two replicates of each excreta sample (i.e., 1.0 kg of fresh manure and 234 mL of urine) were surface applied to grazing plots. The plots were delineated by plastic frames (35 × 25 cm) inserted 8 cm into the soil. Plots were kept 30 cm apart to prevent cross contamination. The amount of feces and urine was approximately equivalent to an average deposition event as measured during the previous 4-wk period (Korir et al., 2015). There were 56 frames in total (2 blocks, each with 12 feces samples, 12 urine samples, and 4 controls), with treatment randomly assigned to each plot within each block. The frames remained fixed in the soil for 2 wk before and 4 wk after urine and feces application. Two weeks after application, we simulated a 20-mm rainfall.

The manure and urine were applied on the same day as collection. Thus, there were three different application periods, with the application date corresponding to the final date of each feeding period in the feed trial. The application dates were as follows: Period 1, 30 June 2014; Period 2, 28 July 2014; and Period 3, 25 Aug. 2014. Precipitation was measured continuously using an ECRN-100 high-resolution rain gauge. Soil moisture content and temperature were measured at the time of gas sampling using a GS3 soil moisture and temperature sensor (Decagon GS3). The soil was analyzed for pH using a Jenway (Staffordshire) 350 pH meter with an epoxy-bodied combination electrode in a 1:5 soil/water solution. Soil bulk density was measured by dividing the oven-dried (12 h at 105°C) weight by the volume of the soil cores (5 cm diam., 20 cm length). Soil texture analysis was done at the Kenya Agricultural Research Institute using the hydrometer technique (Sheldrick and Wang, 1993).

Sample Analysis

Fecal DM was determined by drying in a forced air oven at 105°C. Organic matter content was calculated by subtracting the ash content after total combustion in muffle furnace from the DM. Total N concentration in feed and urine were determined by micro Kjeldhal procedure (see Korir et al. 2015 for more information). The fecal C content was calculated from the

known organic matter using the formula by Jiménez and Pérez García (1992) for compost. Applied C and N were calculated by multiplying the amount of fresh feces applied by the total C or N content (mg C kg FW^{-1} and mg N kg FW^{-1} , respectively). The amount of urine-N applied was calculated by multiplying the volume of urine by the concentration of total Kjeldahl N in each sample.

The CO_2 , N_2O , and CH_4 flux from the excreta and soil were measured using opaque polyvinyl chloride static chambers equipped with fans and vent tubes to allow continuous venting (Rochette and Bertrand, 2008). After the lids were clamped to the frames, a needle was inserted through the sampling port, and 60 mL of the headspace was sampled. The first 40 mL was used to flush out ambient air from an airtight 10-mL glass vial, after which the remaining 20 mL was injected to achieve overpressure of the vial to prevent contamination. Samples were taken immediately on closing of the chambers and again after 9, 18, and 27 min to ensure an adequate and linear response in headspace concentration. Once sampling was completed, the lids were removed. Sampling was done once a day for 2 d before application, once on the day of excreta application, daily for the after 4 d, and then two to three times per week for the subsequent 3 wk. On the application day, sampling occurred in the early afternoon, immediately after application; however, on all other days the sampling occurred in the morning, generally between 8:00 and 11:00 AM.

The gas samples from the chambers were analyzed within 1 wk for CO_2 , CH_4 , and N_2O on a SRI 8610C gas chromatograph fitted with a flame ionization detector for CO_2 (after passing through a methanizer) and CH_4 and an electron capture detector for N_2O . The carrier gas (pure N_2) had a flow rate of 20 mL min^{-1} . Two calibration gases (800 ppm CO_2 , 7040 ppb CH_4 , and 823 ppb N_2O and 400 ppm CO_2 , 4000 ppb CH_4 , and 360 ppb N_2O in synthetic air; Air Liquide) were analyzed alongside the samples, and the relation between the peak areas of the nearest calibration gas and its concentration was used to calculate the concentrations within the samples assuming a linear response in peak area to concentration.

Gas fluxes were estimated by calculating the rate of change in concentration over time using a linear approach while correcting for air pressure and temperature using the ideal gas law. Data were validated by examining the CO_2 concentrations. In cases where the change in concentration was below the precision of the gas chromatograph, we assumed no flux (i.e., minimum detection limits of $5 \mu\text{g CH}_4\text{-C m}^{-2} \text{ h}^{-1}$, $0.4 \text{ mg CO}_2\text{-C m}^{-2} \text{ h}^{-1}$, and $2 \mu\text{g N}_2\text{O-N m}^{-2} \text{ h}^{-1}$). The cumulative gas flux from the urine or feces application was estimated by summing the daily fluxes (using trapezoidal integration between sampling dates) and then subtracting the mean cumulative emissions from the control plots (i.e., those that received no application). The cumulative flux for the 2 d before excreta application was found to be similar for all plots, suggesting that differences between control plots and treatment plots were caused by the added excreta. Emission factors (the amount of CH_4 or N_2O emitted per unit of C/N added) for CH_4 and N_2O were calculated by dividing the amount of C or N lost as either $\text{CH}_4\text{-C}$ or $\text{N}_2\text{O-N}$, respectively, by the amount of C or N added.

Statistical Analysis

We used a random mixed model to analyze the feces and urine cumulative flux data (lmer package, R 3.0.3). All data were checked for normality using the Shapiro–Wilk test. We used ANOVA for CH_4 and CO_2 and Kruskal–Wallis for N_2O (to account for the non-normal distribution) to compare chamber effects for the preapplication period. For fecal flux analysis, species and diet were used as fixed effects, and period and the animal were fitted as random effects. For urine, only the Friesians' data were analyzed. The Boran data were not collected because the collection caused swelling of the animals' urethra, forcing us to halt the urine collection. Consequently, only diet was used as a fixed effect in the model, with period and animal remaining as random effects. We tested for relations between the amount of C or N applied with the amount of CH_4 , CO_2 , and N_2O emitted using a Pearson correlation. Annual emissions for a tropical livestock unit (one tropical livestock unit = one 250-kg steer) were calculated by multiplying the mean 1-mo emission (either $\text{mg CH}_4 28 \text{ d}^{-1}$ or $\text{mg N}_2\text{O} 28 \text{ d}^{-1}$) per kg manure by the amount of manure (kg) produced by each animal, adjusting for differences in LW and extrapolating to 1 yr.

For fecal DM, C, and N content, we used a two-factor ANOVA with species and diet as fixed factors. For the urine-N content, we used a single-factor ANOVA with diet as the single, fixed factor. Homoscedasticity was confirmed through visual observations of residual plots.

Results

The mean LW of the Friesian steers was approximately 50% greater than that of the Boran steers ($P < 0.001$) (Table 1). The Friesian steers also produced about 0.5 kg d^{-1} more feces than the Boran steers ($P < 0.001$). Steers receiving the bi-daily supplementation produced 1.0 kg more feces (Table 1) than steers on the basal diet ($P = 0.02$). The Boran steers' feces had a higher DM content than the Friesian steers' feces ($P < 0.001$) (Table 1), whereas the feces C content was similar among all treatments. Feces from animals fed the basal diet contained 29% less N than both the supplemented diets ($P < 0.001$) (Table 1). Because we applied the feces on a fresh weight basis, the application rates were 98.6 g C and 2.5 g N from the Friesian steer feces, compared with 118.7 g C and 3.1 g N with the Boran feces. Approximately 1 wk after application, we noticed an influx of termites in some of the plots, and by the end of the second week all of the feces plots had evidence of termites. The urine-N concentration and application rates were similar among the different diets (Table 2).

Cumulative CH_4 emissions from the urine plots were similar to control plots (i.e., plots with nothing added) ($P = 0.42$) (Table 3), indicating, as expected, that urine application did not affect CH_4 emissions. However, application of feces did cause greater CH_4 emissions compared with the controls ($P = 0.001$). The differences between control and application was mainly due to a short-term pulse of CH_4 immediately after application. During this pulse, the CH_4 emission rates increased from around 20 to over $600 \mu\text{g m}^{-2} \text{ h}^{-1}$ and then decreased exponentially to near background within about 1 wk (Fig. 1). Cumulative CH_4 fluxes (Table 3) differed by breed and diet, with the feces of the Friesian steers releasing more CH_4 than feces from the Boran steers ($P = 0.032$) and with both the diets with calliandra

supplementation resulting in higher feces CH₄ emissions than the basal diet ($P = 0.03$ and 0.01 for bi-daily and daily supplementation diets, respectively). The interaction between diet and breed had no detectable effect on the CH₄ emissions from the feces application ($P = 0.56$). The presence of termites within the chambers, which was generally noticed 1 to 2 wk after application, did not cause a noticeable increase in CH₄ emissions.

Cumulative 28-d CO₂ fluxes from the urine plots were approximately 34.0 g m⁻² greater than emissions from the control plots ($P = 0.002$), whereas feces application resulted in 42.5 g m⁻² more CO₂ than the control plots ($P = 0.02$) over the 28-d monitoring period (Table 3). Neither the diet nor the species affected CO₂ emissions from the feces and the urine applications ($P > 0.10$ for all). The CO₂ emissions from both the urine and feces applications increased immediately after application and then returned to baseline values after about 7 d (Fig. 1 and 2). Addition of water (or rainfall) increased emissions; however,

emissions in the control plots also increased to a similar degree (Fig. 1 and 2), suggesting that this increase was related to the wetting rather than the excreta.

Both the feces and urine application resulted in greater cumulative N₂O emissions over the 28-d period than the controls ($P = 0.007$ and 0.013 for feces and urine application, respectively) (Table 3). However, even though the amount of N applied as urine was approximately 25% of the amount of N applied as feces (Tables 1 and 2), the N₂O emissions from the urine were higher than emissions from feces, likely because the N in urine is more available than the N in feces. In general, the applications did not cause immediate increases in N₂O emissions; however, the addition of water 2 wk after the application caused a spike in emissions, particularly in Period 3 (Fig. 1 and 2). The N₂O emission rates were highest during the Period 3, when emission rates in the urine plots exceeded 250 μg m⁻² h⁻¹ and emissions from the feces plots exceeded 200 μg m⁻² h⁻¹. This spike in emissions

Table 1. Animal live weight, mean daily deposition, percent dry matter (DM), percent carbon (dry matter basis), total Kjeldahl nitrogen (dry matter basis) content, and C/N ratio of 1 kg of fresh cattle manure added to grassland at the campus of the International Livestock Research Institute, Nairobi, Kenya ($n = 34$).

| Type | Diet | Live weight kg | Deposition kg DM d ⁻¹ | DM % | Total C | | C/N ratio |
|----------|---|-------------------|-------------------------------------|---------------|-----------------------|---------------|--------------|
| | | | | | g kg ⁻¹ FW | | |
| Friesian | Basal diet (wheat straw only) | 181.5 ± 10.3† | 1.11 ± 0.03 | 20.6 ± 0.41a‡ | 97.6 ± 2.23a | 1.72 ± 0.026a | 56.9 ± 1.64a |
| | Daily supplement (calliandra) + basal diet | 185.5 ± 10.9 | 1.39 ± 0.05 | 20.3 ± 0.58a | 96.5 ± 2.41a | 2.56 ± 0.181b | 38.5 ± 2.33b |
| | Bi-daily supplement (calliandra) + basal diet | 181.7 ± 11.0 | 1.47 ± 0.05 | 20.4 ± 0.57a | 96.8 ± 2.79a | 2.46 ± 0.121b | 39.7 ± 2.15b |
| | Mean | 183.0 ± 5.8 | 1.31 ± 0.03 | 20.4 ± 0.28 | 97.0 ± 1.33 | 2.23 ± 0.118 | 45.3 ± 2.41 |
| Boran | Basal diet (wheat straw only) | 126.9 ± 5.4 | 0.76 ± 0.04 | 22.7 ± 1.73b | 107.2 ± 8.68b | 1.98 ± 0.106a | 53.8 ± 2.27a |
| | Daily supplement (calliandra) + basal diet | 121.8 ± 7.7 | 0.93 ± 0.07 | 25.7 ± 0.65b | 121.1 ± 2.41b | 3.00 ± 0.267b | 41.5 ± 3.39b |
| | Bi-daily supplement (calliandra) + basal diet | 128.4 ± 6.0 | 0.95 ± 0.02 | 25.9 ± 0.75b | 122.8 ± 2.79b | 3.08 ± 0.215b | 40.6 ± 2.20b |
| | Mean | 125.7 ± 3.5 | 0.88 ± 0.03 | 24.7 ± 0.75 | 116.8 ± 3.59 | 2.67 ± 0.167 | 45.5 ± 2.07 |

† Values are mean ± SEM.

‡ Different letters indicate significant differences within columns.

Table 2. Urine nitrogen concentration and amount of nitrogen applied from the application of 234 mL of fresh cattle urine to grassland at the campus of the International Livestock Research Institute, Nairobi, Kenya ($n = 15$).

| Type | Diet | Urine N concentration | | Urine N application rate | |
|----------|---|-----------------------|--|--------------------------|--|
| | | g L ⁻¹ | | mg N plot ⁻¹ | |
| Friesian | Basal diet (wheat straw only) | 2.59 ± 0.201† | | 605 ± 47.1 | |
| | Daily supplement (calliandra) + basal diet | 2.60 ± 0.136 | | 609 ± 31.9 | |
| | Bi-daily supplement (calliandra) + basal diet | 2.66 ± 0.173 | | 622 ± 40.5 | |

† Values are mean ± SEM.

Table 3. Cumulative CH₄, CO₂, and N₂O emissions over 28 d from the application of 1 kg of fresh cattle manure or 234 mL of fresh urine to grassland at the campus of the International Livestock Research Institute, Nairobi Kenya.

| Type | Diet | Emission† | | | | | |
|----------|---|--|---|--|---|--|---|
| | | Dung application | | | Urine application | | |
| | | CH ₄ -C | CO ₂ -C | N ₂ O-N | CH ₄ -C | CO ₂ -C | N ₂ O-N |
| | | mg CH ₄ -C kg manure ⁻¹ | g CO ₂ -C kg manure ⁻¹ | mg N ₂ O-N kg manure ⁻¹ | mg CH ₄ -C 234 mL urine ⁻¹ | g CO ₂ -C 234 mL urine ⁻¹ | mg N ₂ O-N 234 mL urine ⁻¹ |
| Friesian | basal diet (wheat straw only) | 34.4 ± 9.7ab‡ | 40.5 ± 5.3 | 1.92 ± 1.19 | -10.4 ± 4.29 | 36.2 ± 6.18 | 8.4 ± 2.09 |
| | daily supplement (calliandra) + basal diet | 75.2 ± 15.5c | 42.7 ± 5.4 | 4.01 ± 1.85 | -5.5 ± 4.24 | 35.5 ± 7.25 | 7.3 ± 2.98 |
| | bi-daily supplement (calliandra) + basal diet | 71.9 ± 24.7c | 31.3 ± 9.4 | 0.93 ± 1.18 | -6.3 ± 6.39 | 30.3 ± 12.32 | 4.0 ± 1.18 |
| Boran | basal diet (wheat straw only) | 11.6 ± 3.9a | 43.1 ± 7.0 | 7.04 ± 4.55 | ND§ | ND | ND |
| | daily supplement (calliandra) + basal diet | 24.6 ± 3.8b | 46.4 ± 8.2 | 7.82 ± 2.74 | ND | ND | ND |
| | bi-daily supplement (calliandra) + basal diet | 34.7 ± 7.8b | 53.0 ± 8.5 | 2.94 ± 1.11 | ND | ND | ND |

† Emissions shown were calculated by subtracting the emissions from the control (no addition) plots from the total emissions from plots with the added feces; mean cumulative (28 d) emissions from control plots: 12.0 ± 5.54 mg CH₄-C m⁻², 99.0 ± 6.28 g CO₂-C m⁻², and 5.1 ± 1.56 mg N₂O-N m⁻² ($n = 34$).

‡ Values are mean ± SEM. Different letters indicate significant differences within columns.

§ ND, not determined.




Fig. 1. Mean CO_2 ($\text{mg CO}_2\text{-C m}^{-2} \text{h}^{-1}$), CH_4 ($\mu\text{g CH}_4\text{-C m}^{-2} \text{h}^{-1}$), and N_2O ($\mu\text{g N}_2\text{O-N m}^{-2} \text{h}^{-1}$) flux rates from 1 kg fresh feces applications to grassland, along with soil moisture (0.05 m depth) and precipitation. Blue arrows indicate application of excreta, black arrows indicate water application, and blue dashed lines differentiate three application periods. The different treatments are as follows: basal, basal diet of chaffed wheat straw; bi-daily, basal diet + bi-daily supplement of calliandra (2% live weight); Borana, feces from a borana steer (*Bos indicus*); daily, basal diet + daily supplement of calliandra (1% LW); Friesian, feces from a Friesian steer (*Bos taurus*); none, no feces application.

coincided with a rain event that deposited over 22 mm of rain over 3 d and a corresponding increase in soil moisture from 0.20 to 0.26% v/v (Fig. 2). For the feces application, the N_2O emissions were greater from the Boran than from the Friesian steers ($P = 0.09$) (Table 3), whereas diet had no effect on N_2O emissions ($P = 0.19$). The urine application increased N_2O emissions compared with the controls; however, diet had no detectable effect on emissions ($P = 0.37$). The interaction between diet and breed had no detectable effect on N_2O emissions from the feces applications ($P = 0.77$).

There were no correlations between the C applied and CO_2 emissions ($P = 0.34$) or between the N applied and N_2O emissions ($P = 0.86$). However, there was a significant negative correlation between the C applied and cumulative CH_4 emissions ($P = 0.008$; $R = -0.32$). However, the range of applied C and N was very narrow (see Table 1), and the correlations could change if a wider range is used.

Discussion

The amount of feces voided is determined by feed intake and digestibility. In turn, feed intake is related mainly to LW (CSIRO, 2007). Although all the steers were yearlings, the Friesian steers were heavier than the Borans (Table 1), which was reflected in differences in feed intake and the amount of feces

voided. Breeds of *Bos indicus* adapted to harsh climatic conditions, such as Borans, have higher water reabsorption efficiencies compared with *Bos taurus* breeds like Friesians (Quarterman et al., 1957), which likely caused the higher DM content in the Boran feces. Because we applied manure on a fresh-weight basis, the higher DM content in the Boran feces resulted in more C being added to those plots. However, even with the greater amount of C applied, the CH_4 emissions from the Boran feces were less (per kg of fresh feces) than the emissions from the Friesian feces. Because most of the CH_4 was produced very rapidly after application, it is likely that the higher moisture content of the Friesian feces formed a more favorable environment for CH_4 production (Lodman et al., 1993) for a longer period of time, allowing for additional methanogenesis (Conrad, 1996).

With the influx of termites, we expected to see a corresponding increase in CH_4 emissions. However, this did not occur, suggesting that the termites we saw produced little to no CH_4 or that they transferred the feces to another location, producing CH_4 somewhere else. Further study to properly evaluate the role of termites in fecal CH_4 emissions is required.

The greater amounts of feces deposited by the Friesians (Table 1) would produce cumulative CH_4 fluxes that were approximately five times greater per steer than cumulative CH_4 fluxes from the feces of a Boran steer. Even when accounting for




Fig. 2. Mean CO_2 ($\text{mg CO}_2\text{-C m}^{-2} \text{h}^{-1}$), CH_4 ($\mu\text{g CH}_4\text{-C m}^{-2} \text{h}^{-1}$), and N_2O ($\mu\text{g N}_2\text{O-N m}^{-2} \text{h}^{-1}$) flux rates from 234-mL urine applications to grassland along with soil moisture and precipitation. Blue arrows indicate application of urine, black arrows indicate water application, and blue dashed lines differentiate the three periods. The different treatments are as follows: basal, basal diet of chaffed wheat straw; bi-daily, basal diet + bi-daily supplement of calliandra (2% live weight); daily, basal diet + daily supplement of calliandra (1% live weight); Friesian, urine from a Friesian steer (*Bos taurus*); none, no feces application.

differences in LW, fecal CH_4 emissions (per kg LW) from the Friesian steers were still 3.5 times higher than from the Boran steers. The different water contents of the feces therefore appear to result in different emission potentials as well. Although it is known that feces properties differ between the two species, to our knowledge this is the first study to report differences in CH_4 emissions between feces from *Bos indicus* and *Bos taurus*.

The measured CH_4 emissions, which ranged from 11.6 to 75.2 $\text{mg CH}_4\text{-C kg feces}^{-1}$ over a 28-d period (Table 3), were much lower than those in a previous study (Jarvis et al., 1995), which found that CH_4 losses from 1 kg of feces in the United Kingdom ranged from 300 to 2040 $\text{mg CH}_4\text{-C kg feces}^{-1}$ over a 10- to 15-d period. However, CH_4 losses measured in our study were much more consistent with two other UK studies that measured CH_4 losses of between 20 and 90 $\text{mg CH}_4\text{-C kg feces}^{-1}$ over study periods of between 15 and 60 d (Holter, 1997; Yamulki et al., 1999) as well as a study in Brazil that measured emission rates between 10 and 60 $\text{mg CH}_4\text{-C kg feces}^{-1}$ over 30 d (Mazzetto et al., 2014).

Dietary composition is known to affect the N concentration, soluble organic C, readily fermentable carbohydrates (Boadi et al., 2004), and C/N ratio of feces, which consequently affects CH_4 emissions (Cardenas et al., 2007). The steers used in this study were fed a basal diet with a low crude protein content (2%),

which is known to increase N use efficiency, reduce the excreta N concentrations, and increase the excreta C/N ratio (Korir et al., 2015). The C/N ratio of the feces was much higher than previous studies (e.g., Jarvis et al., 1995; Qian and Schoenau, 2002; Yamulki et al., 1999) where the ratio typically ranged between 10 and 15 and is likely why these emissions were so low when compared with those from Jarvis et al. (1995). However, these results were consistent with Mazzetto et al. (2014), who also used low-quality tropical fodders. Fecal CH_4 emissions were found to be negatively correlated to the C/N ratio of the feces (Table 1), consistent with previous studies (Jarvis et al., 1995). The C/N ratios in our study ranged from 38 to 54, all of which were greater than the C/N ratios (13–28) found in Jarvis et al. (1995), indicating that the effect of C/N ratio on CH_4 emissions extends across a much wider range than previously thought.

Assuming that emission rates remain the same throughout the rest of the year and that the deposition rates and feces properties remain consistent throughout the year, the annual fecal CH_4 emissions per tropical livestock unit would be approximately $117 \pm 17.0 \text{ g CH}_4 \text{ yr}^{-1} \text{ animal}^{-1}$ for the Boran and $328 \pm 68.1 \text{ g CH}_4 \text{ yr}^{-1} \text{ animal}^{-1}$ for the Friesian breeds (after adjustment for differences in LW). These results are between 9 and 25% of what the IPCC uses as the CH_4 emission factor for feces from African cattle (IPCC, 2006). However, emission

rates, deposition rates, and feces properties are likely to change during the different seasons, and it is important to measure the effects of these seasonal changes as well. Even though the increased temperature typical of the tropics should increase emissions compared with temperate regions, it appears that the effect of the lower-quality fodder and consequently the higher C/N ratio of the feces has a stronger effect on controlling emissions in these systems.

Because the feed trial took place during the dry season, we were limited in our supply of fresh manure and were only able to measure emissions during the dry season. We did add 20 mm of precipitation 2 wk after application to mimic a rain event, which resulted in an increase in emission rates. This, along with changes to fodder quality and therefore manure quality as well, suggests that the emission factors may change if measurements are made throughout the year rather than during just one season. A previous study found higher emissions during the “summer,” when temperatures were higher than during a cooler “winter” period in Brazil (Mazzetto et al., 2014). Also, deposition rates and feces properties are not constant throughout the year (Rufino et al., 2006; Schlecht et al., 2006). Given that the ad libitum diet with additional protein supplementation was a better quality diet than many African cattle receive during the dry season, it is likely that annual field emissions are lower than what we suggest here, although additional investigation is required to verify these assumptions. Excreta emissions on pasturelands, however, are typically very low in comparison to enteric CH_4 emissions (Jarvis et al., 1995; Mazzetto et al., 2014), although they are still important for emission reporting (e.g., IPCC, 2006; Tubiello et al., 2015).

The increase in CO_2 from the feces application was expected because we were adding approximately between 97 and 117 g C to the plots. However, there was very little C applied with the urine application, which also resulted in a similar increase in CO_2 emissions. The addition of the urea-N likely caused increased soil organic matter decomposition, also known as priming (Kuzyakov et al., 2000), which resulted in the large increase in CO_2 emissions.

Urine application increases N_2O emissions compared with control plots (Table 3) likely because the greater N availability provided additional substrate for denitrification. The lack of rain during the first two periods may have limited denitrification; however, the precipitation at the start of the third period likely mobilized additional C and N substrate for denitrification (Birch, 1960; Ruser et al., 2006) while providing anaerobic sites. These peaks in N_2O emissions tend to be the result of denitrification, which requires the correct air–water balance, and the degree of gas diffusivity (Balaine et al., 2013; Butterbach-Bahl et al., 2013; Singurindy et al., 2009). Period 3 emissions were twice as high as Period 1 emissions and at least eight times higher than Period 2 emissions (Fig. 2). Because the amounts of applied N applied were almost equal, it seems that the greater precipitation received in period 3 (45.6 mm) compared with Period 1 (11.6 mm) and Period 2 (6.4 mm; see Fig. 2) caused the higher flux in Period 3. The burst of N_2O after a precipitation event and urine/feces additions to grassland has been observed in other trials as well (Boon et al., 2014; de Klein et al., 2003; Yamulki et al., 1999).

Opposite to the CH_4 flux, the feces of Borans produced more N_2O than that of the Friesians. It is possible that the

higher moisture content of the Friesian feces favored further reduction of N_2O to N_2 . Calliandra addition to the diet was expected to increase both fecal N concentration and N_2O emissions. However, although supplementation resulted in greater N excreted in the feces, it did not cause measurable differences in N_2O emissions. However, we measured only total C and N, whereas it is the labile portion of both that are substrates for denitrification. The high tannin content of calliandra (3.2% for this study) (Korir et al., 2015) has been found to result in greater concentrations of recalcitrant N in the feces (Delve et al., 2001) that may have limited N availability for denitrifiers.

The higher N_2O emissions from urine compared with feces were likely because the N in urine is more easily available (Sordi et al., 2014). This is also suggested by the delay in N_2O emissions from the feces application after the simulated rainfall (Fig. 1 and 2), which did not occur in the urine applications. It is likely that the fecal N needed to be mineralized before denitrification could occur. In addition, the high amounts of C in the feces and the high C/N ratio likely caused rapid N immobilization, resulting in less available NO_3^- and further reduction of N_2O to N_2 (Blackmer and Bremner, 1978; Senbayram et al., 2012).

The EFs for the feces in this study (0.2% from the Boran and 0.1% from Friesians) and from the urine (1.2% from the Friesian only) were lower than the 2% estimated by the IPCC (2006), although they were similar to a Brazilian study that also measured low N_2O emissions from cattle excreta on tropical rangelands (Mazzetto et al., 2015). Because fecal C/N ratios were found to be negatively correlated with N mineralization (Qian and Schoenau, 2002) and N_2O emissions (Chantigny et al., 2013), it is likely that the high C/N ratio of the feces in this tropical system was responsible for the low emissions.

However, because our measurements took place during the dry season, it is possible that emissions during the rainy season may be higher than what was measured here. Although it is likely that the increased water availability during the rainy season may cause increased GHG emissions, the improvement in fodder and therefore manure quality (Rufino et al., 2006; Schlecht et al., 2006) will likely have a larger effect on emission rates. With the exception of Period 3, there was only a small increase in N_2O emissions with a simulated 20-mm rainfall event, suggesting that the lack of rainfall alone may not be causing the low EF. Rather, the low N_2O emissions are more likely related to the low N concentration and high C/N ratio typical of tropical cattle manure (Rufino et al., 2006), which was related to the poor-quality fodder (Korir et al., 2015).

Conclusions

As indicated, the IPCC (2006) EF are at least two times higher for fecal CH_4 , 10 to 20 times higher for fecal N_2O , and about two times higher for urine N_2O . The low EFs in this study were likely due to the poor-quality diet (low crude protein) and the subsequent low excreta N. Also, the native cattle breeds (i.e., *Bos indicus*) have better water retention than imported breeds, which caused decreased CH_4 but increased N_2O emissions. The diets used in this study were consistent with those frequently used in smallholder farms in the region and similar in digestible energy to the low-quality fodder category used by the IPCC to estimate livestock emissions, suggesting that emission factors

used for GHG inventories in this region may need to be revised. However, additional studies performed under different climatic seasons, linked with measurements of enteric fermentation and with measurements performed over extended periods, are required to verify these findings.

Acknowledgments

The authors thank the CGIAR Research Program on Climate Change, Agriculture, and Food Security (CCAFS) and its Standard Assessment of Mitigation Potential and Livelihoods in Smallholder Systems (SAMPLES) programme as well as the Deutsche Gesellschaft für Internationale Zusammenarbeit (GIZ) project on "In situ assessment of GHG emissions from two livestock systems in East Africa" Project No. 81180349 for technical support and financial support of scientists and laboratories, and Marko Kvakic, Suzanne Jacobs, and George Wanyama for help collecting and analyzing the field samples.

References

- AmbiWeb GmbH. 2015. Climate data. Gernsbach, Germany. <http://en.climate-data.org/info/sources/> (accessed 18 Apr. 2015).
- Balaine, N., T.J. Clough, M.H. Beare, S.M. Thomas, E.D. Meenken, and J.G. Ross. 2013. Changes in relative gas diffusivity explain soil nitrous oxide flux dynamics. *Soil Sci. Soc. Am. J.* 77:1496–1505. doi:10.2136/sssaj2013.04.0141
- Birch, H.F. 1960. Nitrification in soils after different periods of dryness. *Plant Soil* 12:81–96. doi:10.1007/BF01377763
- Blackmer, A.M., and J.M. Bremner. 1978. Inhibitory effect of nitrate on reduction of N_2O to N_2 by soil microorganisms. *Soil Biol. Biochem.* 10:187–191. doi:10.1016/0038-0717(78)90095-0
- Boadi, D.A., K.M. Wittenberg, S.L. Scott, D. Burton, K. Buckley, J.A. Small, et al. 2004. Effect of low and high forage diet on enteric and manure pack greenhouse gas emissions from a feedlot. *Can. J. Anim. Sci.* 84:445–453. doi:10.4141/A03-079
- Boon, A., J.S. Robinson, D.R. Chadwick, and L.M. Cardenas. 2014. Effect of cattle urine addition on the surface emissions and subsurface concentrations of greenhouse gases in a UK peat grassland. *Agric. Ecosyst. Environ.* 186:23–32. doi:10.1016/j.agee.2014.01.008
- Butterbach-Bahl, K., E.M. Baggs, M. Dannenmann, R. Kiese, and S. Zechmeister-Boltenstern. 2013. Nitrous oxide emissions from soils: How well do we understand the processes and their controls? *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 368:20130122. doi:10.1098/rstb.2013.0122
- Cardenas, L.M., D. Chadwick, D. Scholefield, R. Fychan, C.L. Marley, R. Jones, et al. 2007. The effect of diet manipulation on nitrous oxide and methane emissions from manure application to incubated grassland soils. *Atmos. Environ.* 41:7096–7107. doi:10.1016/j.atmosenv.2007.04.055
- Chantigny, M.H., D.E. Pelster, M.-H. Perron, P. Rochette, D.A. Angers, L.-É. Parent, et al. 2013. Nitrous oxide emissions from clayey soils amended with paper sludges and biosolids of separated pig slurry. *J. Environ. Qual.* 42:30–39. doi:10.2134/jeq2012.0196
- Conrad, R. 1996. Soil microorganisms as controllers of atmospheric trace gases (H_2 , CO , CH_4 , OCS , N_2O , and NO). *Microbiol. Rev.* 60:609–640.
- CSIRO. 2007. Nutrient requirements of domesticated ruminants. CSIRO, Collingwood, Victoria, Australia.
- Dawson, I.K., S. Carsan, S. Franzel, R. Kindt, P. van Breugel, L. Graudal, et al. 2014. Agroforestry, livestock, fodder production and climate change adaptation and mitigation in East Africa: Issues and options. World Agroforestry Center, Nairobi, Kenya.
- de Klein, C.A.M., L. Barton, R.R. Sherlock, Z. Li, and R.P. Littlejohn. 2003. Estimating a nitrous oxide emission factor for animal urine from some New Zealand pastoral soils. *Soil Res.* 41:381–399. doi:10.1071/SR02128
- Delve, R.J., G. Cadisch, J.C. Tanner, W. Thorpe, P.J. Thorne, and K.E. Giller. 2001. Implications of livestock feeding management on soil fertility in the smallholder farming systems of sub-Saharan Africa. *Agric. Ecosyst. Environ.* 84:227–243. doi:10.1016/S0167-8809(00)00244-9
- Dijkstra, J., O. Oenema, and A. Bannink. 2011. Dietary strategies to reducing N excretion from cattle: Implications for methane emissions. *Curr. Opin. Environ. Sustain.* 3:414–422. doi:10.1016/j.cosust.2011.07.008
- Felius, M. 1995. Cattle breeds: An encyclopedia. Misset, Doetinchem, The Netherlands.
- González-Avalos, E., and L.G. Ruiz-Suárez. 2001. Methane emission factors from cattle manure in Mexico. *Bioresour. Technol.* 80:63–71. doi:10.1016/S0960-8524(01)00052-9
- Government of Kenya. 2010. Strategic plan: 2008–2012. Ministry of Livestock Development, Nairobi, Kenya.
- Holter, P. 1997. Methane emissions from Danish cattle dung pats in the field. *Soil Biol. Biochem.* 29:31–37. doi:10.1016/S0038-0717(96)00267-2
- Iglesias Jiménez, E., and V. Pérez García. 1992. Relationships between organic carbon and total organic matter in municipal solid wastes and city refuse composts. *Bioresour. Technol.* 41:265–272. doi:10.1016/0960-8524(92)90012-M
- IPCC. 2006. 2006 IPCC guidelines for national greenhouse gas inventories. IPCC, Geneva, Switzerland.
- Jaetzold, R., H. Schmidt, B. Hornetz, and C. Shisanya. 2006. Farm management handbook of Kenya. Vol. II. Natural conditions and farm management information. Ministry of Agriculture, Kenya, in Cooperation with the German Agency for Technical Cooperation (GTZ), Nairobi, Kenya.
- Jarvis, S.C., R.D. Lovell, and R. Panayides. 1995. Patterns of methane emission from excreta of grazing animals. *Soil Biol. Biochem.* 27:1581–1588. doi:10.1016/0038-0717(95)00092-5
- Korir, D., J. Goopy, C. Gachui, and K. Butterbach-Bahl. 2015. Supplementation with *Calliandra calothyrsus* improves N retention in cattle fed low protein diets. *Anim. Prod. Sci.* 5:619. doi:10.1071/AN15569
- Kuzyakov, Y., J.K. Friedel, and K. Stahr. 2000. Review of mechanisms and quantification of priming effects. *Soil Biol. Biochem.* 32:1485–1498. doi:10.1016/S0038-0717(00)00084-5
- Lodman, D.W., M.E. Branine, B.R. Carmean, P. Zimmerman, G.M. Ward, and D.E. Johnson. 1993. Estimates of methane emissions from manure of U.S. cattle. *Chemosphere* 26:189–199. doi:10.1016/0045-6535(93)90421-Z
- Mazzetto, A.M., A.S. Barneze, B.J. Feigl, J.W. Van Groenigen, O. Oenema, and C.C. Cerri. 2014. Temperature and moisture affect methane and nitrous oxide emission from bovine manure patches in tropical conditions. *Soil Biol. Biochem.* 76:242–248. doi:10.1016/j.soilbio.2014.05.026
- Mazzetto, A.M., A.S. Barneze, B.J. Feigl, J.W. Van Groenigen, O. Oenema, C.A.M. De Klein, et al. 2015. Use of the nitrification inhibitor dicyandiamide (DCD) does not mitigate N_2O emission from bovine urine patches under Oxisol in Northwest Brazil. *Nutr. Cycling Agroecosyst.* 101:83–92. doi:10.1007/s10705-014-9663-4
- Merry, R.J., M.R.F. Lee, D.R. Davies, R.J. Dewhurst, J.M. Moorby, N.D. Scollan, et al. 2006. Effects of high-sugar ryegrass silage and mixtures with red clover silage on ruminant digestion: 1. In vitro and in vivo studies of nitrogen utilization. *J. Anim. Sci.* 84:3049–3060. doi:10.2527/jas.2005-735
- Myhre, G., D. Shindell, F.-M. Breion, W. Collins, J. Fuglestvedt, J. Huang, et al. 2013. Anthropogenic and natural radiative forcing. In: T.F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, et al., editors, *Climate change 2013: The physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge Univ. Press, Cambridge, UK, and New York. p. 659–740.
- Qian, P., and J.J. Schoenau. 2002. Availability of nitrogen in solid manure amendments with different C:N ratios. *Can. J. Soil Sci.* 82:219–225. doi:10.4141/S01-018
- Quarterman, J., G.D. Phillips, and G.H. Lampkin. 1957. A difference in the physiology of the large intestine between European and indigenous cattle in the tropics. *Nature* 180:552–553. doi:10.1038/180552a0
- Rochette, P., and N. Bertrand. 2008. Soil-surface gas emissions. In: M. Carter and E.G. Gregorich, editors, *Soil sampling and methods of analysis*. CRC Press, Boca Raton, FL. p. 851–861.
- Rochette, P., M.H. Chantigny, N. Ziadi, D.A. Angers, G. Bélanger, É. Charbonneau, et al. 2014. Soil nitrous oxide emissions after deposition of dairy cow excreta in eastern Canada. *J. Environ. Qual.* 43:829–841. doi:10.2134/jeq2013.11.0474
- Rosenstock, T.S., M.C. Rufino, K. Butterbach-Bahl, and E. Wollenberg. 2013. Toward a protocol for quantifying the greenhouse gas balance and identifying mitigation options in smallholder farming systems. *Environ. Res. Lett.* 8:021003. doi:10.1088/1748-9326/8/2/021003
- Rotz, C.A. 2004. Management to reduce nitrogen losses in animal production. *J. Anim. Sci.* 82(E-Suppl):E119–E137.
- Rufino, M.C., P. Brandt, M. Herrero, and K. Butterbach-Bahl. 2014. Reducing uncertainty in nitrogen budgets for African livestock systems. *Environ. Res. Lett.* 9:105008. doi:10.1088/1748-9326/9/10/105008
- Rufino, M.C., E.C. Rowe, R.J. Delve, and K.E. Giller. 2006. Nitrogen cycling efficiencies through resource-poor African crop–livestock systems. *Agric. Ecosyst. Environ.* 112:261–282. doi:10.1016/j.agee.2005.08.028
- Ruser, R., H. Flessa, R. Russow, G. Schmidt, F. Buegger, and J.C. Munch. 2006. Emission of N_2O , N_2 and CO_2 from soil fertilized with nitrate: Effect of compaction, soil moisture and rewetting. *Soil Biol. Biochem.* 38:263–274. doi:10.1016/j.soilbio.2005.05.005

- Schlecht, E., P. Hiernaux, I. Kadaouré, C. Hülsebusch, and F. Mahler. 2006. A spatio-temporal analysis of forage availability and grazing and excretion behaviour of herded and free grazing cattle, sheep and goats in Western Niger. *Agric. Ecosyst. Environ.* 113:226–242 doi:10.1016/j.agee.2005.09.008.
- Senbayram, M., R. Chen, A. Budai, L. Bakken, and K. Dittert. 2012. N₂O emission and the N₂O/(N₂O+N₂) product ratio of denitrification as controlled by available carbon substrates and nitrate concentrations. *Agric. Ecosyst. Environ.* 147:4–12. doi:10.1016/j.agee.2011.06.022
- Sheldrick, B., and C. Wang. 1993. Particle size distribution. In: M.R. Carter, editor, *Soil sampling and methods of analysis*. Lewis Publishers, Ann Arbor, MI, p. 499–511.
- Singurindy, O., M. Molodovskaya, B.K. Richards, and T.S. Steenhuis. 2009. Nitrous oxide emission at low temperatures from manure-amended soils under corn (*Zea mays* L.). *Agric. Ecosyst. Environ.* 132:74–81 doi:10.1016/j.agee.2009.03.001.
- Smith, P., M. Bustamante, H. Ahammad, H. Clark, H. Dong, E.A. Elsidig, et al. 2014. Agriculture, Forestry and Other Land Use (AFOLU). In: O. Edenhofer, R. Pichs-Madruga, Y. Sokona, E. Farahani, S. Kadner, K. Seyboth, et al., editors, *Mitigation of climate change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge Univ. Press, Cambridge, UK, and New York.
- Sordi, A., J. Dieckow, C. Bayer, M.A. Albuquerque, J.T. Piva, J.A. Zanatta, et al. 2014. Nitrous oxide emission factors for urine and dung patches in a subtropical Brazilian pastureland. *Agric. Ecosyst. Environ.* 190:94–103. doi:10.1016/j.agee.2013.09.004
- Stieber, S., D. Murphy, J. Dion, M. Iisd, and D. Sawyer. 2012. Kenya's climate change action plan: Mitigation. Chapter 3: Agriculture. Ministry of Environment and Natural Resources, Government of Kenya, Nairobi, Kenya.
- Tubiello, F.N., M. Salvatore, R. Condor, A. Ferrara, S. Rossi, S. Federici, et al. 2014. Agriculture, forestry and other land use emissions by sources and removals by sinks 1990-2011 Analysis. FAO Statistics Division Working Paper. FAO, Rome, Italy.
- Tubiello, F.N., M. Salvatore, A.F. Ferrara, J. House, S. Federici, S. Rossi, et al. 2015. The contribution of agriculture, forestry and other land use activities to global warming, 1990–2012. *Glob. Change Biol.* 21:2655–2660. doi:10.1111/gcb.12865
- Valentini, R., A. Arneeth, A. Bombelli, S. Castaldi, R. Cazzolla Gatti, F. Chevalier, et al. 2014. A full greenhouse gases budget of Africa: Synthesis, uncertainties, and vulnerabilities. *Biogeosciences* 11:381–407. doi:10.5194/bg-11-381-2014
- Vergé, X.P.C., C. De Kimpe, and R.L. Desjardins. 2007. Agricultural production, greenhouse gas emissions and mitigation potential. *Agric. For. Meteorol.* 142:255–269. doi:10.1016/j.agrformet.2006.06.011
- Yamulki, S., S.C. Jarvis, and P. Owen. 1999. Methane emission and uptake from soils as influenced by excreta deposition from grazing animals. *J. Environ. Qual.* 28:676–682. doi:10.2134/jeq1999.00472425002800020036x