

Bibliography

- Alberda, Th. (1957). The effects of cutting, light intensity and night temperature on growth and soluble carbohydrate content of *Lolium perenne* L. *Plant and Soil* **8**, 199-230.
- Alberda, Th. (1965). The influence of temperature, light intensity and nitrate concentration on dry matter production and chemical composition of *Lolium perenne* L. *Netherlands Journal of Agricultural Science* **13**, 335-60.
- Alberda, Th. (1966a). Responses of grasses to temperature and light. In 'The Growth of Cereals and Grasses.' (Eds. F.L. Milthorpe and J.D. Ivins.) pp. 200-12. (Butterworths: London.)
- Alberda, Th. (1966b). The influence of reserve substances on dry matter production after defoliation. *Proceedings of the 10th International Grassland Congress, Helsinki*, 140-7.
- Anon. (1985). Seed Science and Technology. *Proceedings of the International Seed Testing Association* **13**, 322-463.
- Anon. (1992a). Advisory Survey, New South Wales Department of Agriculture.
- Anon. (1992b). 'TableCurve 2D Automated Curve Fitting and Equation Discovery.' (Jandel Scientific Software: San Rafael.)
- Appadurai, R.R. and Holmes, W. (1964). The influence of stage of growth, closeness of defoliation, and moisture on the growth and productivity of a ryegrass-white clover sward. 1. Effect on herbage yields. *Journal of Agricultural Science* **62**, 327-32.
- Archer, K.A. and Robinson, G.G. (1989). The role of stolons and seedlings in the persistence and production of white clover (*Trifolium repens* L. cv. Huia) in temperate pastures on the Northern Tablelands, New South Wales. *Australian Journal of Agricultural Research* **40**, 605-16.
- Archer, K.A. and Rochester, I.J. (1982). Numbers and germination characteristics of white clover seed recovered from soils on the northern tablelands of New South Wales after drought. *Journal of the Australian Institute of Agricultural Science* **1982**, 99-101.
- Arcioni, S., Damiani, F. and Mariott, D. (1985a). Relationships between some

- physiological and morphological traits in the first phases of plant growth in two differently adapted varieties of *Lolium perenne*. *Genetica-Agraria* **39**, 306.
- Arcioni, S., Damiani, F. and Mariotti, D. (1985b). Ecological adaptation in *Lolium*: morphological and physiological parameters at early stages of growth in two varieties with different degree of persistence. *Agronomie* **5**, 181-6.
- Arcioni, S., Falcinelli, M. and Mariotti, D. (1985c). Ecological adaptation in *Lolium perenne* L.: physiological relationships among persistence, carbohydrate reserves and water availability. *Canadian Journal of Plant Science* **65**, 615-24.
- Arcioni, S., Mariotti, D., Falcinelli, M. and Ceccarelli, S. (1980). Relationship between ecological adaptation and carbohydrate reserve evolution for two different varieties of *Lolium perenne* L. *Annals Amelioration Plantes* **30**, 309-19.
- Auda, H., Blaser, R.E. and Brown, R.H. (1966). Tillering and carbohydrate contents of orchardgrass as influenced by environmental factors. *Crop Science* **6**, 139-43.
- Awad, A.S. and Edwards, D.G. (1977). Reversal of adverse effects of heavy ammonium sulphate application on growth and nutrient status of a kikuyu pasture. *Plant and Soil* **48**, 169-83.
- Baker, B.S. and Jung, G.A. (1968a). Effect of environmental conditions on the growth of four perennial grasses. I. Response to controlled temperature. *Agronomy Journal* **60**, 155-8.
- Baker, B.S. and Jung, G.A. (1968b). Effect of environmental conditions on the growth of four perennial grasses. II. Response to fertility, water and temperature. *Agronomy Journal* **60**, 158-62.
- Baker, B.S. and Jung, G.A. (1970). Response of four perennial grasses to high temperature stress. *Proceedings of the 11th International Grassland Congress, Surfers Paradise*, 499-502.
- Balasko, J.A., Evers, G.W. and Duell, R.W. (1995). Bluegrasses, ryegrasses and bentgrasses. In 'Forages, Volume 1: An Introduction to Grassland Agriculture.' Fifth Edition. (Eds. R.F. Barnes, D.A. Miller and C.J. Nelson.) pp. 357-72. (Iowa State University Press: Iowa.)
- Barker, D.J., Chu, A.C.P. and Korte, C.J. (1985). Some effects of spring defoliation and drought on perennial ryegrass swards. *Proceedings of the New Zealand Grassland Association* **46**, 57-63.

- Barlow, E.W.R., Boersma, L. and Young, J.L. (1976). Root temperature and soil water potential effects on growth and soluble carbohydrate concentration of corn seedlings. *Crop Science* **16**, 59-62.
- Barnard, C. and Frankel, O.H. (1964). Grass, grazing animals, and man in historic perspective. In 'Grasses and Grasslands.' (Ed. C. Barnard.) pp. 1-12. (MacMillan and Company Ltd.: London.)
- Bartholomew, P.E. and Booyesen, P. de V. (1969). The influence of clipping frequency on reserve carbohydrates and regrowth of *Eragrostis curvula*. *Proceedings of the Grassland Society of South Africa* **4**, 35-43.
- Barthram, G.T. and Grant, S.A. (1995) Interactions between variety and the timing of conservation cuts on species balance in *Lolium perenne* - *Trifolium repens* swards. *Grass and Forage Science* **50**, 98-105.
- Baur-Hoch, B., Machler, F. and Nosberger, J. (1990). Effect of carbohydrate demand on the remobilisation of starch in stolons and roots of white clover (*Trifolium repens* L.) after defoliation. *Journal of Experimental Botany* **41**, 573-8.
- Beard, J. (1973). Temperature. In 'Turfgrass: Science and Culture.' pp. 209-60. (Prentice-Hall Inc.: New Jersey.)
- Bettany, A.J. (1995a). Stress responses in cell cultures of *Lolium temulentum*. I. Transcriptional and post-transcriptional changes in gene expression during heat shock and recovery. *Journal of Plant Physiology* **146**, 162-8.
- Bettany, A.J. (1995b). Stress responses in cell cultures of *Lolium temulentum*. II. Induction of thermotolerance and associated changes in gene expression. *Journal of Plant Physiology* **146**, 183-9.
- Biddiscombe, E.F., Rogers, A.L. and Maller, R.A. (1977). Summer dormancy, regeneration and persistence of perennial grasses in south-western Australia. *Australian Journal of Experimental Agriculture and Animal Husbandry* **17**, 795-801.
- Blaser, R.E., Brown, R.H. and Bryant, H.T. (1966). The relationship between carbohydrate accumulation and growth of grasses under different microclimates. *Proceedings of the 10th International Grassland Congress, Helsinki*, 147-50.
- Bommer, D.F.R. (1966). Influence of cutting frequency and nitrogen level on the carbohydrate reserves of three grass species. *Proceedings of the 10th International*

Grasslands Congress, Helsinki, 156-60.

- Booyesen, P. de V. and Nelson, C.J. (1975). Leaf area and carbohydrate reserves in regrowth of tall fescue. *Crop Science* **15**, 262-6.
- Boyd, A.G. and Frame, J. (1983). Response of white clover to various management factors. In 'Efficient Grassland Farming.' (Ed. A.J. Corrall) *Proceedings of the 9th General Meeting of the European Grassland Federation*, 213-6. (British Grassland Society: Great Britain.)
- Bredemeijer, G.M.M. and Esselink, G. (1994). Phosphofructokinase in relation to sugar accumulation in cold-hardened *Lolium perenne* L. cultivars. *Journal of Plant Physiology* **143**, 112-8.
- Brereton, A.J., Carton, O.T. and Conway, A. (1985). The effect of grass tiller density on the performance of white clover. *Proceedings of the 15th International Grassland Congress, Kyoto*, 756-7.
- Brink, G.E. (1995). White clover growth and morphology under contrasting cutting regimes. *Crop Science* **35**, 1100-3.
- Brock, J.L. and Caradus, J.R. (1996). Influence of grazing management and drought on white clover population performance and genotypic frequency. In 'White Clover: New Zealand's Competitive Edge.' (Ed. D.R. Woodfield.) pp. 79-82. (Agronomy Society of New Zealand: Christchurch.)
- Brock, J.L. and Fletcher, R.H. (1993). Morphology of perennial ryegrass (*Lolium perenne*) plants in pastures under intensive sheep grazing. *Journal of Agricultural Science, Cambridge* **120**, 301-10.
- Brock, J.L. and Hay, M.J.M. (1996). A review of the role of grazing management on the growth and performance of white clover cultivars in lowland New Zealand pastures. In 'White Clover: New Zealand's Competitive Edge.' (Ed. D.R. Woodfield.) pp. 65-70. (Agronomy Society of New Zealand: Christchurch.)
- Brock, J.L. and Thomas, V.J. (1991). The pasture ryegrass plant, what is it? *Proceedings of the New Zealand Grassland Association* **53**, 111-6.
- Brock, J.L., Hay, M.J.M., Thomas, V.J. and Sedcole, J.R. (1988). Morphology of white clover (*Trifolium repens* L.) plants in pastures under intensive sheep grazing. *Journal of Agricultural Science, Cambridge* **111**, 273-83.

- Brock, J.L., Hume, D.E. and Fletcher, R.H. (1996). Seasonal variation in the morphology of perennial ryegrass (*Lolium perenne*) and cocksfoot (*Dactylis glomerata*) plants and populations in pastures under intensive sheep grazing. *Journal of Agricultural Science, Cambridge* **126**, 37-51.
- Brougham, R.W. (1958a). Interception of light by the foliage of pure and mixed stands of pasture plants. *Australian Journal of Agricultural Research* **9**, 39-52.
- Brougham, R.W. (1958b). Leaf development in swards of white clover (*Trifolium repens* L.). *New Zealand Journal of Agricultural Research* **1**, 707-18.
- Brougham, R.W., Ball, P.R. and Williams, W.M. (1978). The ecology and management of white clover-based pastures. In 'Plant Relations in Pastures.' (Ed. J.R. Wilson.) (Commonwealth Scientific and Industrial Research Organisation: Melbourne.)
- Brouwer, R. (1966). Root growth of grasses and cereals. In 'The Growth of Cereals and Grasslands.' (Ed. F.L. Milthorpe and J.D. Ivins.) pp. 153-66. (Butterworths: London.)
- Bullock, J.M., Clearhill, B. and Silvertown, J. (1994). Tiller dynamics of two grasses - responses to grazing, density and weather. *Journal of Ecology* **82**, 331-40.
- Butler, G.W., Greenwood, R.M. and Soper, K. (1959). Effects of shading and defoliation on the turnover of root and nodule tissue of plants of *Trifolium repens*, *Trifolium pratense*, and *Lotus uliginosus*. *New Zealand Journal of Agricultural Research* **2**, 415-26.
- Campbell, A.G. (1969). Grazing interval, stocking rate, and pasture production. *New Zealand Journal of Agricultural Research* **12**, 67-74.
- Caradus, J.R. (1977). Structural variation of white clover root systems. *New Zealand Journal of Agricultural Research* **20**, 213-9.
- Caradus, J.R. (1980). Distinguishing between grass and legume species for efficiency of phosphorus use. *New Zealand Journal of Agricultural Research* **23**, 75-81.
- Carlson, G.E., Gibson, P.B. and Baltensperger, D.D. (1985). White clover and other perennial clover. In 'Forages: the Science of Grassland Agriculture.' (Eds. M.E. Heath, R.F. Barnes and D.S. Metcalfe.) pp. 118-27. (Iowa State University Press: Iowa.)
- Carroll, J.C. (1943). Effects of drought, temperature and nitrogen on turf grasses. *Plant Physiology* **18**, 19-36.

- Chapman, D.F. (1983). Growth and demography of *Trifolium repens* stolons in grazed hill pastures. *Journal of Applied Ecology* **20**, 597-608.
- Clark, S.G. and McFadden, M.J. (1997). Yield and stolon characteristics of white clover (*Trifolium repens* L.) cultivars in perennial pasture in Victoria. *Australian Journal of Experimental Agriculture* **37**, 159-64.
- Clement, C.R., Hopper, M.J., Jones, L.H.P. and Leafe, E.L. (1978). The uptake of nitrate by *Lolium perenne* from flowing nutrient solution. II. Effect of light, defoliation, and relationship to CO₂ flux. *Journal of Experimental Botany* **29**, 1173-83.
- Clifford, P.E. (1977). Tiller bud suppression in reproductive plants of *Lolium multiflorum* Lam. cv. Westerwoldicum. *Annals of Botany* **41**, 605-15.
- Colby, W.G., Drake, M., Oohara, H. and Yoshida, N. (1966). Carbohydrate reserves in orchardgrass. *Proceedings of the 10th International Grassland Congress, Helsinki*, 151-5.
- Colby, W.G., Fenner, H., Drake, M., Oohara, H., Yoshida, N., Fukanga, K. and Oohara, Y. (1974). Role of soluble carbohydrate reserves in the growth and performance of timothy (*Phleum pratense*). *Proceedings of the 12th International Grassland Congress, Moscow* **1**, 322-30.
- Collins, R.P. and Rhodes, I. (1995). Stolon characteristics related to winter survival in white clover. *Journal of Agricultural Science, Cambridge* **124**, 11-6.
- Colman, R.L. (1964). Root knot nematodes as a cause of poor growth of clover on the far north coast of N.S.W. *The Agricultural Gazette of New South Wales*, October 1964, 1367-8.
- Colman, R.L. and Lazenby, A. (1970). Factors affecting the response of some tropical and temperate grasses to fertiliser nitrogen. *Proceedings of the 11th International Grassland Congress, Surfers Paradise*, 392-7.
- Colvill, K.E. and Marshall, C. (1981). The patterns of growth, assimilation of ¹⁴CO₂ and distribution of ¹⁴C-assimilate within vegetative plants of *Lolium perenne* at low and high density. *Annals of Applied Biology* **99**, 179-90.
- Colvill, K.E. and Marshall, C. (1984). Tiller dynamics and assimilate partitioning in *Lolium perenne* with particular reference to flowering. *Annals of Applied Biology* **104**, 543-57.

- Cooper, J.P. and Tainton, N.M. (1968). Light and temperature requirements for the growth of tropical and temperate grasses. *Herbage Abstracts* **38**, 167-76.
- Corleto, A. and Laude, H.M. (1974). Evaluating growth potential after drought stress. *Crop Science* **14**, 224-7.
- Cornish, P.S. (1987). Root growth and function in temperate pastures. In 'Temperate Pastures: Their Production, Use and Management.' (Eds. J.L. Wheeler, C.J. Pearson and G.E. Robards.) pp. 79-98. (Australian Wool Corporation/Commonwealth Scientific and Industrial Research Organisation: Australia.)
- Corre, N., Bouchart, V., Ourry, A. and Boucaud, J. (1996). Mobilisation of nitrogen reserves during regrowth of defoliated *Trifolium repens* L. and identification of potential vegetative storage proteins. *Journal of Experimental Botany* **47**, 1111-8.
- Crider, F.J. (1955). Root - growth stoppage resulting from defoliation of grass. United States Department of Agriculture Technical Bulletin 1102. pp. 3-23.
- Culvenor, R.A. and Simpson, R.J. (1991). Mobilisation of nitrogen in swards of *Trifolium subterraneum* L. during regrowth after defoliation. *New Phytologist* **117**, 81-90.
- Curtis, O.F. (1944). The food content of forage crops as influenced by the time of day at which they are cut. *Journal of the American Society of Agronomy* **36**, 401-16.
- Danckwerts, J.E. (1993). Reserve carbon and photosynthesis: their role in regrowth of *Themeda triandra*, a widely distributed subtropical graminaceous plant. *Functional Ecology* **7**, 634-41.
- Danckwerts, J.E. and Gordon, A.J. (1987). Long-term partitioning, storage and remobilisation of ¹⁴C assimilated by *Lolium perenne* (cv. Melle). *Annals of Botany* **59**, 55-66.
- Danckwerts, J.E. and Gordon, A.J. (1989). Long-term partitioning, storage and remobilisation of ¹⁴C assimilated by *Trifolium repens* (cv. Blanca). *Annals of Botany* **64**, 533-44.
- Davidson, J.L. and Milthorpe, F.L. (1966a). Leaf growth in *Dactylis glomerata* following defoliation. *Annals of Botany* **30**, 173-84.
- Davidson, J.L. and Milthorpe, F.L. (1966b). The effect of defoliation on the carbon balance in *Dactylis glomerata*. *Annals of Botany* **30**, 185-98.

- Davidson, R.L. (1969). Effect of root/leaf temperature differentials on root/shoot ratios in some pasture grasses and clover. *Annals of Botany* **33**, 561-9.
- Davies, A.G. (1960). Conditions influencing primary growth and regrowth in perennial ryegrass. *Report of the Welsh Plant Breeding station for 1959*, pp. 110-6.
- Davies, A. (1965). Carbohydrate levels and regrowth in perennial ryegrass. *Journal of Agricultural Science* **65**, 213-21.
- Davies, A. (1966). The regrowth of swards of S24 perennial ryegrass subjected to different pretreatments. *Journal of Agricultural Science Cambridge* **67**, 139-44.
- Davies, A. (1974). Leaf tissue remaining after cutting and regrowth in perennial ryegrass. *Journal of Agricultural Science, Cambridge* **82**, 165-72.
- Davies, I. (1969). The influence of management on tiller development and herbage growth. Welsh Plant Breeding Station, Aberystwyth Technical Bulletin Number 3.
- Deinum, B. (1966). Climate, nitrogen and grass: research into the influence of light intensity, temperature, water supply and nitrogen on the production and chemical composition of grass. *Mededelingen Landbouwhogeschool Wageningen 66-11*, 1-91.
- Dennis, W.D. and Woledge, J. (1985). The effect of nitrogenous fertiliser on the photosynthesis and growth of white clover/perennial ryegrass swards. *Annals of Botany* **55**, 171-8.
- Dilz, K. (1966). The effect of nitrogen nutrition and clipping frequency on regrowth of perennial ryegrass. *Proceedings of the 10th International Grassland Congress, Helsinki*, 160-4.
- Di Mascio, J.A., Sweeney, P.M., Danneberger, T.K. and Kamalay, J.C. (1994). Analysis of heat shock response in perennial ryegrass using maize heat shock protein clones. *Crop Science* **34**, 798-804.
- Downes, R.W. (1970). Differences between tropical and temperate grasses in rates of photosynthesis and transpiration. *Proceedings of the 11th International Grassland Congress, Surfers Paradise*, 527-30.
- Downes, R.W. and Hesketh, J.D. (1968). Enhanced photosynthesis at low oxygen concentrations: differential response of temperate and tropical grasses. *Planta* **78**, 79-84.

- Eagles, C.F. and Williams, J. (1992). Hardening and dehardening of *Lolium perenne* in response to fluctuating temperatures. *Annals of Botany* **70**, 333-8.
- Earle, D.F. and McGowan A.A. (1979). Evaluation and calibration of an automated rising plate meter for estimating dry matter yield of pasture. *Australian Journal of Experimental Agriculture and Animal Husbandry* **19**, 337-43.
- Edmond, D.B. (1964). Some effects of sheep treading on the growth of 10 pasture species. *New Zealand Journal of Agricultural Research* **7**, 1-16.
- Edwards, P.J. (1965). The construction and use of light-proof boxes in grassland research work. *Proceedings of the 9th International Grassland Congress, Sao Paulo*, 1483-5.
- Evans, D.R. and Williams, T.A. (1987) The effect of cutting and grazing managements on dry matter yield of white clover varieties (*Trifolium repens*) when grown with S23 perennial ryegrass. *Grass and Forage Science* **42**, 153-9.
- Evans, D.R., Williams, T.A. and Evans, S.A. (1992). Evaluation of white clover varieties under grazing and their role in farm systems. *Grass and Forage Science* **47**, 342-52.
- Evans, L.T. (1960). The influence of temperature on flowering in species of *Lolium* and in *Poa pratensis*. *Journal of Agricultural Science, Cambridge* **54**, 410-6.
- Evans, L.T., Wardlaw, I.F. and Williams, C.N. (1964). Environmental control of growth. In 'Grasses and Grasslands.' (Ed. C. Barnard.) pp. 102-25. (MacMillan and Co.: London.)
- Evans, P.S. (1970). Root growth of *Lolium perenne* L. I. Effect of plant age, seed weight, and nutrient concentration on root weight, length, and number of apices. *New Zealand Journal of Botany* **8**, 344-56.
- Evans, P.S. (1971). Root growth of *Lolium perenne* L. II. Effects of defoliation and shading. *New Zealand Journal of Agricultural Research* **14**, 552-62.
- Evans, P.S. (1972). Root growth of *Lolium perenne* L. III. Investigation of the mechanism of defoliation - induced suppression of elongation. *New Zealand Journal of Agricultural Research* **15**, 347-55.
- Evans, P.S. (1973). The effect of repeated defoliation to three different levels on root growth of five pasture species. *New Zealand Journal of Agricultural Research* **16**, 31-4.

- Evans, P.S. (1977). Comparative root morphology of some pasture grasses and clovers. *New Zealand Journal of Agricultural Research* **20**, 331-5.
- Evans, P.S. (1978). Plant root distribution and water use patterns of some pasture and crop species. *New Zealand Journal of Agricultural Research* **21**, 261-5.
- Eyles, A.G. and Cameron, D.G. (1985). Pasture research in northern Australia - its history, achievements and future emphasis. (Ed. J.B. Hacker.) Commonwealth Scientific and Industrial Research Organisation Division of Tropical Crops and Pastures Research Report number 4. (Commonwealth Scientific and Industrial Research Organisation: Brisbane.)
- Fairey, N.A. (1985). Productivity, quality and persistence of perennial ryegrass as influenced by cutting/fertility management and ploidy. *Canadian Journal of Plant Science* **65**, 565-71.
- Frame, J. (1966a). The effects of cutting and grazing techniques on productivity of grass/clover swards. *Proceedings of the 9th International Grassland Congress, Sao Paulo*, 1511-6.
- Frame, J. (1966b). The evaluation of herbage production under cutting and grazing regimes. *Proceedings of the 10th International Grassland Congress, Helsinki*, 291-6.
- Frame, J. and Boyd, A.G. (1987). The effect of fertiliser nitrogen rate, white clover variety and closeness of cutting on herbage productivity from perennial ryegrass/white clover swards. *Grass and Forage Science* **42**, 85-96.
- Frame, J. and Newbould, P. (1986). Agronomy of White Clover. *Advances in Agronomy* **40**, 1-88.
- Frankow-Lindberg, B.E. (1997). Assimilate partitioning in three white clover cultivars in the autumn, and the effect of defoliation. *Annals of Botany* **79**, 83-7.
- Fulkerson, W.J. (1994). Effect of redefoliation on the regrowth and water soluble carbohydrate content of *Lolium perenne*. *Australian Journal of Agricultural Research* **45**, 1809-15.
- Fulkerson, W.J. and Bryant, R. (1994). Defoliation frequency up to, and after, stem elongation in *Lolium perenne*. Wollongbar Agricultural Institute Dairy Research Report 1993/94. pp. 10-1. (New South Wales Agriculture: Australia.)

- Fulkerson, W.J. and Lowe, K.F. (1994) Limits to dairy production in the tropics: forage and crop feeding systems. Invited Speakers' Abstracts S18, p. 9. 24th International Dairy Congress Brief Communications. (Australian National Committee of the International Dairy Federation: Melbourne.)
- Fulkerson, W.J. and Reeves, M. (1996). Management and productivity of white clover in a kikuyu grass sward in subtropical Australia. *Proceedings of the New Zealand Grassland Association* **57**, 199-201.
- Fulkerson, W.J. and Slack, K. (1993). Estimating mass of temperate and tropical pastures in the subtropics. *Australian Journal of Experimental Agriculture* **33**, 865-9.
- Fulkerson, W.J. and Slack, K. (1994a). Management of *Lolium perenne*/*Trifolium repens* pastures in the subtropics. II. Effect of summer defoliation, irrigation duration, seedbed preparation and pasture type. *Australian Journal of Agricultural Research* **45**, 721-8.
- Fulkerson, W.J. and Slack, K. (1994b). Leaf number as a criterion for determining defoliation time for *Lolium perenne*: 1. Effect of water-soluble carbohydrates and senescence. *Grass and Forage Science* **49**, 373-7.
- Fulkerson, W.J. and Slack, K. (1995). Leaf number as a criterion for determining defoliation time for *Lolium perenne*: 2. Effect of defoliation frequency and height. *Grass and Forage Science* **50**, 16-20.
- Fulkerson, W.J. and Slack, K. (1996a). Productivity and persistence of irrigated lotus (*Lotus pedunculatus*) and white clover (*Trifolium repens*) in a kikuyu (*Pennisetum clandestinum*) pasture. *Tropical Grasslands* **30**, 353-62.
- Fulkerson, W.J. and Slack, K. (1996b). Maintaining white clover (*Trifolium repens*) in a kikuyu (*Pennisetum clandestinum*) pasture in the subtropics. *Proceedings of the 8th Australian Agronomy Conference, Toowoomba*, 270-2.
- Fulkerson, W.J. and Yeoman, M. (1994). Relationship between seeding rate and production and persistence of perennial ryegrass (*Lolium perenne*)/white clover (*Trifolium repens*) pastures under grazing. Wollongbar Agricultural Institute Dairy Research Report 1993/94. pp. 13-14. (New South Wales Agriculture: Australia.)
- Fulkerson, W.J., Lowe, K.J., Ayres, J.F. and Launders, T. (1993a). Northern dairy feedbase 2001. 3. Winter pastures and crops. *Tropical Grasslands* **27**, 162-79.

- Fulkerson, W.J., Norton, M. and Slack, K. (1993c). White clover evaluation - genotype x environment matching. Wollongbar Agricultural Institute Dairy Research Report 1992/93. pp. 22-3. (New South Wales Agriculture: Australia.)
- Fulkerson, W.J., Slack, K. and Lowe, K.F. (1994). Variation in the response of *Lolium* genotypes to defoliation. *Australian Journal of Agricultural Research* **45**, 1309-17.
- Fulkerson, W.J., Slack, K., Moore, K. and Rolfe, C. (1993b). Management of *Lolium perenne*/*Trifolium repens* pastures in the subtropics. I. Effect of defoliation interval, seeding rate and application of N and lime. *Australian Journal of Agricultural Research* **44**, 1947-58.
- Gallagher, J.A., Volenec, J.J., Turner, L.B. and Pollock, C.J. (1994). Starch-hydrolytic enzymes in white clover (*Trifolium repens* L.) stolons. The Working Group on Pasture Ecology Newsletter 33, 12-14.
- Gao, Y. and Wilman, D. (1994). Leaf development in eight related grasses. *Journal of Agricultural Science, Cambridge* **123**, 41-6.
- Garden, D.L. (1988). Effects of stocking rate and site characteristics on the pathway of persistence of *Trifolium repens* L. in a subtropical environment. Masters Thesis. The University of Queensland.
- Garwood, E.A. (1967). Seasonal variation in appearance and growth of grass roots. *Journal of the British Grassland Society* **22**, 121-30.
- Garwood, E.A. and Sinclair, J. (1979). Use of water by six grass species. 2. Root distribution and use of soil water. *Journal of Agricultural Science, Cambridge* **93**, 25-35.
- Gay, A.P. and Thomas, H. (1995). Leaf development in *Lolium temulentum* L.: photosynthesis in relation to growth and senescence. *New Phytologist* **130**, 159-68.
- Genstat Committee. (1993). Genstat 5 Release 3 Reference Manual. pp. 539-83. (Oxford University Press: Great Britain.)
- Gifford, R.M. and Marshall, C. (1973). Photosynthesis and assimilate distribution in *Lolium multiflorum* Lam. following differential tiller defoliation. *Australian Journal of Biological Science* **26**, 517-26.
- Gonzalez, B., Boucaud, J., Salette, J., Langlois, J. and Duyme, M. (1989). Changes in stubble carbohydrate content during regrowth of defoliated perennial ryegrass (*Lolium perenne* L.) on two nitrogen levels. *Grass and Forage Science* **44**, 411-5.

- Gordon, A.J., Kessler, W. and Minchin, F.R. (1990). Defoliation-induced stress in nodules of white clover. 1. Changes in physiological parameters and protein synthesis. *Journal of Experimental Botany* **41**, 1245-53.
- Gordon, A.J., Ryle, G.J.A., Mitchell, D.F., Lowry, K.H. and Powell, C.E. (1986). The effect of defoliation on carbohydrate, protein and leghaemoglobin content of white clover nodules. *Annals of Botany* **58**, 141-54.
- Grant, S.A., Barthram, G.T. and Torvell, L. (1981). Components of regrowth in grazed and cut *Lolium perenne* swards. *Grass and Forage Science* **36**, 155-68.
- Griffith, S.M. (1992). Changes in post-anthesis assimilates in stem and spike components of Italian ryegrass (*Lolium multiflorum* Lam.). I. Water soluble carbohydrates. *Annals of Botany* **69**, 243-8.
- Hartwig, U.A., Heim, I., Luscher, A. and Nosberger, J. (1994). The nitrogen-sink is involved in the regulation of nitrogenase activity in white clover after defoliation. *Physiologia Plantarum* **92**, 375-82.
- Hay, M.J.M. and Newton, P.C.D. (1996). Effect of severity of defoliation on the viability of reproductive and vegetative axillary buds of *Trifolium repens* L. *Annals of Botany* **78**, 117-23.
- Hay, M.J.M., Brock, J.L. and Thomas, V.J. (1989b). Density of *Trifolium repens* plants in mixed swards under intensive grazing by sheep. *Journal of Agricultural Science, Cambridge* **113**, 81-6.
- Hay, M.J.M., Chu, A.C.P., Knighton, M.V. and Wewala, S. (1989a). Variation with season and node position in carbohydrate content of white clover stolons. *Proceedings of the 16th International Grassland Congress, Nice*, 1059-60.
- Hay, M.J.M., Kim, M., McLean, R.W. and Gonzalez-Rodriguez, A. (1993). Defoliation of young branch and parent stolons of white clover in rotationally grazed pastures. *Proceedings of the 17th International Grassland Congress, Palmerston North*, 128-9.
- Hayes, M.J. and Williams, E.D. (1995). Effects of timing and intensity of defoliation on growth and development of white clover seedlings. *Grass and Forage Science* **50**, 414-23.
- Heide, O.M. (1994). Control of flowering and reproduction in temperate grasses. *New Phytologist* **128**, 347-62.

- Hodgkinson, K.C. and Baas Becking, H.G. (1977). Effect of defoliation on root growth of some arid zone perennial plants. *Australian Journal of Agricultural Research* **29**, 31-42.
- Hoen, K. (1966). The effect of temperature and soil moisture on summer survival of two varieties of *Phalaris tuberosa*. *Australian Journal of Experimental Agriculture and Animal Husbandry* **6**, 280-2.
- Hoen, K. (1968). Summer dormancy in *Phalaris tuberosa* L. *Australian Journal of Agricultural Research* **19**, 227-39.
- Hoglund, J.H. and Brock, J.L. (1987). Nitrogen fixation in managed grasslands. In 'Ecosystems of the World 17B: Managed Grasslands - Analytical Studies..' (Ed. R.W. Snaydon.) pp. 187-96. (Elsevier: Amsterdam.)
- Hollowell, E.A. (1966). White clover (*Trifolium repens* L.), annual or perennial? *Proceedings of the 10th International Grassland Congress, Helsinki*, 184-7.
- Holt, D.A. and Hilst, A.R. (1969). Daily variation in carbohydrate content of selected forage crops. *Agronomy Journal* **611**, 239-42.
- Horst, G.L. and Nelson, C.J. (1979). Compensatory growth of tall fescue following drought. *Agronomy Journal* **71**, 559-63.
- Hoveland, C.S. (1970). Dormancy and seasonal growth of *Phalaris* species in Alabama. *Proceedings of the 11th International Grassland Congress, Surfers Paradise*, 608-11.
- Hughes, R. and Jackson, D.K. (1974). Impact of grazing management on sward survival. *Journal of the British Grassland Society* **29**, 76.
- Hume, D.E. (1991). Effect of cutting on production and tillering in prairie grass (*Bromus willdenowii* Kunth) compared to two ryegrass (*Lolium*) species. 1. Vegetative plants. *Annals of Botany* **67**, 533-41.
- Hunt, L.A. and Brougham, R.W. (1967). Some changes in the structure of a perennial ryegrass sward frequently but leniently defoliated during the summer. *New Zealand Journal of Agricultural Research* **10**, 397-404.
- Hutchinson, K.J. (1970). The persistence of perennial species under intensive grazing in a cool temperate environment. *Proceedings of the 11th International Grassland Congress, Surfers Paradise*, 611-4.

- Hyder, D.N. (1972). Defoliation in relation to vegetative growth. In 'The Biology and Utilisation of Grasses.' (Eds. V.B. Younger and C.M. McKell.) pp. 304-17. (Academic Press Inc.: New York.)
- Irwin, J.A.G. and Jones, R.M. (1977). The role of fungi and nematodes as factors associated with death of white clover (*Trifolium repens*) stolons over summer in south-eastern Queensland. *Australian Journal of Experimental Agriculture and Animal Husbandry* **17**, 789-94.
- Jacques, W.A. and Edmond, D.B. (1952). Root development in some common New Zealand pasture plants. V. The effect of defoliation and root pruning on cocksfoot (*Dactylis glomerata*) and perennial ryegrass (*Lolium perenne*). *New Zealand Journal of Science and Technology* **34**, 231-48.
- Jewiss, O.R. (1966). Morphological and physiological aspects of growth of grasses during the vegetative phase. In 'The Growth of Cereals and Grasses.' (Eds. F.L. Milthorpe and J.D. Ivins.) pp. 39-54. (Butterworths: London.)
- Johansson, G. (1993). Carbon distribution in grass (*Festuca pratensis* L.) during regrowth after cutting - utilisation of stored and newly assimilated carbon. *Plant and Soil* **151**, 11-20.
- Johnson, I.R. (1990). Plant respiration in relation to growth, maintenance, iron uptake and nitrogen assimilation. *Plant, Cell and Environment* **13**, 319-28.
- Johnston, A. and Booth, C. (1983). 'Plant Pathologist's Pocketbook.' pp. 237, 239. (Commonwealth Mycological Institute: Great Britain.)
- Jones, R.J., Davies, J.G., Waite, R.B. and Fergus, I.F. (1968). The production and persistence of grazed irrigated pasture mixes in south-eastern Queensland. *Australian Journal of Experimental Agriculture and Animal Husbandry* **8**, 177-89.
- Jones, R.M. (1980). Survival of seedlings and primary taproots of white clover (*Trifolium repens*) in subtropical pastures in south-east Queensland. *Tropical Grasslands* **14**, 19-22.
- Jones, R.M. (1982). White clover (*Trifolium repens*) in subtropical south-east Queensland. I. Some effects of site, season and management practices on the population dynamics of white clover. *Tropical Grasslands* **16**, 118-27.
- Jones, R.M. and Evans, T.R. (1977). Soil seed levels of *Lotononis bainesii*, *Desmodium intortum* and *Trifolium repens* in subtropical pastures. *The Journal of the*

- Australian Institute of Agricultural Science* **43**, 164-6.
- Julander, O. (1945). Drought resistance in range and pasture grasses. *Plant Physiology* **20**, 573-99.
- Kang, J.H. and Brink, G.E. (1995). White clover morphology and physiology in response to defoliation interval. *Crop Science* **35**, 264-9.
- Kays, S. and Harper, J.L. (1974). The regulation of plant and tiller density in a grass sward. *The Journal of Ecology* **62**, 97-105.
- Ketellapper, H.J. (1960). The effect of soil temperature on the growth of *Phalaris tuberosa* L. *Physiologia Plantarum* **13**, 641-7.
- Kim, T.H., Ourry, A., Boucaud, J. and Lemaire, G. (1991). Changes in source-sink relationship for nitrogen during regrowth of lucerne (*Medicago sativa* L.) following removal of shoots. *Australian Journal of Plant Physiology* **18**, 593-602.
- Kim, T.H., Ourry, A., Boucaud, J. and Lemaire, G. (1993). Partitioning of nitrogen derived from N₂ fixation and reserves in nodulated *Medicago sativa* L. during regrowth. *Journal of Experimental Botany* **44**, 266, 555-62.
- King, J., Lamb, W.I.C. and McGregor, M.T. (1978). Effect of partial and complete defoliation on regrowth of white clover plants. *Journal of the British Grassland Society* **33**, 49-55.
- Kingsbury, L.R. (1965). Pasture quality in terms of soluble carbohydrates and volatile fatty acid production. *Proceedings of the New Zealand Society of Animal Production* **25**, 119-34.
- Knight, R. (1966). The growth of cocksfoot (*Dactylis glomerata* L.) in a winter-rainfall environment. *Proceedings of the 10th International Grassland Congress, Helsinki*, 308-11.
- Korte, C.J. and Chu, A.C.P. (1983). Some effects of drought on perennial ryegrass swards. *Proceedings of the New Zealand Grassland Association* **44**, 211-6.
- Lambers, H. (1985). Respiration in intact plants and tissues: its regulation and dependence on environmental factors, metabolism and invaded organisms. In 'Encyclopaedia of Plant Physiology Volume 18.' (Eds. A. Pirson and M.H. Zimmerman.) pp. 418-73. (Springer-Verlag: Berlin.)
- Lambert, J.P., Rumball, P.J. and Boyd, A.F. (1979). Comparison of ryegrass-white clover pastures with and without paspalum and kikuyu grass. 1. Pasture production. *New*

Zealand Journal of Experimental Agriculture 7, 259-302.

- Lambourne, D. (1985). Pasture Systems for Milk Production. The University of Sydney Post-Graduate Committee in Veterinary Science, Proceedings number 78, Dairy Cattle Production.
- Lancashire, J.A. (1974). Some effects of grazing interval and crown rust on a perennial ryegrass/white clover pasture during summer and autumn. *Proceedings of the 12th International Grassland Congress, Moscow* 3, 272-9.
- Lange, O.L. (1965). The heat resistance of plants, its determination and variability. In 'Methodology of Plant Eco-Physiology.' (Ed. F. Eckhardt.) pp. 399-405. (Montpellier Symposium: United Nations Educational, Scientific and Cultural Organisation: Paris.)
- Langer, R.H.M. (1966). Mineral nutrition in grasses and cereals. In 'The Growth of Cereals and Grasses.' (Eds. F.L. Milthorpe and J.D. Ivins.) pp. 213-26. (Butterworths: London).
- Langer, R.H.M. (1979). 'How Grasses Grow.' Second Edition. The Institute of Biology. Studies in Biology No. 34. (Edward Arnold Ltd.: London.)
- Langer, R.H.M. (1990). Pasture Plants. In 'Pastures: Their Ecology and Management.' (Ed. R.H.M. Langer.) pp. 39-74. (Oxford University Press: Auckland.)
- Laude, H.M. (1953). The nature of summer dormancy in perennial grasses. *Botanical Gazette* 114, 284-92.
- Lawson, A.R., Kelly, K.B. and Sale, P.W.G. (1995). The effect of defoliation frequency on reserve utilisation and regrowth of two white clover cultivars. Institute of Sustainable Irrigated Agriculture Report 1993-95. p. 97. Victorian Department of Agriculture, Energy and Minerals.
- Lee, C.K., Cunningham, P.J. and Anderson, M.W. (1993). Breeding temperate forages for winter growth and persistence. *Proceedings of the 10th Australian Plant Breeding Conference*, 67-71.
- Leon, M., Laine, P., Ourry, A. and Boucaud, J. (1995). Increased uptake of native soil nitrogen by roots of *Lolium multiflorum* Lam. after nitrogen fertilisation is explained by a stimulation of the uptake process itself. *Plant and Soil* 173, 197-203.

- Levitt, J. (1980). High-temperature or heat stress. In 'Response of Plants to Environmental Stresses. Volume 1: Chilling, Freezing and High Temperature Stresses.' Second Edition. pp. 347-93. (Academic Press Inc.: Orlando.)
- L'Huillier, P.J. (1987). Tiller appearance and death of *Lolium perenne* in mixed swards grazed by dairy cattle at two stocking rates. *New Zealand Journal of Agricultural Research* **30**, 15-22.
- Lowe, K.F. and Bowdler, T.M. (1984). Performance of temperate and tropical grasses and legumes under two irrigation frequencies in south-eastern Queensland. *Tropical Grasslands* **18**, 46-55.
- Lowe, K.F. and Bowdler, T.M. (1995). Growth, persistence and rust sensitivity of irrigated, perennial temperate grasses in the Queensland subtropics. *Australian Journal of Experimental Agriculture* **35**, 571-8.
- Lowe, K.F., Bowdler, T.M., Casey, N.D., Fulkerson, W.J. and Launders, T. (1995). Evaluation of temperate species in the subtropics. Annual summary of the evaluation of temperate grasses and legumes conducted by the Queensland Department of Primary Industries and New South Wales Agriculture.
- Lowe, K.F., Bowdler, T.M., Reason, G.K. and Moss, R.J. (1993). The value of adapted, annual ryegrasses for sub-tropical dairy production. *Proceedings of the 17th International Grassland Congress, Palmerston North*, 407-8.
- Lowe, K.F., Bowdler, T.M., Shaw, R.J. and Geritz, A.F. (1992). Effect of pugging on a black earth soil: establishment and production of a subsequent annual ryegrass crop. Queensland Department of Primary Industries Project Report Q092011. (Queensland Department of Primary Industries: Brisbane.)
- Lucanus, R., Mitchell, K.J., Pritchard, G.G. and Calder, D.M. (1960). Factors influencing survival of strains of ryegrass during the summer. *New Zealand Journal of Agricultural Research* **3**, 185-93.
- Ludlow, M.M. (1978). Light relations of pasture plants. In 'Plant Relations in Pastures.' (Ed. J.R. Wilson.) pp. 35-49. (Commonwealth Scientific and Industrial Research Organisation: Melbourne.)
- Madison, J.H. (1962). Mowing of turfgrass. II. Responses of three species of grass. *Agronomy Journal* **54**, 250-2.

- Marshall, C. (1987). Physiological aspects of pasture growth. In 'Ecosystems of the World 17B: Managed Grasslands - Analytical Studies.' (Ed. R.W. Snaydon.) pp. 29-46. (Elsevier: Amsterdam.)
- Marshall, C. and Sagar, G.R. (1965). The influence of defoliation on the distribution of assimilates in *Lolium multiflorum* Lam. *Annals of Botany* **29**, 365-72.
- Matthew, C., Chu, A.C.P., Hodgson, J. and MacKay, A.D. (1991). Early summer pasture control: what suits the plant? *Proceedings of the New Zealand Grassland Association* **53**, 73-7.
- Matthew, C., Quilter, S.J., Korte, C.J., Chu, A.C.P. and MacKay, A.D. (1989). Stolon formation and significance for sward tiller dynamics in perennial ryegrass. *Proceedings of the New Zealand Grassland Association* **50**, 255-9.
- Mawson, W.F.Y. and White, B.J. (1971). Climatic and biological limitations to dairy production in a tropical environment. *Tropical Grasslands* **5**, 145-58.
- May, L.H. (1960). The utilisation of carbohydrate reserves in pasture plants after defoliation. *Herbage Abstracts* **30**, 239-45.
- McGrath, D. (1992). A note on the influence of nitrogen application and time of cutting on water soluble carbohydrate production by Italian ryegrass. *Irish Journal of Agricultural and Food Research* **31**, 189-92.
- McKenzie, F.R. (1996). The influence of grazing frequency and intensity on the vigour of *Lolium perenne* L. under subtropical conditions. *Australian Journal of Agricultural Research* **47**, 975-93.
- McKenzie, F.R. (1997). Influence of grazing frequency and intensity on tiller appearance and death rates of *Lolium perenne* L. under subtropical conditions. *Australian Journal of Agricultural Research* **48**, 337-42.
- McWilliam, J.R. (1968). The nature of the perennial response in Mediterranean grasses. II. Senescence, summer dormancy, and survival in *Phalaris*. *Australian Journal of Agricultural Research* **19**, 397-409.
- McWilliam, J.R. (1978). Response of pasture plants to temperature. In 'Plant Relations in Pastures.' (Ed. J.R. Wilson.) pp. 17-34. (Commonwealth Scientific and Industrial Research Organisation: Melbourne.)
- McWilliam, J.R. and Kramer, P.J. (1968). The nature of the perennial response in Mediterranean grasses. I. Water relations and summer survival in *Phalaris*.

Australian Journal of Agricultural Research **19**, 381-95.

- Mitchell, K.J. (1953a). Influence of light and temperature on the growth of ryegrass (*Lolium* spp.). I. Pattern of vegetative development. *Physiologia Plantarum* **6**, 21-46.
- Mitchell, K.J. (1953b). Influence of light and temperature on the growth of ryegrass (*Lolium* spp.). II. The control of lateral bud development. *Physiologia Plantarum* **6**, 425-43.
- Mitchell, K.J. (1956). Growth of pasture species under controlled environment. I. Growth at various levels of constant temperature. *New Zealand Journal of Science and Technology* **38**, 203-16.
- Moustafa, E., Ball, R. and Field, T.R.O. (1969). The use of acetylene reduction to study the effect of nitrogen fertiliser and defoliation on nitrogen fixation by field-grown white clover. *New Zealand Journal of Agricultural Research* **12**, 691-6.
- Murata, Y. and Iyama, J. (1963). Studies of the photosynthesis of forage crops. II. Influence of air temperature upon the photosynthesis of some forage and grain crops. *Proceedings of the Crop Science Society of Japan* **31**, 315-22.
- Murphy, P. (1982). Growth reserves in white clover. *An Foras Taluntais, Soils and Grassland Production Report*, p. 20.
- Newton, P.C.D. and Hay, M.J.M. (1994). Patterns of nodal rooting in *Trifolium repens* (L.) and correlations with stages in the development of axillary buds. *Grass and Forage Science* **49**, 270-6.
- Newton, P.C.D. and Hay, M.J.M. (1996). Clonal growth of white clover: factors influencing the viability of axillary buds and the outgrowth of a viable bud to form a branch. *Annals of Botany* **78**, 111-5.
- Newton, P.C.D., Hay, M.J.M., Thomas, V.J. and Dick, H.B. (1992). Viability of axillary buds of white clover (*Trifolium repens*) in grazed pasture. *Journal of Agricultural Science, Cambridge* **119**, 345-54.
- Nielsen, K.F. and Humphries, E.C. (1966). Effects of root temperature on plant growth. *Soils and Fertilisers* **29**, 1-7.
- Norris, I.B. (1981). Morphological variation in white clover. *Report of the Welsh Plant Breeding Station for 1980*, 120-1.

- Northcote, K.H. (1971). 'A Factual Key for the Recognition of Australian Soils.' Third Edition. (Rellim Technical Publications: Glenside.)
- Ong, C.K. (1978). The physiology of tiller death in grasses. 1. The influence of tiller age, size and position. *Journal of the British Grassland Society* **33**, 197-203.
- Ong, C.K. and Marshall, C. (1979). The growth and survival of severely-shaded tillers in *Lolium perenne* L. *Annals of Botany* **43**, 147-55.
- Ong, C.K., Marshall, C. and Sagar, G.R. (1978). The physiology of tiller death in grasses. 2. Causes of tiller death in a grass sward. *Journal of the British Grassland Society* **33**, 205-11.
- Oram, R.N. and Freebairn, R.D. (1984). Genetic improvement of drought survival ability in *Phalaris aquatica* L. *Australian Journal of Experimental Agriculture and Animal Husbandry* **24**, 403-9.
- Ostrowski, H. (1972). White clover (*Trifolium repens*) in subtropical Australia - a review. *Tropical Grasslands* **6**, 97-106.
- Oswalt, D.L., Bertrand, A.R. and Teel, M.R. (1959). Influence of nitrogen fertilisation and clipping on grass roots. *Proceedings of the Soil Science Society of America* **23**, 228-30.
- Ourry, A., Bigot, J. and Boucaud, J. (1989a). Protein mobilisation from stubble and roots, and proteolytic activities during post-clipping regrowth of perennial ryegrass. *Journal of Plant Physiology* **134**, 298-303.
- Ourry, A., Boucaud, J. and Salette, J. (1988). Nitrogen mobilisation from stubble and roots during regrowth of defoliated perennial ryegrass. *Journal of Experimental Botany* **39**, 803-9.
- Ourry, A., Gonzalez, B., Bigot, J., Boucaud, J. and Salette, J. (1989b). Nitrogen and carbohydrate mobilisations during regrowth of defoliated *Lolium perenne* L. *Proceedings of the 16th International Grassland Congress, Nice*, 513-4.
- Parfitt, R.L. (1980). A note on the losses from a phosphate cycle under grazed pasture. *New Zealand Journal of Experimental Agriculture* **8**, 215-7.
- Parsons, A.J., Collett, B. and Lewis, J. (1984). Changes in the structure and physiology of a perennial ryegrass sward when released from a continuous stocking management: implications for the use of exclusion cages in continually stocked swards. *Grass and Forage Science* **39**, 1-9.

- Patterson, H.D. and Thompson, R. (1971). Recovery of interblock information when block sizes are unequal. *Biometrika* **58**, 545-54.
- Peacock, J.M. (1975). Temperature and leaf growth in *Lolium perenne*. II. The site of temperature perception. *Journal of Applied Ecology* **12**, 115-23.
- Percival, N.S. and McClintock, M.B. (1982). Role of grazing management in manipulating the balance of ryegrass and paspalum in pastures. *New Zealand Journal of Experimental Agriculture* **10**, 365-70.
- Piano, E. and Annicchiarico, P. (1995). Persistence of Ladino white clover ecotypes and its relationship with other agronomic traits. *Grass and Forage Science* **50**, 195-8.
- Pinxterhuis, I., Chapman, D.F. and Hay, M.J.M. (1993). Structure of white clover populations and plants in pastures grazed by beef cattle. *Proceedings of the 17th International Grassland Congress, Palmerston North*, 138-9.
- Pollock, C. (1992). Thermotolerance in tropical cereals. Agricultural Food Research Council Institute of Grassland and Environmental Research 1991 Report, p. 46.
- Pollock, C.J. and Jones, T. (1979). Seasonal patterns of fructan metabolism in forage grasses. *New Phytologist* **83**, 9-15.
- Pollock, C.J., Eagles, C.F., Howarth, C.J., Schunmann, P.H.D. and Stoddart, J.L. (1993). Temperature stress. In 'Plant Adaptation to Environmental Stress.' (Eds. L. Fowden, T. Mansfield and J. Stoddart.) pp. 109-32. (Chapman and Hall: London.)
- Pontis, H.G. and Del Campillo, E. (1985). Fructans. In 'Biochemistry of Storage Carbohydrates in Green Plants.' (Eds. P.M. Dey and R.A. Dixon) pp. 205-27. (Academic Press: London.)
- Prestidge, R.A., Thom, E.R., Marshall, S.L., Taylor, M.J., Willoughby, B. and Wildermoth, D.D. (1992). Influence of *Acremonium lolii* infection in perennial ryegrass on germination, emergence, survival, and growth of white clover. *New Zealand Journal of Agricultural Research* **35**, 225-34.
- Prud'homme, M.P., Gonzalez, B., Billard, J.P. and Boucaud, J. (1992). Carbohydrate content, fructose and sucrose enzyme activities in roots, stubble and leaves of ryegrass (*Lolium perenne* L.) as affected by source/sink modification after cutting. *Journal of Plant Physiology* **140**, 282-91.
- Raven, P.H., Evert, R.F. and Eichhorn, S.E. (1986). 'Biology of Plants.' (Eds. S. Anderson and G. Hefta.) Fourth Edition. (Worth Publishers Inc.: New York.)

- Reason, G.K., Bodero, J., Lowe, K.F., Rayment, G.K. and McGuigan, K. (1989). Milk production from nitrogen fertilised raingrown grass pastures. Queensland Department of Primary Industries Final Report DAQ49. (Queensland Department of Primary Industries: Brisbane.)
- Reid, D. (1959). Studies on the cutting management of grass-clover swards. 1. The effects of varying the closeness of cutting on the yields from an established grass-clover sward. *Journal of Agricultural Science* **53**, 299-312.
- Reid, D. (1962). Studies on the cutting management of grass-clover swards. III. The effects of prolonged close and lax cutting on herbage yields and quality. *Journal of Agricultural Science* **59**, 359-68.
- Richards, J.H. (1993). Physiology of plants recovering from defoliation. In 'Grasslands for Our World.' (Ed. M.J. Barker.) pp. 46-54. (SIR Publishing: New Zealand.)
- Richards, J.H. and Caldwell, M.M. (1985). Soluble carbohydrates, concurrent photosynthesis and efficiency in regrowth following defoliation: a field study with *Agropyron* species. *Journal of Applied Ecology* **22**, 907-20.
- Robin, Ch., Hay, M.J.M. and Newton, P.C.D. (1994b). Effect of light quality (red: far-red ratio) and defoliation treatments applied at a single phytomer on axillary bud outgrowth in *Trifolium repens* L. *Oecologia* **100**, 236-42.
- Robin, Ch., Hay, M.J.M., Newton, P.C.D. and Greer, D.H. (1994a). Effect of light quality (red: far-red ratio) at the apical bud of the main stolon on morphogenesis of *Trifolium repens* L. *Annals of Botany* **74**, 119-23.
- Robson, M.J. (1973a). The growth and development of simulated swards of perennial ryegrass. I. Leaf growth and dry weight changes as related to the ceiling yield of a seedling sward. *Annals of Botany* **37**, 487-500.
- Robson, M.J. (1973b). The growth and development of simulated swards of perennial ryegrass. II. Carbon assimilation and respiration in a seedling sward. *Annals of Botany* **37**, 501-18.
- Robson, M.J. (1981). Potential production - what is it and can we increase it? In 'Plant Physiology and Herbage Production.' (Ed. C.E. Wright.) pp. 5-18. (British Grassland Society: Great Britain)
- Rosa, J.T. (1921). Investigations on the hardening process in vegetable plants. Research Bulletin 48, University of Missouri College of Agriculture.

- Roush, M.L. and Radosevich, S.R. (1985). Relationships between growth and competitiveness of four annual weeds. *Journal of Applied Ecology* **22**, 895-905.
- Ryan, B.F., Joiner, B.L. and Ryan, T.A. (1985). 'Minitab Handbook.' Second Edition. (PWS-KENT Publishing Company: Boston.)
- Ryle, G.J.A. (1964). A comparison of leaf and tiller growth in several perennial grasses as influenced by nitrogen and temperature. *Journal of the British Grassland Society* **19**, 281-90.
- Ryle, G.J.A. and Powell, C.E. (1975) Defoliation and regrowth in the graminaceous plant: the role of current assimilate. *Annals of Botany* **39**, 297-310.
- Sandli, N., Svenning, M.M., Rosnes, K. and Junttila, O. (1993). Effect of nitrogen supply on frost resistance, nitrogen metabolism and carbohydrate content in white clover (*Trifolium repens*). *Physiologia Plantarum* **88**, 661-7.
- Sarathchandra, S.U., DiMenna, M.E., Burch, G., Brown, J.A., Watson, R.N., Bell, N.L. and Cox, N.R. (1995). Effects of plant-parasitic nematodes and rhizosphere microorganisms on the growth of white clover (*Trifolium repens* L.) and perennial ryegrass (*Lolium perenne* L.). *Soil Biology and Biochemistry* **27**, 9-16.
- Searle, S.R. (1971). 'Linear Models.' (John Wiley and Sons: New York.)
- Silsbury, J.H. (1961). A study of dormancy, survival, and other characteristics in *Lolium perenne* L. at Adelaide, S.A. *Australian Journal of Agricultural Research* **12**, 1-9.
- Silsbury, J.H. (1964). Tiller dynamics, growth and persistency of *Lolium perenne* L. and of *Lolium rigidum* Gaud. *Australian Journal of Agricultural Research* **15**, 9-20.
- Silsbury, J.H. (1970). Leaf growth in pasture grasses. *Tropical Grasslands* **4**, 17-36.
- Simons, R.G., Davies, A. and Troughton, A. (1972). The effect of the height of defoliation on two clones of perennial ryegrass. *Journal of Agricultural Science, Cambridge* **79**, 509-14.
- Singh, D.K. and Sale, P.W.G. (1997a). Defoliation frequency and the response by white clover to increasing phosphorus supply. 2. Non-structural carbohydrate concentrations in plant parts. *Australian Journal of Agricultural Research* **48**, 119-24.
- Singh, D.K. and Sale, P.W.G. (1997b). Defoliation frequency and the response by white clover to increasing phosphorus supply. 1. Leaf dry matter yield and plant morphology responses. *Australian Journal of Agricultural Research* **48**, 111-8.

- Smetham, M.L. (1973). Grazing management. In 'Pastures and Pasture Plants.' (Ed. R.H.M. Langer.) pp. 179-228. (A.H. and A.W. Reed: Wellington.)
- Smith, D. (1969). Removing and analysing total non-structural carbohydrates from plant tissue. *Wisconsin Agricultural Experimental Station, Research Report* **41**, 1-11.
- Smith, D. (1970). Influence of cool and warm temperatures and temperature reversal at inflorescence emergence on yield and chemical composition of timothy and brome grass at anthesis. *Proceedings of the 11th International Grassland Congress, Surfers Paradise*, 510-4.
- Smith, D. (1972). Carbohydrate reserves of grasses. In 'The Biology and Utilisation of Grasses.' (Eds. V.B. Younger and C.M. McKell.) pp. 318-33. (Academic Press Inc.: New York.)
- Smith, D. (1973). The nonstructural carbohydrates. In 'Chemistry and Biochemistry of Herbage.' Volume 1. (Eds. G.W. Butler and R.W. Bailey.) pp. 105-55. (Academic Press Inc.: New York.)
- Smith, D. (1974). Growth and development of timothy tillers as influenced by level of carbohydrate reserves and leaf area. *Annals of Botany* **38**, 595-606.
- Smith, D. and Jewiss, O.R. (1966). Effects of temperature and nitrogen supply on the growth of timothy (*Phleum pratense* L.). *Annals of Applied Biology* **58**, 145-57.
- Smouter, H., Simpson, R.J. and Pearce, G.R. (1995). Water-soluble carbohydrates and *in vitro* digestibility of annual ryegrass (*Lolium rigidum* Gaudin) sown at varying densities. *Australian Journal of Agricultural Research* **46**, 611-25.
- Smucker, A.J.M., McBurney, S.L. and Srivastava, A.K. (1982). Quantitative separation of roots from compacted soil profiles by the hydropneumatic elutriation system. *Agronomy Journal* **74**, 500-3.
- Spedding, C.R.W. (1971). Grassland Ecology. (Oxford University Press: New Zealand.)
- Sprague, V.G. and Sullivan, J.T. (1950). Reserve carbohydrates in orchard grass clipped periodically. *Plant Physiology* **25**, 92-102.
- Stevenson, C.A. and Laidlaw, A.S. (1985). The effect of moisture stress on stolon and adventitious root development in white clover (*Trifolium repens* L.) *Plant and Soil* **85**, 249-57.
- Stuckey, I.H. (1941). Seasonal growth of grass roots. *American Journal of Botany* **28**, 486-91.

- Sullivan, J.T. and Sprague, V.G. (1943). Composition of the roots and stubble of perennial ryegrass following partial defoliation. *Plant Physiology* **18**, 656-70.
- Sullivan, J.T. and Sprague, V.G. (1949). The effect of temperature on the growth and composition of the stubble and roots of perennial ryegrass. *Plant Physiology* **24**, 706-19.
- Tallowin, J.R.B., Kirkham, F.W. and Brookman, S.K.E. (1986). Sward damage by sod-pulling - the effect of nitrogen. In 'Grazing.' (Ed. J. Frame.) pp. 44-8. (British Grassland Society: Great Britain.)
- Thom, E.R. (1991). Effect of early spring grazing frequency on the reproductive growth and development of a perennial ryegrass tiller population. *New Zealand Journal of Agricultural Research* **34**, 383-9.
- Thom, E.R., Sheath, G.W. and Bryant, A.M. (1989). Seasonal variations in total nonstructural carbohydrate and major element levels in perennial ryegrass and paspalum in a mixed pasture. *New Zealand Journal of Agricultural Research* **32**, 157-65.
- Thom, E.R., Sneath, G.W., Bryant, A.M. and Cox, N.R. (1986). Renovation of pastures containing paspalum. 3. Effect of defoliation management and irrigation on ryegrass growth and persistence. *New Zealand Journal of Agricultural Research* **29**, 599-611.
- Thomas, H. (1991). Accumulation and consumption of solutes in swards of *Lolium perenne* during drought and after rewatering. *New Phytologist* **118**, 35-48.
- Troughton, A. (1981). Length of life of grass roots. *Grass and Forage Science* **36**, 117-20.
- Turner, N.C. and Begg, J.E. (1978). Responses of pasture plants to water deficits. In 'Plant Relations in Pastures.' (Ed. J.R. Wilson.) pp. 50-66. (Commonwealth Scientific and Industrial Research Organisation: Melbourne.)
- Van den Bosch, J. and Mercer, C.F. (1996a). Variation in white clover for root-knot nematode resistance. *New Zealand Journal of Agricultural Research* **39**, 137-48.
- Van den Bosch, J. and Mercer, C.F. (1996b). Third generation progress in breeding white clover for resistance to root-knot nematode. In 'White Clover: New Zealand's Competitive Edge.' (Ed. D.R. Woodfield.) pp. 163-6. (Agronomy Society of New Zealand: Christchurch.)

- Van Loo, E.N. (1992). Tillering, leaf expansion and growth of plants of two cultivars of perennial ryegrass grown using hydroponics at two water potentials. *Annals of Botany* **70**, 511-8.
- Vartha, E.W and Bailey, R.W. (1980). Soluble carbohydrate content and composition in above- and below-ground tissues of Westerwolds ryegrass during winter. *New Zealand Journal of Agricultural Research* **23**, 93-6.
- Vine, D.A. (1983). Sward structure changes within a perennial ryegrass sward: leaf appearance and death. *Grass and Forage Science* **38**, 231-42.
- Volaire, F. (1994a). Accumulation of water soluble carbohydrates in young plants of two contrasting populations of *Dactylis glomerata* L. subjected to water-stress. *Proceedings of the 3rd Congress of the European Society of Agronomy*, 272-3.
- Volaire, F. (1994b). Effects of summer drought and spring defoliation on carbohydrate reserves, persistence and recovery of two populations of cocksfoot (*Dactylis glomerata*) in a Mediterranean environment. *Journal of Agricultural Science, Cambridge* **122**, 207-15.
- Volaire, F. (1995). Growth, carbohydrate reserves and drought survival strategies of contrasting *Dactylis glomerata* populations in a Mediterranean environment. *Journal of Applied Ecology* **32**, 56-66.
- Volaire, F. and Gandoin, J.M. (1996). The effect of age of the sward on the relationship between water-soluble carbohydrate accumulation and drought survival in two contrasted populations of cocksfoot (*Dactylis glomerata* L.). *Grass and Forage Science* **51**, 190-8.
- Volaire, F. and Thomas, H. (1995). Effects of drought on water relations, mineral uptake, water-soluble carbohydrate accumulation and survival of two contrasting populations of cocksfoot (*Dactylis glomerata* L.). *Annals of Botany* **75**, 513-24.
- Volenc, J.J. (1986). Nonstructural carbohydrates in stem base components of tall fescue during regrowth. *Crop Science* **26**, 122-7.
- Volenc, J.J., Ourry, A. and Joern, B.C. (1996). Minireview: A role for nitrogen reserves in forage regrowth and stress tolerance. *Physiologia Plantarum* **97**, 185-93.
- Waite, R. and Boyd, J. (1953). The water-soluble carbohydrates of grasses. I. Changes occurring during the normal life-cycle. *Journal of the Science of Food and Agriculture* **4**, 197-204.

- Ward, C.Y. and Blaser, R.E. (1961). Carbohydrate food reserves and leaf area in regrowth of orchard grass. *Crop Science* **1**, 366-70.
- Watson, R.N., Harris, S.L., Bell, N.L. and Neville, F.J. (1993). Pasture pests reduce white clover performance. *Proceedings of the 45th Ruakura Dairy Farmer's Conference*, 57-61.
- Watson, R.N., Harris, S.L., Bell, N.L. and Neville, F.J. (1996). Deferred grazing to enhance white clover content in pastures. In 'White Clover: New Zealand's Competitive Edge.' (Ed. D.R. Woodfield.) p. 154. (Agronomy Society of New Zealand: Christchurch.)
- Wehner, D.J. and Watschke, T.L. (1981). Heat tolerance of Kentucky bluegrass, perennial ryegrass, and annual bluegrass. *Agronomy Journal* **73**, 79-84.
- Weinmann, H. (1948). Underground development and reserves of grasses: a review. *Journal of the British Grassland Society* **3**, 115-40.
- Weinmann, H. (1952). Carbohydrate reserves in grasses. *Proceedings of the 6th International Grassland Congress. Pennsylvania* **1**, 655-60.
- White, L.M. (1973). Carbohydrate reserves of grasses: a review. *Journal of Range Management* **26**, 13-8.
- Williams, P.H., Gregg, P.E.H. and Hedley, M.J. (1990). Mass balance modelling of potassium losses from grazed dairy pasture. *New Zealand Journal of Agricultural Research* **33**, 661-8.
- Wilman, D. and Acunap, G.M. (1993). Effects of cutting height on the growth of leaves and stolons in perennial ryegrass - white clover swards. *Journal of Agricultural Science, Cambridge* **21**, 39-46.
- Wilman, D. and Asiegbu, J.E. (1982a). The effects of clover variety, cutting interval and nitrogen application on herbage yields, proportions and heights in perennial ryegrass-white clover swards. *Grass and Forage Science* **37**, 1-13.
- Wilman, D. and Asiegbu, J.E. (1982b). The effects of variety, cutting interval and nitrogen application on the morphology and development of stolons and leaves of white clover. *Grass and Forage Science* **37**, 15-27.
- Wilman, D. and Fisher, A. (1996). Effects of interval between harvests and application of fertiliser N in spring on the growth of perennial ryegrass in a grass/white clover sward. *Grass and Forage Science* **51**, 52-7.

- Wilson, D. (1975). Variation in leaf respiration in relation to growth and photosynthesis of *Lolium*. *Annals of Applied Biology* **80**, 323-38.
- Wilson, J.R. and Ford, C.W. (1971). Temperature influences on the growth, digestibility, and carbohydrate composition of two tropical grasses, *Panicum maximum* var. *Trichoglume* and *Setaria sphacelata*, and two cultivars of the temperate grass *Lolium perenne*. *Australian Journal of Agricultural Research* **22**, 563-71.
- Woledge, J. (1977). The effects of shading and cutting treatments on the photosynthetic rates of ryegrass leaves. *Annals of Botany* **41**, 1279-86.
- Yarrow, N.H. and Penning, P.D. (1974). Managing grass-clover swards to produce differing clover proportions. *Grass and Forage Science* **49**, 496-501.