

## REFERENCES

- Abyzov, S. S. (1993). Microorganisms in the antarctic ice. In *Antarctic Microbiology*. pp. 265-295. Edited by Friedman, E. I. Wiley-Liss, New York.
- Achstetter, T., Ehmann, C., Osaki, A. and Wolf, D. H. (1984). Proteolysis in eukaryotic cells: Proteinase yscE, a new yeast peptidase. *J. Biol. Chem.* **259**, 13344-13348.
- Aebi, H. (1984). Catalase *in vitro*. *Meth. Enzymol.* **105**, 121-126.
- Aguilar, A. (1996). Extremophile research in the European Union: from fundamental aspects to industrial expectations. *FEMS Microbiol. Rev.* **18**, 89-92.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. (1990). Basic local alignment search tool. *J. Mol. Biol.* **215**, 403-410.
- App, H. and Holzer, H. (1989). Purification and characterization of neutral trehalase from the yeast *ABYS 1* mutant. *J. Biol. Chem.* **264**, 17583-17588.
- Araki, T. (1992). An analysis of the effect of changes in growth temperature on proteolysis *in vivo* in the psychrophilic bacterium *Vibrio* sp. Strain ANT-300. *J. Gen. Microbiol.* **138**, 2075-2082.
- Argüelles, J. C. (1994). Heat-shock response in a yeast tps1 mutant deficient in trehalose synthesis. *FEBS Lett.* **350**, 266-270.
- Argüelles, J. C. (1997). Thermotolerance and trehalose accumulation induced by heat shock in yeast cells of *Candida albicans*. *FEMS Lett.* **146**, 65-71.
- Arthur, H. (1979). Psychrophilic and thermophilic yeasts – Biochemical and ultrastructural studies. *PhD thesis* Department of Chemistry and Biochemistry, James Cook University, North Queensland, Australia
- Arthur, H. and Watson, K. (1976). Thermal adaptation in yeast: growth temperatures, membrane lipid, and cytochrome composition of psychrophilic, mesophilic and thermophilic yeasts. *J. Bacteriol.* **128**, 56-68.
- Arthur, H., Watson, K., McArthur, C. R. and Clark-Walker, G. D. (1978). Naturally occurring respiratory deficient *Candida slooffii* strains resemble petite mutants. *Nature* **271**, 750-752.
- Aruoma, O. I., Halliwell, B. and Dizdaroglu, M. (1989). Iron ion-dependent modification in bases in DNA by superoxide radical-generating system hypoxanthine/xanthine oxidase. *J. Biol. Chem.* **264**, 13024-13028.

- Ashburner, M. and Bonner, J. J. (1979). The induction of gene activity in *Drosophila* by heat shock. *Cell* **17**, 241-254.
- Atencio, D. and Yaffe, M. (1992) MAS5, a yeast homolog of DnaJ involved in mitochondrial import. *Mol. Cell. Biol.* **12**, 283-291.
- Attfield, P. V. (1987). Trehalose accumulates in *Saccharomyces cerevisiae* during exposure to agents that induce the heat shock response. *FEBS Lett.* **225**, 259-263.
- Attfield, P. V., Kletsas, S. and Hazell, B. W. (1994). Concomitant appearance of intrinsic thermotolerance and storage of trehalose in *Saccharomyces cerevisiae* during early respiratory phase of batch-culture is *CIF1*-dependent. *Microbiol.* **140**, 2625-2632.
- Attfield, P. V., Raman, A. and Northcott, C. J. (1992). Construction of *Saccharomyces cerevisiae* strains that accumulate relatively low concentrations of trehalose, and their application in testing the contribution of the disaccharide to stress tolerance. *FEMS Lett.* **94**, 271-276.
- Auffray, Y., Lecesne, E., Hartke, A. and Bcuitbonnes, P. (1995). Basic features of the *Streptococcus thermophilus* heat shock response. *Curr. Microbiol.* **30**, 87-91.
- Ausubel, F., Brent, R., Kingston, R., Moore, D. D., Seidman, J. G., Smith, J. A. and Stuhl, K. (eds) (1988). *Current Protocols in Molecular Biology*. Wiley-Interscience, New York.
- Bachair, A., Finley, D. and Varshavsky, A. (1986). In vivo half-life of a protein is a function of its amino-terminal residue. *Science* **234**, 179-186.
- Barnes, C. A., Johnson, G. C. and Singer, R. A. (1990). Thermotolerance is independent of induction of the full spectrum of heat shock proteins of cell cycle blockage in the yeast *Saccharomyces cerevisiae*. *J. Bacteriol.* **172**, 4352-4358.
- Barnett, J. A., Payne, R. W. and Yarrow, D (1990). *Yeast: Characteristics and identification*. 2nd edn. Cambridge University Press, Cambridge.
- Barraclough, R. and Ellis, R. J. (1980). Proein synthesis in chloroplasts. XI. Assembly of newly-synthesized large subunits of ribulose bisphosphate carboxylase in intact isolated pea chloroplasts. *Biochem. Biophys. Acta* **608**, 18-31.
- Beavan, M. J., Charpentier, C. and Rose, A. H. (1982). Production and tolerance of ethanol in relation to phospholipid fatty-acyl composition in *Saccharomyces cerevisiae* NCYC 431. *J. Gen. Microbiol.* **128**, 1447-1455.
- Becker, J. and Craig, E. A. (1994). Heat-shock proteins as molecular chaperones. *Eur. J. Biochem.* **219**, 11-23.

- Belazzi, T., Wagner, A., Wieser, R., Schanz, M., Adam, G., Hartig, A. and Ruis, H. (1991). Negative regulation of transcription of the *Saccharomyces cerevisiae* catalase T (CTT1) gene by cAMP is mediated by a positive control element. *EMBO J.* **10**, 585-592.
- Bell, W., Klaassen, P., Ohnacker, M., Boller, T., Herweijer, M., Schoppoml, P., van der Zee, P. and Wiemken, A. (1992). Characterization of the 56-kDa subunit of yeast trehalose-6-phosphate synthase and cloning of its gene reveal its identity with the product of CIF1, a regulator of carbon catabolite inactivation. *Eur. J. Biochem.* **209**, 951-959.
- Bentley, N. J., Fitch, I. T. and Tuite, M. F. (1992). The small heat-shock protein Hsp26 of *Saccharomyces cerevisiae* assembles into a high molecular weight aggregate. *Yeast* **8**, 95-106.
- Berg, G. R., Inniss, W. E. and Heikkila, J. J. (1987). Stress proteins and thermotolerance in psychrotrophic yeasts from arctic environments. *Can. J. Microbiol.* **33**, 383-389.
- Bienz, M. (1984). Developmental control of the heat shock response in *Xenopus*. *Proc. Natl. Acad. Sci. USA* **81**, 3138-3142.
- Bienz, M. and Pelham, H. R. B. (1987). Mechanisms of heat-shock gene activation in higher eukaryotes. *Adv. Genet.* **34**, 31-72.
- Bilinski, T., Krawiek, Z., Liczmanski, A., and Litwinska, J. (1985). Is hydroxyl radical generated by the Fenton reaction *in vivo*? *Biochem. Biophys. Res. Comm.* **130**, 533-539.
- Birnboim, H. C. and Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nuc. Acids Res.* **7**, 1513-1523.
- Blumberg, H. and Silver, P. A. (1991). A homologue of the bacterial heat-shock gene *DnaJ* that alters protein sorting in yeast. *Nature* **349**, 627-630.
- Bond, U. and Schlesinger, M. J. (1987). Heat shock proteins and development. *Adv. Genet.* **24**, 1-29.
- Boorstein, W. R. and Craig, E. A. (1990). Regulation of a yeast HSP70 gene by a cAMP responsive transcriptional control element. *EMBO J.* **9**, 2543-2553.
- Boorstein, W. R., Ziegelhoffer, T. and Craig, E. A. (1994). Molecular evolution of the HSP70 multigene family. *J. Mol. Evolution* **38**, 1-17.
- Borkovich, K. A., Farrelly, F. W., Finkelstein, D. B., Taulien, J. and Lindquist, S. (1989). Hsp82 is an essential protein that is required in higher concentration for growth of cells at higher temperatures. *Mol. Cell. Biol.* **9**, 3919-3930.

- Bose, S., Weikl, T., Bügl, H. and Bucher, J. (1996). Chaperone function of Hsp90-associated proteins. *Science* **274**, 1715-1717.
- Bossier, P., Fernandes, L., Rocha, D. and Rodrigues-Pousada, C. (1993). Overexpression of YAP2, coding for a new yAP protein, and YAP1 in *Saccharomyces cerevisiae* alleviates growth inhibition caused by 1, 10-phenanthroline. *J. Biol. Chem.* **268**, 23640-23645.
- Bossier, P., Fitch, I. T., Boucherie, H. and Tuite, M. F. (1989). Structure and expression of a yeast gene encoding the small heat-shock protein Hsp26. *Gene* **78**, 323-330.
- Boucherie, H., Bataille, N., Fitch, I. T., Periot, M. and Tuite, M. F. (1995). Differential synthesis of glyceraldehyde-3-phosphate dehydrogenase polypeptides in stressed yeast cells. *FEMS Lett.* **125**, 127-134.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of micogram quantities of protein utilizing the principle of protein dye-binding. *Anal. Biochem.* **72**, 248-254.
- Breimer, L. H. (1988). Ionizing radiation-induced mutation. *Br. J. Cancer* **57**, 6-18.
- Bridson, E. Y. (1990). *The Oxoid Manual*. pp. 2-5. Unipath Ltd., Basingstoke, England.
- Brock, T. D. (1978). *Thermophilic Microorganisms and Life at High Temperatures*. Springer-Verlag, New York.
- Broeze, R. J., Solomon, C. J. and Pope, D. H. (1978). Effects of low temperature on in vivo and in vitro protein synthesis in *Escherichia coli* and *Pseudomonas fluorescens*. *J. Bacteriol.* **134**, 861-874.
- Brot, N., Weissbach, L., Werth, J. and Weissbach, H. (1981). Enzymatic reduction of protein-bound methionine sulfoximide. *Proc. Natl. Acad. Sci. USA* **78**, 2155-2158.
- Brugge, J. S. (1986). Interaction of the Rous sarcoma virus protein, pp60<sup>src</sup>, with the cellular protein pp50 and pp90. *Curr. Topics Microbiol. Immunol.* **123**, 1-22.
- Burdon, R. H., Gill, V. and Evans, C. R. (1990). Active oxygen species and heat shock protein induction. In *Stress Proteins – Induction and Function*. pp. 19-25. Edited by Schlesinger, M. J., Santoro, M. G. and Caraci, E. Springer-Verlag, Berlin.
- Bush, G. L. and Meyer, D. I. (1996). The refolding activity of the yeast heat shock protein Ssa1 and Ssa2 defines their role in protein translocation. *J. Cell. Biol.* **135**, 1229-1237.
- Cabib, E. and Leloir, F. L. (1958). The biosynthesis of trehalose-6-phosphate. *J. Biol. Chem.* **231**, 259-275.

- Cairns, A., Howarth, C. J. and Pollock, C. J. (1995). Submerged batch culture of the psychrophile *Monographella nivalis* in a defined medium: growth, carbohydrate utilization and responses to temperature. *New Phytol.* **129**, 299-308.
- Cameron, R. E. (1974). Application of low-latitude microbial ecology to high-latitude deserts. In *Polar Deserts and Modern Man*. pp. 71-90. Edited by Smiley, T. L. and Zumberge, J. H. University of Arizona Press, Tucson.
- Cameron, S., Levin, L., Zoller, M. and Wigler, M. (1988). cAMP-independent control of sporulation, glycogen metabolism and heat shock resistance in *Saccharomyces cerevisiae*. *Cell* **53**, 555-566.
- Caplan, A., Cyr, D. and Douglas, M. (1992). YDJ1p facilitated polypeptide translocation across different intracellular membranes by a conserved mechanism. *Cell* **71**, 1143-1155.
- Caplan, A. and Douglas, M. (1991). Characterization of *YDJ1*: A yeast homologue of the bacterial *dnaJ* protein. *J. Cell. Biol.* **114**, 609-621.
- Carmognol, F., Sinet, P. M. and Jerome, H. (1983). Selenium-dependent and non-dependent glutathione peroxidase in human tissue extracts. *Biochim. Biophys. Acta* **759**, 49-57.
- Carpenter, J. F. (1993). Stabilization of proteins during freezing and dehydration. Application of lessons from nature. *Cryobiology* **30**, 220-221.
- Carpenter, J. F., Martin, B., Crowe, L. M. and Crowe, J. H. (1987). Stabilization of phosphofructokinase during air-drying with sugars and sugar/transition metal mixtures. *Cryobiology* **24**, 455-467.
- Carri, M. T., Galiazzo, F., Ciriolo, M. R. and Rotilio, G. (1991). Evidence for co-regulation of Cu,Zn superoxide dismutases and metallothionein gene expression in yeast through transcriptional control by copper via the ACE1 factor. *FEBS Lett.* **278**, 263-266.
- Cavicchi, S., Guerra, D., La Torre, V. and Huey, R. B. (1995). Chromosomal analysis of heat-shock tolerance in *Drosophila melanogaster* evolving at different temperatures in the laboratory. *Evolution* **49**, 676-684.
- Chang, H. C. and Lindquist, S. (1994). Conservation of Hsp90 macromolecular complexes in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **269**, 24983-24988.
- Chang, H. C., Nathan, D. F. and Lindquist, S. (1997). In vivo analysis of the *HSP90* cochaperone Sti1(p60). *Mol. Cell. Biol.* **17**, 318-325.

- Chau, V., Tobias, J. W., Bachmair, A., Mariot, D., Ecker, D. J., Gonda, D. K. and Varshavsky, A. (1989). A multiubiquitin chain is confined to specific lysine in a targeted short-lived protein. *Science* **243**, 1576-1583.
- Chen, Y. and Piper, P. W. (1995). Consequence of the overexpression of ubiquitin in yeast: Elevated tolerances of osmostress, ethanol and canavanine, yet reduced tolerances of cadmium, arsenite and paromycin. *Biochim. Biophys. Acta* **1268**, 59-64.
- Cheng, M. Y., Hartl, F.-V., Martin, J., Pollock, R. A., Kalousek, F., Neupert, W., Hallberg, E. M., Hallberg, R. L. and Horwich, A. I. (1989). Mitochondrial heat-shock protein hsp 60 is essential for assembly of proteins imported into yeast mitochondria. *Nature* **337**, 620-625.
- Cheng, L., Hirst, K. and Piper, P. (1992). Authentic temperature-regulation of a heat shock gene inserted into yeast on a high copy number vector. Influences of overexpression of HSP90 protein on high temperature growth and thermotolerance. *Biochim. Biophys. Acta* **1132**, 26-34.
- Christman, M. F., Morgan, R. W., Jacobsen, F. S. and Ames, B. N. (1985). Positive control of a regulon for defences against oxidative stress and some heat shock proteins in *Salmonella typhimurium*. *Cell* **41**, 753-762.
- Clark, M. W., Yip, M. L. R., Campbell, J. and Abelson, J. (1990). SSB-1 of the yeast *Saccharomyces cerevisiae* is a nucleolar-specific, silver-binding protein that is associated with the snR10 and snR11 small nuclear RNAs. *J. Cell. Biol.* **111**, 1741-1751.
- Cohen, I. R. and Young, D. B. (1991). Autoimmunity, microbial immunity and the immunological homunculus. *Immunol. Today* **12**, 105-110.
- Cohen, S. N., Chang, A. C. Y. and Hsu, L. (1972). Nonchromosomal antibiotic resistance in bacteria: Genetic transformation of *Escherichia coli* by R-factor DNA. *Proc. Natl. Acad. Sci. USA* **69**, 2110-2114.
- Cole, M. B. and Keenan, M. J. H. (1986). Synergistic effects of weak acid preservatives and pH on the growth of *Zygosaccharomyces bailii*. *Yeast* **2**, 93-100.
- Collinson, L. P. and Dawes, I. W. (1992). Inducibility of the response of yeast cells to peroxide stress. *J. Gen. Microbiol.* **138**, 329-335.
- Cooney, D. G. and Emerson, R. (1964). *Thermophilic fungi*. W. H. Freeman, San Francisco.

- Coote, P. J., Jones, M. V., Edgar, K. and Cole, M. B. (1992). *TPK* gene products mediate cAMP-independent thermotolerance in *Saccharomyces cerevisiae*. *J. Gen. Microbiol.* **138**, 2551-2557.
- Coote, P. J., Jones, M. V., Seymour, I. J., Rowe, D. L., Ferdinando, D. P., McArthur, A. J. and Cole, M. B. (1994). Activity of the plasma membrane H<sup>+</sup>-ATPase is a key physiological determinant of thermotolerance in *Saccharomyces cerevisiae*. *Microbiol.* **140**, 1881-1890.
- Costa, V., Reis, E., Quintanilha, A., and Moradas-Ferreira, P. (1993). Acquisition of ethanol tolerance in *Saccharomyces cerevisiae*: the key role of the mitochondrial superoxide dismutase. *Arch. Biochem. Biophys.* **300**, 608-614.
- Craig, E. A. (1989). Essential roles of 70 kDa heat inducible proteins. *BioEssays* **11**, 48-52.
- Craig, E. A. (1992). The heat-shock response of *Saccharomyces cerevisiae*. In *The molecular and cellular biology of the yeast Saccharomyces cerevisiae: Gene expression*. Vol. II. pp. 501-537. Edited by Jones, E. W., Pringle, J. R. and Broach, J. R. Cold Spring Harbor Laboratory Press, New York.
- Craig, E. A., Baxter, B. K., Becker, J., Halladay, J. and Ziegelhoffer, T. (1994). Cytosolic hsp70s of *Saccharomyces cerevisiae*: Roles in protein synthesis, posttranslocation, proteolysis, and regulation. In *The Biology of Heat Shock Proteins and Molecular Chaperones*. pp. 31-52. Edited by Morimoto, R. I., Tissières, A. and Georgopoulos, C. Cold Spring Harbor Laboratory Press, Cold Spring Harbor.
- Craig, E. A., Gambill, B. D. and Nelson, J. (1993). Heat shock proteins: Molecular chaperones of protein biogenesis. *Microbiol. Rev.* **57**, 402-414.
- Craig, E. A. and Gross, C. A. (1991). Is hsp70 the cellular thermometer? *Trends Biochem. Sci.* **16**, 135-140.
- Craig, E. A. and Jacobsen, K. (1984). Mutations of the heat inducible 70 kilodalton genes of yeast confer temperature sensitive growth. *Cell* **38**, 841-849.
- Craig, E. A. and Jacobsen, K. (1985). Mutations in cognate genes of *Saccharomyces cerevisiae* hsp70 result in reduced growth rates at low temperatures. *Mol. Cell. Biol.* **5**, 3517-3524.
- Craig, E. A., Kang, P. J. and Boorstein, W. (1990). A review of the role of 70 kDa heat shock proteins in protein translocation across membranes. *Anton. van Leeuwenhoek* **58**, 191-193.

- Craig, E. A., Kramer, J. and Kosic-Smithers, J. (1987). SSC1, a member of the 70-kDa heat shock protein multigene family of *Saccharomyces cerevisiae*, is essential for growth. *Proc. Natl. Acad. Sci. USA* **84**, 4156-4160.
- Daniels, C. J., McKee, A. H. Z. and Doolittle, W. F. (1984). Archaeabacterial heat-shock proteins. *EMBO J.* **3**, 745-749.
- Davidson, J. F., Whyte, B., Bissinger, P. H. and Schiestl, R. H. (1996). Oxidative stress is involved in heat-induced cell death in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* **93**, 5116-5121.
- Davies, K. J. A. (1986). Intracellular proteolytic systems may function as secondary antioxidant defenses: a hypothesis. *J. Free Rad. Biol. Med.* **2**, 155-173.
- Davies, K. J. A. and Lin, S. W. (1988). Degradation of oxidatively denatured proteins in *Escherichia coli*. *J. Bacteriol.* **168**, 1120-1127.
- Dean, R. T. and Pollak, J. K. (1985). Endogenous free radical generation may influence proteolysis in mitochondria. *Biochem. Biophys. Res. Comm.* **126**, 1082-1089.
- Deegenaars, M. L. (1991). Preliminary characterization of the heat shock response in the thermophilic yeast, *Candida pintolopesii* and the psychrophilic yeast, *Candida psychrophila*. Honours thesis, University of New England, Armidale, Australia.
- Deegenaars, M. L. and Watson, K. (1992). Characterization of the stress response in yeast from extreme environments. Proc. 16th Int. Conf. on Yeast Genetic and Mol. Biol.. Vienna, 14-9B.
- Deegenaars, M. L. and Watson, K. (1997). Stress proteins and stress tolerance in an Antarctic psychrophilic yeast, *Candida psychrophila*. *FEMS Lett.* **151**, 191-196.
- Deshaines, R. J., Koch, B. D., Werner-Washburne, M., Craig, E. A., Schekman, R. (1988). A subfamily of stress proteins facilitates translocation of secretory and mitochondrial precursor polypeptides. *Nature* **332**, 800-805.
- Destruelle, M., Holzer, H. and Klionsky, D. J. (1994). Identification and characterization of a novel yeast gene: the *YGP1* gene product is a highly glycosylated secreted protein that is synthesized in response to nutrient limitation. *Mol. Cell. Biol.* **14**, 2740-2754.
- Destruelle, M., Holzer, H. and Klionsky, D. J. (1995). Isolation and characterization of a novel yeast gene, *ATH1*, that is required for vacuolar and trehalose activity. *Yeast* **11**, 1015-1025.
- De Virgilio, C., Burckert, N., Bell, W., Jeno, P., Boller, T. and Wiemken, A. (1993). Disruption of *TPS2*, the gene encoding the 100-kDa subunit of trehalose-6-phosphate

- synthase/phosphatase complex in *Saccharomyces cerevisiae*, causes accumulation of trehalose-6-phosphate and loss of trehalose-6-phosphate phosphatase activity. *Eur. J. Biochem.* **212**, 315-323.
- De Virgilio, C., Burckert, N., Boller, T. and Wiemken, A. (1991). A method to study the rapid phosphorylation-related modulation of neutral trehalase activity by temperature shifts in yeast. *FEBS Lett.* **291**, 355-358.
- De Virgilio, C., Hottiger, T., Dominguez, J., Boller, T. and Wiemken, A. (1994). The role of trehalose synthesis for the acquisition of thermotolerance in yeast. I. Genetic evidence that trehalose is a thermoprotectant. *Eur. J. Biochem.* **219**, 179-186.
- De Virgilio, C., Piper, P., Boller, T. and Wiemken, A. (1991). Acquisition of thermotolerance in *Saccharomyces cerevisiae* without heat shock protein hsp104 and in the absence of protein synthesis. *FEBS Lett.* **288**, 86-90.
- De Virgilio, C., Simmen, U., Hottiger, T., Boller, T. and Wiemken, A. (1990). Heat shock induces enzymes of trehalose metabolism, trehalose accumulation, and thermotolerance in *Schizosaccharomyces pombe*, even in the presence of cycloheximide. *FEBS Lett.* **273**, 107-110.
- Diddens, H. A. and Lodder, J. (1942). *Die eukaryotischen Hefen, II Hälfte*. North Holland Publ. Co., Amsterdam.
- Dietz, T. J. (1994). Acclimation of the threshold induction temperatures for 70-kDa and 90-kDa heat shock proteins in the fish *Gillichthys mirabilis*. *J. Exp. Biol.* **188**, 333-338.
- Dietz, T. J. and Somero, G. N. (1992). The threshold induction temperature of the 90-kDa heat shock protein is subject to acclimatization in eurythermal goby fishes (genus *Gillichthys*). *Proc. Natl. Acad. Sci. USA* **89**, 3389-3393.
- di Menna, M. E. (1966a). Three new yeasts from Antarctic soils: *Candida nivalis*, *Candida gelida*, and *Candida frigida*. *Anton. van Leeuwenhoek* **32**, 25-28.
- di Menna, M. E. (1966b). Yeasts in Antarctic soils. *Anton. Van Leeuwenhoek* **32**, 29-38.
- Dowds, B. C. A. (1994). The oxidative stress response in *Bacillus subtilis*. *FEMS Lett.* **124**, 255-263.
- Durnez, P., Pernambuco, M. B., Oirs, E., Argüelles, J. C., Mergelsberg, H. and Thevelein, J. M. (1994). Activation of trehalase during growth induction by nitrogen sources in the yeast *Saccharomyces cerevisiae* depends on the free catalytic subunits of cAMP-dependent protein kinase, but not on functional Ras proteins. *Yeast* **10**, 1049-1064.

- Elliot, B. and Futcher, B. (1993). Stress resistance of yeast cells is largely independent of cell cycle phase. *Yeast* **9**, 33-42.
- Elliot, B., Haltiwanger, R. S. and Futcher, B. (1996). Synergy between trehalose and hsp 104 for thermotolerance in *Saccharomyces cerevisiae*. *Genetics* **144**, 923-933.
- Ellis, R. J. (1990). Molecular chaperones: The plant connection. *Science* **250**, 954-959.
- Ellis, R. J. (1992). Cytosolic chaperonin confirmed. *Nature* **358**, 191-192.
- Ellis, R. J. (1996). Stress proteins as molecular chaperones. In *Stress Proteins in Medicine*. pp. 1-26. Edited by van Eden, W. and Young, D. B. Marcel Dekker, Inc., New York.
- Ellis, R. J. and van der Vies, S. M. (1991). Molecular chaperones. *Ann. Rev. Biochem.* **60**, 321-347.
- Eraso, P. and Gancedo, C. (1987). Activation of yeast plasma membrane ATPase by acid pH during growth. *FEBS Lett.* **224**, 187-192.
- Farr, S. B. and Kogoma, T. (1991). Oxidative stress responses in *Escherichia coli* and *Salmonella typhimurium*. *Microbiol. Rev.* **55**, 561-585.
- Farrelly, F. W. and Finkelstein, D. B. (1984). Complete sequence of the heat shock-inducible HSP90 gene of *Saccharomyces cerevisiae*. *J. Biol. Chem.* **259**, 5745-5751.
- Feige, U. and Mollenhauer, J. (1992). Heat shock proteins. Introduction. *Experientia* **48**, 621-622.
- Feinberg, A. P. and Vogelstein, B. (1983). A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* **132**, 6-13.
- Fell, J. W. (1976). Yeasts in oceanic regions. In *Recent Advances in Aquatic Mycology*. pp. 93-124. Edited by Gareth Jones, E. B. Eek Sciences, London.
- Fell, J. W., Statzell, A. C., Hunter, I. L. and Phaff, H. J. (1969). *Leucosporidium* gen. N. the heterobasidiomycetes stage of several yeasts of the genus *Candida*. *Anton. van Leeuwenhoek* **35**, 433-462.
- Feofilova, E. P. (1994). Biochemical adaptation of mycelial fungi to temperature stress. *Microbiol.* **63**, 421-432.
- Feofilova, E. P., Tereshina, V. M. and Gornova, I. B. (1994). Changes in carbohydrate composition of fungi during adaptation to thermostress. *Microbiol.* **63**, 442-445.
- Ferbeyre, G., Villareal, A. and Morales, J. (1993). A rapid procedure for preparing high molecular weight DNA from yeast for Southern analysis. *BioTech.* **14**, 386.

- Fernandes, P. M. B., Panek, A. D. and Kurtenbach, E. (1997). Effect of hydrostatic pressure on a mutant of *Saccharomyces cerevisiae* deleted in the trehalose-6-phosphate synthase gene. *FEBS Lett.* **152**, 17-21.
- Finley, D., Ozkaynak, E. and Varshavsky, A. (1987). The yeast polyubiquitin gene is essential for resistance to high temperature, starvation, and other stresses. *Cell* **48**, 1035-1946.
- Flaherty, K. M., De Luca, F. C. and McKay, D. B. (1990). Three-dimensional structure of the ATPase fragment of a 70K heat-shock cognate protein. *Nature* **346**, 623-628.
- Flattery O'Brien, J., Collinson, L. P. and Dawes, I. W. (1993). *Saccharomyces cerevisiae* has an inducible response to menadione which differs from that to hydrogen peroxide. *J. Gen. Microbiol.* **139**, 501-507.
- François, J., Neves, M.-J. and Hers, H.-G. (1984). The mechanism by which glucose increases fructose 2,6-bisphosphate concentration in *Saccharomyces cerevisiae*. *Eur. J. Biochem.* **145**, 187-193.
- François, J., van Schaftingen, E. and Hers, H.-G. (1986). Effect of benzoate on the metabolism of fructose 2,6-bisphosphate in yeast. *Eur. J. Biochem.* **154**, 141-145.
- Frankenberg, D., Frankenberg-Schwager, M. and Harbich, R. (1993). Mechanisms of oxygen radiosensitization in irradiated yeast. I. DNA double-strand breakage. *Int. J. Radiat. Biol.* **64**, 511-521.
- Franzmann, P. D. (1995). Due south for useful microorganisms. *Today's Life Science* **June**, 27-30.
- Freeman, M. L., Spitz, D. R. and Meredith, M. J. (1990). Does heat shock enhance oxidative stress? Studies with ferrous and ferric iron. *Radiat. Res.* **124**, 288-293.
- Frydman, J. and Hartl, F.-U. (1994). Molecular chaperone functions of hsp70 and hsp60 in protein folding. In *The Biology of Heat Shock Proteins and Molecular Chaperones*. pp. 251-283. Edited by Morimoto, R. I., Tissières, A. and Georgopoulos, C., Cold Spring Harbor Laboratory Press, Cold Spring Harbor.
- Fuge, E. K., Braun, E. L. and Werner-Washburne, M. (1994). Protein synthesis in long-term stationary-phase cultures of *Saccharomyces cerevisiae*. *J. Bacteriol.* **176**, 5802-5813.
- Garbe, T. R. (1992). Heat shock proteins and infection: Interactions of pathogen and host. *Experientia* **48**, 635-639.

- Gaston, J. S. H. and Pearce, J. H. (1996). Immune response to heat shock proteins in reactive arthritis. In *Stress Proteins in Medicine*. pp. 103-117. Edited by van Eden, W. and Young, D. B. Marcel Dekker, Inc., New York.
- Georgopoulos, C., Liberek, K., Zylacz, M and Ang, D. (1994). Properties of the heat shock proteins of *Escherichia coli* and the auto-regulation of the heat shock response. In *The Biology of Heat Shock Proteins and Molecular Chaperones*. pp. 209-249. Edited by Morimoto, R. I., Tissières, A. and Georgopoulos, C. Cold Spring Harbor Laboratory Press, Cold Spring Harbor.
- Georgopoulos, C. and McFarland, H. (1993). Heat shock proteins in multiple sclerosis and other autoimmune diseases. *Immunol. Today* **14**, 373-375.
- Gething, M.-J. and Sambrook, J. (1992). Protein folding in the cell. *Nature* **355**, 33-45.
- Giménez-Jurado, G. and van Uden, N. (1989). *Leucosporidium fellii* sp. Nov., a basidiomycetous yeast that degrades L(+)-tartaric acid. *Anton. van Leeuwenhoek* **55**, 133-141.
- Goldstein, J., Pollitt, N. S. and Inouye, M. (1990). Major cold shock protein of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **87**, 283-287.
- Gottschal, J. C. and Prins, R. A. (1991). Thermophiles: A life at elevated temperatures. *TREE* **6**, 157-162.
- Goto, S., Sugiyama, J. and Iizuka, H. (1969). A taxonomic study of Antarctic yeasts. *Mycologia* **61**, 748-774.
- Gounalaki, N. and Thireos, G. (1994). *Yap1p*, a yeast transcriptional activator that mediates multidrug resistance, regulates the metabolic stress response. *EMBO J.* **13**, 4036-4041.
- Gounot, A-M. (1991). Bacterial life at low temperature: physiological aspects and biotechnological implications. *J. Appl. Bacteriol.* **71**, 386-397.
- Gralla, E. B. and Kosman, D. J. (1992). Molecular genetics of superoxide dismutases in yeasts and related fungi. *Adv. Genet.* **30**, 251-319.
- Gralla, E. B., Thiele, D. J., Silar, P. and Valentine, J. S. (1991). ACE1, a copper-dependent transcription factor, activates expression of the yeast copper, zinc superoxide dismutase gene. *Proc. Natl. Acad. Sci. USA* **88**, 8558-8562.
- Grant, C. M., Firoozan, M. and Tuite, M. F. (1989). Mistranslation induces the heat shock response in the yeast *Saccharomyces cerevisiae*. *Mol. Microbiol.* **3**, 215-219.

- Grant, C. M., MacIver, F. H. and Dawes, I. W. (1996). Glutathione is an essential metabolite required for resistance to oxidative stress in the yeast *Saccharomyces cerevisiae*. *Curr. Genet.* **29**, 511-515.
- Gross, C. and Watson, K. (1996). Heat shock protein synthesis and trehalose accumulation are not required for induced thermotolerance in derepressed *Saccharomyces cerevisiae*. *Biochem. Biophys. Res. Comm.* **220**, 766-772.
- Gupta, R. (1990). Sequence and structural homology between a mouse t-complex protein TCP-1 and the chaperonin family of bacteria (groEL, 60-65 kDa heat shock antigen) and eukaryotic proteins. *Biochem. Int.* **20**, 833-839.
- Hall, B. G. (1983). Yeast thermotolerance does not require protein synthesis. *J. Bacteriol.* **156**, 1363-1365.
- Halliwell, B., and Aruoma, O. (1993). *DNA and Free Radicals*. Ellis Horwood, London.
- Halliwell, B. and Gutteridge, J. M. C. (1986). Oxygen free radicals and iron in relation to biology and medicine: some problems and concepts. *Arch. Biochem. Biophys.* **246**, 501-514.
- Halliwell, B. and Gutteridge, J. M. C. (1989). *Free radicals in Biology and medicine*. Clarendon Press, Oxford.
- Hartl, F.-U. (1995). Mechanisms and pathways of chaperone-mediated protein folding. In *Cold Spring Harbor Symposia on Quantitative Biology*. Vol. LX. pp. 429-439. Cold Spring Harbor Laboratory Press, New York.
- Hartl, F.-U. (1996). Molecular chaperones in cellular protein folding. *Nature* **381**, 571-580.
- Hartman, D. J., Hoogenraad, N. J., Condron, R. and Hoj, P. B. (1992). Identification of a mammalian 10-kDa heat shock protein, a mitochondrial chaperonin 10 homologue essential for assisted folding of trimeric ornithine transcarbamoylase *in vitro*. *Proc. Natl. Acad. Sci. USA* **89**, 3394-3398.
- Hayes, S. A. and Dice, J. F. (1996). Roles of molecular chaperones in protein degradation. *J. Cell. Biol.* **132**, 255-258.
- Heinemeyer, W., Kleinschmidt, J. A., Saidowsky, J., Escher, C. and Wolf, D. H. (1991). Proteinase yscE, the yeast proteasome/multicatalytic-multifunctional proteinase: Mutants unravel its function in stress induced proteolysis and uncover its necessity for cell survival. *EMBO J.* **10**, 555-562.

- Hemmingsen, S. M., Woolford, C., van der Vies, S. M., Tilly, K., Dennis, D. T., Georgopoulos, C. P., Hendrix, R. W. and Ellis, R. J. (1988). Homologous plant and bacterial chaperone oligomeric protein assembly. *Nature* **333**, 330-334.
- Hendrick, J. P. and Hartl, F.-U. (1993). Molecular chaperone functions of heat shock proteins. *Ann. Rev. Biochem.* **62**, 349-384.
- Hendrix, R. W. (1979). Purification and properties of GroE, a host protein involved in bacteriophage assembly. *J. Mol. Biol.* **129**, 375-392.
- Herbert, R. A. (1992). A perspective on the biotechnological potential of extremophiles. *Trends Biotech.* **10**, 395-402.
- Hershko, A. (1991). The ubiquitin pathway for protein degradation. *Trends Biochem. Sci.* **16**, 265-267.
- Hightower, L. E., Sadis, S. E. and Takenaka, I. M. (1994). Interaction of vertebrate hsc 70 and hsp 70 with unfolded proteins and peptides. In *The Biology of Heat Shock Proteins and Molecular Chaperones*. pp. 179-207. Edited by Morimoto, R. I., Tissières, A. and Georgopoulos, C. Cold Spring Harbor Laboratory Press, Cold Spring Harbor.
- Hilt, W., Enenkel, C., Gruhler, A., Singer, T. and Wolf, D. H. (1992). The PRE4 gene codes for a subunit of the yeast proteasome necessary for peptidyl glutamyl-peptide hydrolyzing activity. Mutations link the proteasome to stress and ubiquitin dependent proteolysis. *J. Biol. Chem.* **268**, 3479-3486.
- Hilt, W. and Wolf, D. H. (1992). Stress-induced proteolysis in yeast. *Mol. Microbiol.* **6**, 2437-2442.
- Hohn, T., Hohn, B., Engel, A., Wortz, M. and Smith, P. R. (1979). Isolation and characterisation of the host protein GroE involved in bacteriophage lambda assembly. *J. Mol. Biol.* **129**, 359-373.
- Hottiger, T., Boller, T. and Wiemken, A. (1987). Rapid changes of heat and dessication tolerance with changes of trehalose content in *Saccharomyces cerevisiae* cells subjected to temperature shifts. *FEBS Lett.* **220**, 113-115.
- Hottiger, T., Boller, T. and Wiemken, A. (1989). Correlation of trehalose content and heat resistance in yeast mutants altered in the RAS / adenylate cyclase pathway: is trehalose a thermoprotectant? *FEBS Lett.* **255**, 431-434.
- Hottiger, T., De Virgilio, C., Bell, W., Boller, T. and Wiemken, A. (1992). The 70-kilodalton heat-shock proteins of the SSA subfamily negatively modulate heat-shock-

- induced accumulation of trehalose and promote recovery from heat stress in the yeast *Saccharomyces cerevisiae*. *Eur. J. Biochem.* **210**, 125-132.
- Hottiger, T., De Virgilio, C., Hall, M., Boller, T. and Wiemken, A. (1994). The role of trehalose synthesis for the acquisition of thermotolerance in yeast. II. Physiological concentrations of trehalose increase the thermal stability of proteins *in vitro*. *Eur. J. Biochem.* **219**, 187-193.
- Howarth, C. J. and Ougham, H. J. (1993). Gene expression under temperature stress. *New Phytologist* **125**, 1-26.
- Hurley, R., de Louvois, J. and Mulhall, A. (1987). Yeasts as human and animal pathogens. In *The Yeasts*. 2nd ed. Vol. 2. pp. 207-281. Edited by Rose, A. H. and Harrison, J. S. Academic Press, London.
- Iida, H. and Yahara, I. (1985). Yeast heat shock protein of  $M_r$  48,000 is an isoprotein of enolase. *Nature* **315**, 688-690.
- Imlay, J. A. and Lim, S. (1988). DNA damage and oxygen radical toxicity. *Science* **240**, 1302-1309.
- Ingolia, T. D. and Craig, E. A. (1982). Four small *Drosophila* heat shock proteins are related to each other and to mammalian  $\alpha$ -crystallin. *Proc. Natl. Acad. Sci. USA* **79**, 2360-2364.
- Ingram, L. O. (1986). Microbial tolerance to alcohols: role of the cell membrane. *Trends Biotechnol.* **4**, 40-44.
- Inniss, W. E. (1975). Interaction of temperature and psychrophilic microorganisms. *Ann. Rev. Microbiol.* **29**, 445-465.
- Iwahashi, H., Obuchi, K., Fujii, S. and Komatsu, Y. (1997). Effect of temperature on the role of HSP104 and trehalose in barotolerance of *Saccharomyces cerevisiae*. *FEBS Lett.* **416**, 1-5.
- Iwahashi, H., Wu, Y. and Tanguay, R.M. (1995). Detection and expression of the 70 kDa heat shock protein ssb1p at different temperatures in *Saccharomyces cerevisiae*. *Biochem. Biophys. Res. Comm.* **213**, 484-489.
- Jain, S. (1988). Evidence for lipid peroxidation during *in vivo* aging of human erythrocytes. *Biochim. Biophys. Acta* **937**, 201-210.
- Jamieson, D. J. (1992). *Saccharomyces cerevisiae* has distinct adaptive responses to hydrogen peroxide and menadione. *J. Bacteriol.* **174**, 6678-6681.

- Jamieson, D. J., Stephen, D. W. S. and Terrière, E. D. (1996). Analysis of the adaptive oxidative stress response of *Candida albicans*. *FEMS Lett.* **138**, 83-88.
- Jamieson, D. J., Rivers, S. L. and Stephen, D. W. S. (1994). Analysis of *Saccharomyces cerevisiae* proteins induced by peroxide and superoxide stress. *Microbiol.* **140**, 3277-3283.
- Jentsch, S. (1992). The ubiquitin-conjugation system. *Ann. Rev. Genet.* **26**, 179-207.
- Jerez, C. A. (1988). The heat shock response in meso- and thermoacidophilic chemolithotrophic bacteria. *FEMS Lett.* **56**, 289-294.
- Jones, P. G., Cashel, M., Glaser, G. and Neidhardt, F. C. (1992). Function of a relaxed-like state following temperature downshifts in *Escherichia coli*. *J. Bacteriol.* **174**, 3903-3914.
- Jones, P. G. and Inouye, M. (1994). The cold-shock response – a hot topic. *Mol. Microbiol.* **11**, 811-818.
- Jones, P. G., Van Bogelen, R. A. and Neidhardt, F. C. (1987). Induction of proteins in response to low temperature in *Escherichia coli*. *J. Bacteriol.* **169**, 2092-2095.
- Julseth, C. R. and Inniss, W. E. (1990a). Heat shock protein induction and the acquisition of thermotolerance in the psychrotrophic yeast *Trichosporon pullulans*. *Curr. Microbiol.* **20**, 391-396.
- Julseth, C. R. and Inniss, W. E. (1990b). Induction of protein synthesis in response to cold-shock in the psychrotrophic yeast *Trichosporon pullulans*. *Can. J. Microbiol.* **36**, 519-524.
- Jungmann, J., Reins, H. A., Lee, J., Romeo, A., Hassett, R., Kosman, D. and Jentsch, S. (1993). Mac-1, a nuclear regulatory protein related to Cu-dependent transcription factors is involved in Cu/Fe utilization and stress resistance in yeast. *EMBO J.* **12**, 5051-5056.
- Kang, P. J., Osterman, J., Shilling, J., Neuupert, W., Craig, E. A. and Pfanner, N. (1990). Requirements for HSP70 in the mitochondrial matrix for translocation and folding of precursor proteins. *Nature* **348**, 137-143.
- Kapoor, M. and Lewis, J. (1986). Activation of the protein synthesis pattern in *Neurospora crassa* cells by hyperthermal and oxidative stress. *Can. J. Microbiol.* **33**, 162-168.
- Kapoor, M. and Lewis, J. (1987). Heat shock induces peroxidase activity in *Neurospora crassa* and confers tolerance towards oxidative stress. *Biochem. Biophys. Res. Comm.* **147**, 904-910.

- Kappus, H. (1985). Lipid peroxidation: mechanisms, analysis – enzymology and biological relevance. In *Oxidative Stress*. pp. 273-310. Edited by Sies, H. Academic Press, London.
- Kates, M. (1964). Bacterial lipids. *Adv. Lipid Res.* **2**, 17-19.
- Kates, M. and Baxter, R. M. (1962). Lipid composition of mesophilic and psychrophilic yeasts (*Candida* sp.) as influenced by environmental temperature. *Can. J. Biochem. Physiol.* **40**, 1213-1227.
- Kato, M., Miura, Y., Kettoku, M., Shindo, K., Iwamatsu, A. and Kobayashi, K. (1996). Purification and characterization of new trehalose-producing enzymes isolated from the hyperthermophilic archae, *Sulfolobus solfataricus* KM1. *Biosci. Biotech. Biochem.* **60**, 546-550.
- Kaufmann, S. H. E. (1991). *Heat Shock Proteins and Immune Response*. Springer-Verlag, Berlin.
- Kaufmann, S. H. E. (1992). The cellular immune response to heat shock proteins. *Experientia* **48**, 640-649.
- Kelly, R. M. and Brown, S. H. (1993). Enzymes from high-temperature microorganisms. *Curr. Opin. Biotech.* **4**, 188-192.
- Kobayashi, N and McEntee, K. (1990). Evidence for a heat shock transcription factor-independent mechanism for heat shock induction of transcription in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* **87**, 6550-6554.
- Kobayashi, N and McEntee, K. (1993). Identification of *cis* and *trans* components of a novel heat shock stress regulatory pathway in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **13**, 248-256.
- Kondo, K. and Inouye, M. (1991). *TIP1*, a cold shock-inducible gene of *Saccharomyces cerevisiae*. *J. Biol. Chem.* **266**, 17537-17544.
- Kondo, K. and Inouye, M. (1992). Yeast NSR1 protein that has structural similarity to mammalian nucleolin is involved in pre-rRNA processing. *J. Biol. Chem.* **267**, 16252-16258.
- Kondo, K., Kowalski, L. R. Z. and Inouye, M. (1992). Cold shock induction of yeast NSR1 protein and its role in pre-rRNA processing. *J. Biol. Chem.* **267**, 16259-16265.
- Kopp, M., Müller, H. and Holzer, H. (1993). Molecular analysis of the neutral trehalase gene from *Saccharomyces cerevisiae*. *J. Biol. Chem.* **268**, 4766-4774.

- Kowalski, L. R. Z., Kondo, K. and Inouye, M. (1995). Cold-shock induction of a family of TIP1-related proteins associated with the membrane in *Saccharomyces cerevisiae*. *Mol. Microbiol.* **15**, 341-353.
- Kramhoft, B. and Zeuthen, E. (1971). Synchronization of cell division in the fission yeast, *Schizosaccharomyces pombe*, using heat shocks. *C.R. Trav. Lab. Carlsberg* **38**, 351-368.
- Kristjansson, J. B. (1989). Thermophilic organisms as sources of thermostable enzymes. *Trends Biotech.* **7**, 349-353.
- Kuge, S. and Jones, N. (1994). YAP1 dependent activation of TRX2 is essential for the response of *Saccharomyces cerevisiae* to oxidative stress by hydroperoxides. *EMBO J.* **13**, 655-664.
- Kullik, I. and Storz, G. (1994). Transcriptional regulators of the oxidative stress response in prokaryotes and eukaryotes. *Redox Report* **1**, 23-29.
- Kurtz, S. and Lindquist, S. (1984). The changing pattern of gene expression in sporulating yeast. *Proc. Natl. Acad. Sci. USA* **81**, 7323-7327.
- Kurtz, S., Rossi, J., Petko, L. and Lindquist, S. (1986). An ancient developmental induction: Heat shock proteins induced in sporulation and oogenesis. *Science* **231**, 1154-1157.
- Labbe-Rosine, R. and Labbe, P. (1990). Tetrapyrrole and heme biosynthesis in the yeast *Saccharomyces cerevisiae*. In *Biosynthesis of Heme and Chlorophylls*. pp. 235-285. Edited by Dailey, H. A. McGraw-Hill Book Co., New York.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of the bacteriophage T4. *Nature* **227**, 680-685.
- Laemmli, U. K. and Favre, M. (1973). Maturation of the head of bacteriophage T4. I. DNA packaging events. *J. Mol. Biol.* **80**, 575-599.
- Langer, T., Buchner, J. and Bukau, B. (1996). Chaperone function on Crete: a meeting report. *Cell Stress Chap.* **1**, 5-12.
- Laskey, R. A., Honda, B. M., Mills, A. D. and Fitch, J. T. (1978). Nucleosomes are assembled by an acidic protein which binds histones and transfers them to DNA. *Nature* **275**, 416-420.
- La Teana, A., Brandi, A., Falconi, M., Spurio, R., Pon, C. L. and Gualerzi, C. O. (1991). Identification of a cold shock transcriptional enhancer of the *Escherichia coli* gene encoding nucleoid protein N-HS. *Proc. Natl. Acad. Sci. USA* **88**, 10907-10911.

- Lawrence, R. A. and Burk, R. F. (1976). Glutathione peroxidase activity in selenium-deficient rat liver. *Biochem. Biophys. Res. Comm.* **71**, 952-958.
- Lee, J., Dawes, I. W. and Roe, J.-H. (1995). Adaptive response of *Schizosaccharomyces pombe* to hydrogen peroxide and menadione. *Microbiol.* **141**, 3127-3132.
- Lee, J.-S., Hah, Y-C. and Roe, J.-H. (1993). The induction of oxidative enzymes in *Streptomyces coelicolor* upon hydrogen peroxide treatment. *J. Gen. Microbiol.* **139**, 1013-1018.
- Levine, R. L., Williams, J. A., Stadtman, E. R., and Shacter, E. (1994). Carbonyl assays for determination of oxidatively modified proteins. *Meth. Enzymol.* **233**, 346-357.
- Lewis, J. G., Learmonth, R. P., Attfield, P. V. and Watson, K. (1997). Stress co-tolerance and trehalose content in baking strains of *Saccharomyces cerevisiae*. *J. Ind. Microbiol. Biotechnol.* **18**, 30-36.
- Lewis, J. G., Learmonth, R. P. and Watson, K. (1993). The role of growth phase and ethanol in freeze-thaw stress resistance of *Saccharomyces cerevisiae*. *Appl. Env. Microbiol.* **59**, 1065-1071.
- Li, G. C. and Hahn, G. M. (1978). Ethanol-induced tolerance to heat and adriamycin. *Nature* **274**, 699-701.
- Li, G. C., Shiu, E. C. and Hahn, G. M. (1980). Similarities in cellular inactivation by hyperthermia or by ethanol. *Radiat. Res.* **82**, 257-268.
- Lillie, S. and Pringle, J. R. (1980). Reserve carbohydrate metabolism in *Saccharomyces cerevisiae*: Responses to nutrient limitation. *J. Bacteriol.* **143**, 1384-1394.
- Lindquist, S. (1980). Translational efficiency of heat induced messages in *Drosophila melanogaster* cells. *J. Mol. Biol.* **137**, 151-158.
- Lindquist, S. (1991). Genetic analysis of heat shock protein functions in yeast. In *Heat Shock*. pp. 123-131. Edited by Maresca, B. and Lindquist, S. Springer-Verlag, Berlin.
- Lindquist, S. and Craig, E. A. (1988). The heat shock proteins. *Ann. Rev. Genet.* **55**, 1151-1191.
- Lindquist, S. and Kim, G. (1996). Heat-shock protein 104 expression is sufficient for thermotolerance in yeast. *Proc. Natl. Acad. Sci. USA* **93**, 5301-5306.
- Lindquist, S., DiDomenico, B., Bugaisky, G., Kurtz, S., Sonoda, S. (1982). Regulation of the heat-shock response in *Drosophila* and yeast. In *Heat shock: From bacteria to man*. pp. 167-175. Edited by Schlesinger, M. J., Ashburner, M. and Tissieres, A. Cold Spring Harbor Laboratory Press, New York.

- Lindquist, S. and Kim, G. (1996). Heat-shock protein 104 expression is sufficient for thermotolerance in yeast. *Proc. Natl. Acad. Sci. USA* **93**, 5301-5306.
- Lindquist, S., Patino, M. M., Chernoff, Y. O., Kowal, A. S., Singer, M. A., Liebman, S. W., Lee, K.-H. and Blake, T. (1995). The role of Hsp104 in stress tolerance and [PSI<sup>+</sup>] propagation in *Saccharomyces cerevisiae*. In *Cold Spring Harbor Symposia on Quantitative Biology*. Vol. LX. pp. 451-460. Cold Spring Harbor Laboratory Press, Cold Spring Harobor.
- Lodder, J. and Kreger-van Rij, N. J. W. (1952). *The Yeasts, a taxonomic study*. pp. 1-33. North Holland Publ. Co., Amsterdam.
- Londesborough, J. and Varimo, K. (1984). Characterization of two trehalases in bakers' yeast. *Biochem. J. A.*, 511-518.
- Lubben, T. H., Gattenby, A. A., Donaldson, G. K., Lorimer, G. H. and Viitanen, P. V. (1990). Identification of a groES-like chaperonin in mitochondria that facilitates protein folding. *Proc. Natl. Acad. Sci. USA* **87**, 7683-7687.
- Luke, M., Suttin, A., Arndt, K. (1991). Characterization of *SIS1*, a *Saccharomyces cerevisiae* homologue of bacterial *dnaJ* protein. *J. Cell. Biol.* **114**, 623-638.
- MacNamara, E. E. (1973). Macro- and microclimates of the Antarctic coastal oasis, Molodezhnaya. *Biuletyn Peryglacjalny* **23**, 201-236.
- Mager, W. H. and de Kruijff, A. A. J. (1995). Stress-induced transcriptional activation. *Microbiol. Rev.* **59**, 506-531.
- Mager, W. H. and Moradas-Ferreira, P. (1993). Stress response of yeast. *Biochem. J.* **290**, 1-13.
- Marchler, G., Schüller, C., Adam, G. and Ruis, H. (1993). A *Saccharomyces cerevisiae* UAS element controlled by protein kinase A activates transcription in response to a variety of stress conditions. *EMBO J.* **12**, 1997-2003.
- Maresca, B. and Carratù, L. (1990). Heat shock and adaptation during temperature-activated dimorphism in the fungus *Histoplasma capsulatum*. In *Stress proteins - Induction and function*. pp. 45-56. Edited by Schlesinger, M. J., Santoro, M.G. and Garaci, E. Springer-Verlag, Berlin Heidelberg.
- Maresca, B. and Kobayashi, G. S. (1996). Heat shock proteins in fungal infections. In *Stress proteins in medicine*. pp. 287-299. Edited by van Eden, W. and Young, D. B. Marcel Dekker, Inc., New York.

- Margesin, R. and Schinner, F. (1994). Properties of cold-adapted microorganisms and their potential role in biotechnology. *J. Biotechnol.* **33**, 1-14.
- Martin, J., Horwich, A. L. and Hartl, F.-U. (1992). Prevention of protein denaturation under heat stress by the chaperonin Hsp60. *Science* **258**, 995-998.
- Mathews, C. K. and van Holde, K. E. (1990). Biochemistry. Benjamin/Cummings, California.
- M<sup>c</sup>Alister, L. and Holland, M. J. (1985a). Isolation and characterization of yeast strains carrying mutations in the glyceraldehyde-3-phosphate dehydrogenase genes. *J. Biol. Chem.* **260**, 15013-15018.
- M<sup>c</sup>Alister, L. and Holland, M. J. (1985b). Differential expression of the three yeast glyceraldehyde-3-phosphate dehydrogenase genes. *J. Biol. Chem.* **260**, 15019-15027.
- M<sup>c</sup>Alister, L., Strausberg, S., Kulaga, A. and Finkelstein, D. B. (1979). Altered patterns of protein synthesis induced by heat shock of yeast. *Curr. Genet.* **1**, 63-74.
- M<sup>c</sup>Callum, K. L., Heikkila, J. J. and Inniss, W. E. (1986). Temperature dependent pattern of heat shock protein synthesis in psychrophilic and psychrotrophic microorganisms. *Can. J. Microbiol.* **32**, 516-521.
- M<sup>c</sup>Callum, K. L. and Inniss, W. E. (1990). Thermotolerance, cell filamentation, and induced protein synthesis in psychrophilic and psychrotrophic bacteria. *Arch. Microbiol.* **153**, 585-590.
- M<sup>c</sup>Clanahan, T. and M<sup>c</sup>Entee, K. (1986). DNA damage and heat shock dually regulate gene expression in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **6**, 90-96.
- M<sup>c</sup>Cord, J. M. and Day, E. D. (1978). Superoxide-dependent production of hydroxyl radical catalysed by iron-EDTA complex. *FEBS Lett.* **86**, 139-142.
- M<sup>c</sup>Cord, J. M. and Fridovich, J. (1969). Superoxide dismutase. An enzyme function for erythrocuprein (Hemocuprein). *J. Biol. Chem.* **244**, 6049-6056.
- M<sup>c</sup>Kay, D. B. (1991). Structure of the 70-kilodalton heat-shock-related proteins. *Springer. Semin. Immunopathol.* **13**, 1-9.
- M<sup>c</sup>Kay, D. B., Wilbanks, S. M., Flaherty, K. M., Ha, J.-H., O'Brien, M. C. and Shirvane, L. L. (1994). Stress-70 proteins and their interactions with nucleotides. In *The Biology of Heat Shock Proteins and Molecular Chaperones*. pp. 153-177. Edited by Morimoto, R. I., Tissières, A. and Georgopoulos, C., Cold Spring Harbor Laboratory Press, Cold Spring Harbor.

- McMullen, T. W. and Hallberg, R. L. (1988). A highly evolutionary conserved mitochondrial protein is structurally related to the protein encoded by the *Escherichia coli* *groEL* gene. *Mol. Cell. Biol.* **8**, 371-380.
- Meister, A. (1988). Glutathione metabolism and its selective modification. *J. Biol. Chem.* **263**, 17205-17208.
- Meister, A. and Anderson, M. E. (1983). Glutathione. *Ann. Rev. Biochem.* **52**, 711-760.
- Mendonça-Hagler, L. C. and Phaff, H. J. (1975). Deoxyribonucleic acid base composition and deoxyribonucleic acid/deoxyribonucleic acid hybrid formation in psychrophobic and related yeasts. *Int. J. Syst. Bacteriol.* **25**, 222-229.
- Meyer, E. D., Sinclair, N. A. and Nagy, B. (1975). Comparison of the survival and metabolic activity of psychrophilic and mesophilic yeasts subjected to freeze-thaw stress. *Appl. Microbiol.* **29**, 739-744.
- Miller, M. J., Xuong, N. H. and Geiduschek, E. P. (1982). Quantitative analysis of the heat shock response of *Saccharomyces cerevisiae*. *J. Bacteriol.* **151**, 311-327.
- Miralles, V. J. and Serrano, R. (1995). A genomic locus in *Saccharomyces cerevisiae* with four genes up-regulated by osmotic stress. *Mol. Microbiol.* **17**, 653-662.
- Mishra, P. (1993). Tolerance of fungi to ethanol. In *Stress Tolerance in Fungi*. pp. 189-208. Jenning, D. H. Marcel Dekker, New York.
- Mishra, P. and Prasad, R. (1987). Alterations in fatty acyl composition can selectively affect amino acid transport in *Saccharomyces cerevisiae*. *Biochem. Int.* **15**, 499-508.
- Mishra, P. and Prasad, R. (1988). Role of phospholipid head groups in ethanol tolerance of *Saccharomyces cerevisiae*. *J. Gen. Microbiol.* **134**, 3205-3211.
- Mitchel, R. E. J. and Morrison, D. P. (1982). Heat-shock induction of ionizing radiation resistance in *Saccharomyces cerevisiae*, and correlation with stationary growth phase. *Radiat. Res.* **90**, 284-291.
- Mittenbühler, K. and Holzer, H. (1988). Purification and characterization of acid trehalase from the yeast *SUC2* mutant. *J. Biol. Chem.* **263**, 8537-8543.
- Mollenhauer, J. and Schulmeister, A. (1992). The humoral immune response to heat shock proteins. *Experientia* **48**, 644-649.
- Mongkolsuk, S., Vattanaviboon, P. and Praituan, W. (1997). Induced adaptive and cross-protection responses against oxidative stress killing a bacterial phytopathogen *Xanthomonas oryzae* pv. *oryzae*. *FEMS Lett.* **146**, 217-222.

- Moody, C. S. and Hassan, H. M. (1982). Mutagenicity of oxygen free radicals. *Proc. Natl. Acad. Sci. USA* **79**, 2855-2859.
- Moore, R. T. (1980). Taxonomic proposals for the classification of marine yeasts and other yeast like fungi including the smuts. *Botanica Marina* **23**, 361-373.
- Moradas-Ferreira, P., Costa, V., Piper, P. and Mager, W. (1996). The molecular defenses against reactive oxygen species in yeast. *Mol. Microbiol.* **19**, 651-658.
- Morimoto, R. I., Tissières, A. and Georgopoulos, C. (1990). The stress response, function of the proteins, and perspectives. In *Stress Proteins in Biology and Medicine*. pp. 1-36. Edited by Morimoto, R. I., Tissières, A. and Georgopoulos, C., Cold Spring Harbor Laboratory Press, New York.
- Morita, R. Y. (1975). Psychrophilic bacteria. *Microbiol. Rev.* **39**, 144-167.
- Moye-Rowley, W. S., Harshman, K. D. and Parker, C. S. (1989). Yeast *YAP1* encodes a novel form of the jun family of transcriptional activator proteins. *Genes Dev.* **3**, 283-292.
- Mukai, H., Kuno, T., Tanaka, H., Hirata, D., Miyakawa, T. and Tanaka, C. (1993). Isolation and characterization of SSE1 and SSE2, new members of the yeast HSP70 multigene family. *Gene* **132**, 57-66.
- Musgrove, J. E., Johnson, R. A. and Ellis, R. J. (1987). Dissociation of the ribulose bisphosphate carboxylase large-subunit binding protein into dissimilar subunits. *Eur. J. Biochem.* **163**, 529-534.
- Mutoh, N., Nakagawa, C. W. and Hayashi, Y. (1995). Adaptive response of *Schizosaccharomyces pombe* to hydrogen peroxide. *FEMS Lett.* **132**, 67-72.
- Nash, C. H., Grant, D. W. and Sinclair, N. A. (1968). Thermolability of protein synthesis in a cell-free system from the obligately psychrophilic yeast *Candida gelida*. *Can. J. Microbiol.* **15**, 339-343.
- Nelson, R. J., Heschl, M. and Craig, E. A. (1992a). Isolation and characterization of extragenic suppressors of mutations in the SSA hsp70 genes of *Saccharomyces cerevisiae*. *Genetics* **131**, 277-285.
- Nelson, R. J. and Ziegelhoffer, T., Nicolet, C., Werner-Washburne, M. and Craig, E. A. (1992b). The translation machinery and seventy kilodalton heat shock protein cooperated in protein synthesis. *Cell* **71**, 97-105.
- Neupert, W., Hartl, F.-U., Craig, E. A. and Pfanner, N. (1990). How do polypeptides cross the mitochondrial membranes? *Cell* **63**, 447-450.

- Neves, M.-J. and François, J. (1992). On the mechanism by which a heat shock induces trehalose accumulation in *Saccharomyces cerevisiae*. *Biochem. J.* **288**, 559-864.
- Neves, M. J., Jorge, J. A., François, J. M. and Terenzi, H. F. (1991). Effects of heat shock on the level of trehalose and glycogen, and on the induction of thermotolerance in *Neurospora crassa*. *FEBS Lett.* **283**, 19-22.
- Nicolet, C. M. and Craig, E. A. (1989). Isolation and characterization of *STI1*, a stress-inducible gene from *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **9**, 3638-3646.
- Normington, K., Kohno, K., Kozutsumi, V., Gething, M.-J. and Sambrook, J. (1989). *S. cerevisiae* encodes an essential protein homologous in sequence and function to mammalian BiP. *Cell* **57**, 1223-1236.
- Novák, E. K. and Zsolt, J. (1961). A new system proposed for yeasts. *Acta Bot. Acad. Sci. Hung.* **7**, 93-145.
- Nwaka, S. and Holzer, H. (1998). Molecular biology of trehalose and trehalases in the yeast *Saccharomyces cerevisiae*. *Prog. Nuc. Acid. Res. Mol. Biol.* **53**, 197-237 (in press).
- Nwaka, S., Kopp, M., Burgert, M., Deuchler, I., Kienle, I. and Holzer, H. (1994). Is thermotolerance of yeast dependent on trehalose accumulation? *FEBS Lett.* **344**, 225-228.
- Nwaka, S., Mechler, B., van Ahsen, O. and Holzer, T. (1996). The heat shock factor and mitochondrial Hsp70 are necessary for survival of heat shock in *Saccharomyces cerevisiae*. *FEBS Lett.* **399**, 259-263.
- Ostermann, J., Horwich, A. L., Neupert, W. and Hartl, F.-U. (1989). Protein folding in mitochondria requires complex formation with hsp60 and ATP hydrolysis. *Nature* **341**, 125-130.
- Ozkaynak, E., Finley, D., Solomon, M. J. and Varshavsky, A. (1987). The yeast ubiquitin genes: A family of natural gene fusions. *EMBO J.* **6**, 1429-1439.
- Pahl, H. L. and Baeuerle, P. A. (1994). Oxygen and the control of gene expression. *BioEssays* **16**, 497-502.
- Panaretou, B. and Piper, P. W. (1990). Plasma-membrane ATPase action affects several stress tolerances of *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe* as well as the extent and duration of the heat shock response. *J. Gen. Microbiol.* **136**, 1763-1770.

- Panaretou, B. and Piper, P. W. (1992). The plasma membrane of yeast acquires a novel heat-shock protein (hsp30) and displays a decline in proton-pumping ATPase levels in response to both heat shock and the entry into stationary phase. *Eur. J. Biochem.* **206**, 635-640.
- Panek, A. D. and Mattoon, J. R. (1977). Regulation of energy metabolism in *Saccharomyces cerevisiae*. Relationships between catabolite repression, trehalose synthesis, and mitochondrial development. *Arch. Biochem. Biophys.* **183**, 306-316.
- Parag, H. A., Raboy, B. and Kulka, R. G. (1987). Effect of heat shock on protein degradation in mammalian cells: Involvement of the ubiquitin system. *EMBO J.* **6**, 55-61.
- Parry, J. M., Davies, P. J. and Evans, W. E. (1976). The effects of "cell age" upon the lethal effects of physical and chemical mutagens in the yeast, *Saccharomyces cerevisiae*. *Mol. Gen. Genet.* **146**, 27-35.
- Parsell, D. A., Kowal, A. S., Singer, M. A., Lindquist, S. (1994). Protein disaggregation mediated by heat-shock protein Hsp104. *Nature* **372**, 475-477.
- Parsell, D. A. and Lindquist, S. (1994). Heat shock proteins and stress tolerance. In *The Biology of Heat Shock Proteins and Molecular Chaperones*. pp. 457-494. Edited by Morimoto, R. I., Tissières, A. and Georgopoulos, C. Cold Spring Harbor Laboratory Press, Cold Spring Harbor.
- Parsell, D. A., Sanchez, Y., Stitzel, J. D. and Lindquist, S. (1991). Hsp104 is a highly conserved protein with two essential nucleotide-binding sites. *Nature* **353**, 270-273.
- Patchett, R. A., Watson, N., Fernandez, P. S., Kroll, R. G. (1996). The effect of temperature and growth rate on the susceptibility of *Listeria monocytogenes* to environmental stress conditions. *Lett. Appl. Microbiol.* **22**, 121-124.
- Pelham, H. R. B. (1984). Hsp70 accelerates the recovery of nucleolar morphology after heat shock. *EMBO J.* **3**, 3095-3100.
- Pelham, H. R. B. (1986). Speculation on the functions of the major heat shock and glucose regulated stress proteins. *Cell* **46**, 959-961.
- Pelham, H. R. B. (1989). Heat shock and the sorting of luminal ER proteins. *EMBO J.* **8**, 3171-3176.
- Perisic, O., Xiao, H. and Lis, J. T. (1989). Stable binding of Drosophila heat shock factor to head-to-head and tail-to-tail repeats of a conserved 5bp recognition unit. *Cell* **59**, 797-806.

- Petko, L. and Lindquist, S. (1986). Hsp26 is not required for growth at high temperature, nor for thermotolerance, spore development, or germination. *Cell* **45**, 885-894.
- Picard, D., Khursheed, B., Garabedian, M. J., Fortin, M. G., Lindquist, S. and Yamamoto, K. R. (1990). Reduced levels of hsp 90 compromise steroid receptor action *in vivo*. *Nature* **348**, 166-168.
- Piper, P. (1990). Interdependence of several heat shock gene activations, cyclic AMP decline and changes at the plasma membrane of *Saccharomyces cerevisiae*. *Anton. van Leeuwenhoek* **58**, 195-201.
- Piper, P., Curran, M., Davies, A., Lockheart, A., Reid, G. (1986). Transcription of the phosphoglycerate kinase gene of *Saccharomyces cerevisiae* increases when fermentative cultures are stressed by heat shock. *Eur. J. Biochem.* **161**, 525-531.
- Piper, P. W. (1993). Molecular events associated with acquisition of heat tolerance by the yeast *Saccharomyces cerevisiae*. *FEMS Microbiol. Rev.* **11**, 339-356.
- Piper, P. W. (1995). The heat shock and ethanol stress responses of yeast exhibit extensive similarity and functional overlap. *FEMS Lett.* **134**, 121-127.
- Piper, P. W., Talreja, K., Panaretou, B., McRadas-Ferreira, P., Byrne, K., Praekelt, U. M., Meacock, P., Récnacq, M. and Boucherie, H. (1994). Induction of major heat-shock proteins of *Saccharomyces cerevisiae*, including plasma membrane Hsp30, by ethanol levels above a critical threshold. *Microbiol.* **140**, 3031-3038.
- Plesofsky-Vig, N. and Brambl, R. (1993). Heat shock proteins in fungi. In *Stress Tolerance of Fungi*. pp. 45-68. Edited by Jennings, D. H. Marcel Dekker, New York.
- Plesofsky-Vig, N., Vig, J. and Brambl, R. (1992). Phylogeny of the  $\alpha$ -crystallin-related heat shock proteins. *J. Mol. Evol.* **35**, 537-545.
- Plesset, J., Ludwig, J. R., Cox, B. S. and McLaughlin, C. S. (1987). Effect of cell cycle position on thermotolerance in *Saccharomyces cerevisiae*. *J. Bacteriol.* **169**, 779-784.
- Plesset, J., Palm, C. and McLaughlin, C. S. (1982). Induction of heat shock proteins and thermotolerance by ethanol in *Saccharomyces cerevisiae*. *Biochem. Biophys. Res. Comm.* **108**, 1340-1345.
- Polaina, J. and Conde, J. (1982). Genes involved in the control of nuclear fusion during the sexual cycle of *Saccharomyces cerevisiae*. *Mol. Gen. Genet.* **186**, 253-258.
- Polla, B. S. (1991). Heat shock proteins in host-parasite interactions. *Parasitol. Today and Immunol. Today Combined March issues*, A38-A41.

- Potier, P., Drevet, P., Gounot, A.-M., and Hipkiss, A. R. (1987). Protein turnover in a psychrotrophic bacterium: proteolytic activity in extracts of cells grown at different temperatures. *FEMS Lett.* **44**, 267-271.
- Praekelt, U. M. and Meacock, P. A. (1990). HSP12, a new small heat shock gene of *Saccharomyces cerevisiae*: Analysis of structure, regulation and function. *Mol. Gen. Genet.* **223**, 97-106.
- Pratt, W. B., Scherrer, L. C., Hutchinson, K. A. and Kalman, F. C. (1992). A model of glucocorticoid receptor unfolding and stabilization by a heat shock protein complex. *J. Steroid Biochem. Mol. Biol.* **41**, 223-229
- Ramírez, C. and González, A. (1984). Four new species of *Apotrichum* isolated from the decayed wood in the evergreen rainy Valdivian forest of southern Chile. *Mycopathologia* **88**, 71-78.
- Reading, D. S., Hallberg, R. L. and Myers, A. M. (1989). Characterization of the yeast HSP60 gene coding for a mitochondrial assembly factor. *Nature* **337**, 655-659.
- Rechsteiner, M., Rogers, S. and Rote, K. (1987). Protein structure and intracellular stability. *Trends Biochem. Sci.* **12**, 390-394.
- Régnacq, M. and Boucherie, H. (1993). Isolation and sequence of HSP30, a yeast heat-shock gene coding for a hydrophobic membrane protein. *Curr. Genet.* **23**, 435-442.
- Reed, K. (1990). Basic blotting – A quick fix. *Today's Life Science May*, 52-60.
- Richter-Ruoff, B., Heinemeyer, W. and Wolf, D. H. (1992). The proteasome/multicatalytic-multifunctional protease: In vivo function in the ubiquitin-dependent N-end rule pathway of protein degradation in eukaryotes. *FEBS Lett.* **302**, 192-196.
- Ritossa, F. M. (1962). A new puffing pattern induced by a temperature shock and DNP in *Drosophila*. *Experientia* **18**, 571-573.
- Rocha, E. R., Selby, T., Coleman, J. P. and Smith, C. J. (1996). Oxidative stress response in an anaerobe, *Bacteroides fragilis*: a role for catalase in protection against hydrogen peroxide. *J. Bacteriol.* **178**, 6895-6903.
- Rollo, F., Sassaroli, S. and Ubaldi, M. (1995). Molecular phylogeny of the fungi of the Iceman's grass clothing. *Curr. Genet.* **28**, 289-297.
- Rose, A. H. (1987). Responses to the chemical environment. In *The Yeast: Yeast and the Environment*. Vol. 2. pp. 5-40. Edited by Rose, A. H. and Harrison, J. S. Academic Press, London.

- Rose, M. D., Misra, L. M. and Vogel, J. P. (1989). *KAR2*, a karogamy gene, is the yeast homolog of the mammalian BiP/GRP78 gene. *Cell* **57**, 1211-1221.
- Rossi, J. M. and Lindquist, S. (1989). The intracellular location of yeast heat-shock protein 26 varies with metabolism. *J. Cell. Biol.* **108**, 425-439.
- Rothblatt, J. A., Deshaies, R. J., Sanders, S. L., Daum, G. and Schekman, R. (1989). Multiple genes are required for proper insertion of secretory proteins into the endoplasmic reticulum in yeast. *J. Cell. Biol.* **109**, 2641-2652.
- Rothman, J. E. (1989). Polypeptide chain binding proteins: Catalysts of protein folding and related processes in cells. *Cell* **59**, 591-601.
- Ruis, H. and Hamilton, B. (1992). Regulation of yeast catalase genes. In *Molecular Biology of Free Radical Scavenging Systems*. pp. 153-172. Edited by Scandalios, J. G. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Ruis, H. and Schüller, C. (1995). Stress signalling in yeast. *Bioessays* **17**, 959-965.
- Russell, N. J. (1990). Cold adaptation of microorganisms. *Phil. Trans. Roy. Soc. London B* **326**, 595-611.
- Russell, N. J. (1992). Physiology and molecular biology of psychrophilic micro-organisms. In *Molecular Biology and Biotechnology of Extremophiles*. pp. 203-224. Edited by Herbert, R. A. and Sharp, R. J. Blackie and Son, London.
- Russo, P., Lakkinen, N., Sareneva, H., Paakkola, J. and Makarow, M. (1992a). A heat shock gene from *Saccharomyces cerevisiae* encoding a secretory glycoprotein. *Proc. Natl. Acad. Sci. USA* **89**, 3671-3675.
- Russo, P., Lakkinen, N., Sareneva, H., Paakkola, J. and Makarow, M. (1992b). A heat shock gene from *Saccharomyces cerevisiae* encoding a secretory glycoprotein. Published errata. *Proc. Natl. Acad. Sci. USA* **89**, 8857.
- Russo, P., Simonen, M., Uimari, A., Reesalu, T. and Makarow, M. (1993). Dual regulation by heat and nutrient stress of the yeast *HSP150* gene encoding a secretory glycoprotein. *Mol. Gen. Genet.* **239**, 273-280.
- Sadler, I., Chiang, A., Kurihara, T., Rothblatt, J., Way, J. and Silver, P. (1989). A yeast gene important for protein assembly into the endoplasmic reticulum and the nucleus has homology to DnaJ, an *Escherichia coli* heat shock protein. *J. Cell. Biol.* **109**, 2665-2675.
- Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989). *Molecular Cloning – A Laboratory Manual*. 2nd Ed. Edited by Ford, N. Cold Spring Harbor Laboratory Press, New York.

- Sanchez, Y. and Lindquist, S. (1990). Hsp 104 required for induced thermotolerance. *Science* **248**, 1112-1115.
- Sanchez, Y., Parsell, D. A., Taulein, J., Vogel, J. L., Craig, E. A. and Lindquist, S.L. (1993). Genetic evidence for a functional relationship between Hsp 104 and Hsp 70. *J. Bacteriol.* **175**, 6484-6491.
- Sanchez, Y., Taulein, J., Borkovich, K. A. and Lindquist, S. (1992). Hsp 104 is required for tolerance to many forms of stress. *EMBO J.* **11**, 2357-2364.
- Sanders, S., Whitfield, K., Vogel, J., Rose, M. and Schekman, R. (1992). Sec61p and BiP directly facilitate polypeptide translocation into the ER. *Cell* **69**, 353-366.
- Schenberg-Francino, A. and Moustacchi, E. (1972). Lethal and mutagenic effects of elevated temperature on haploid yeast. I. Variations in sensitivity during the cell cycle. *Mol. Gen. Genet.* **115**, 243-257.
- Schirmer, E. D., Glover, J. R., Singer, M. A. and Lindquist, S. (1996). Hsp100/Clp proteins: A common mechanism explains diverse functions. *Trends Biochem. Sci.* **21**, 289-296.
- Schirmer, E. C., Lindquist, S. and Vierling, E. (1994). An Arabidopsis heat shock protein complements a thermotolerance defect in yeast. *The Plant Cell* **6**, 1899-1909.
- Schlesinger, M. J. (1990). Heat shock proteins. *J. Biol. Chem.* **265**, 12111-12114.
- Schnell, N., Krems, B. and Entain, K-D. (1992). The PAR1 (YAP1/SNQ3) gene of *Saccharomyces cerevisiae*, a c-jun homologue, is involved in oxygen metabolism. *Curr. Genet.* **21**, 269-273.
- Schoel, B. and Kaufmann, S. H. E. (1996). The unique role of heat shock proteins in infections. In *Stress proteins in medicine*. pp. 27-51. Edited by van Eden, W. and Young, D. B. Marcel Dekker, Inc., New York.
- Schüller, C., Brewster, J. L., Alexander, M. R., Gustin, M. C. and Ruis, H. (1994). The HOG pathway controls osmotic regulation of transcription via the stress response element (STRE) of the *Saccharomyces cerevisiae CTT1* gene. *EMBO J.* **13**, 4382-4389.
- Schumacher, R. J., Hansen, W. J., Freeman, B. C., Alnerri, E., Litwack, G. and Toft, D. O. (1996). Cooperative action of Hsp70, Hsp90 and DnaJ proteins in protein renaturaton. *Biochem.* **35**, 14889-14898.
- Sen, S., Thangavelu, M., Pinder, S. and Roser, B. (1993). Extraordinary stability of enzymes dried in trehalose - simplified molecular biology. *Bio-Technology* **10**, 1007-1012.

- Sherman, F. (1956). The heat inactivation and production of cytochrome deficiency in yeast. *Exp. Cell. Res.* **11**, 659-660.
- Sherman, F. (1959). The effects of elevated temperatures of yeast. II. Induction of respiratory-deficient mutants. *J. Cell. Comp. Physiol.* **54**, 37-52.
- Shinnik, T. M., Vodkin, M. H. and Williams, J. C. (1988). The *Mycobacterium tuberculosis* 65-kilodalton antigen is a heat shock protein which corresponds to common antigen and to the *Escherichia coli* GroEL protein. *Infect. Immun.* **56**, 446-451.
- Sies, H. (1986). Biochemistry of oxidative stress. *Angew. Chem. Int. Ed. Engl.* **25**, 1058-1071.
- Simon, M., Adam, G., Rapatz, W., Spevak, W. and Ruis, H. (1991). The *Saccharomyces cerevisiae* *ADRI* gene is a positive regulator of transcription of genes encoding peroxisomal proteins. *Mol. Cell. Biol.* **11**, 699-704.
- Sinclair, N. A. and Stokes, J. L. (1963). Role of oxygen in the high cell yields of psychrophiles and mesophiles at low temperatures. *J. Bacteriol.* **85**, 164-167.
- Sinclair, N. A. and Stokes, J. L. (1965). Obligately psychophilic yeasts from the polar regions. *Can. J. Microbiol.* **11**, 259-269.
- Slade, H. J. K., Jones, D. T. and Woods, D. R. (1984). Effect of oxygen and peroxide on *Bacteroides fragilis* cell and phage survival after treatment with DNA damaging agents. *FEMS Lett.* **24**, 159-163.
- Slater, T. F. (1979). Mechanisms of protection against damage produced in biological systems by oxygen-derived radicals. In *Oxygen Free Radicals and Tissue Damage*. pp. 142-176. Excerpta Medica, Amsterdam.
- Smith, B. J. and Yaffe, M. P. (1991). Uncoupling thermotolerance from the induction of heat shock proteins. *Proc. Natl. Acad. Sci. USA* **88**, 11091-11094.
- Somero, G. N. (1995). Proteins and temperature. *Annu. Rev. Physiol.* **57**, 43-68.
- Sorger, P. K., Lewis, M. J. and Pelham, H. R. B. (1987). Heat shock factor is regulated differently in yeast and HeLa cells. *Nature* **329**, 81-84.
- Sorger, P. K. and Nelson, H. C. M. (1989). Trimerization of a yeast transcriptional activator via a coiled-coil motif. *Cell* **59**, 807-813.
- Sorger, P. K. and Pelham, H. R. B. (1988). Yeast heat shock factor is an essential DNA-binding protein that exhibits temperature-dependent phosphorylation. *Cell* **54**, 855-864.
- Souza, N. O. and Panek, A. D. (1968). Location of trehalase and trehalose in yeast cells. *Arch. Biochem. Biophys.* **125**, 22-28.

- Spiro, R. G. (1966). Analysis of sugars found in glycoproteins. *Meth. Enzymol.* **8**, 3-26.
- Steels, E. S., Learmonth, R. and Watson, K. (1994). Stress tolerance and membrane lipid unsaturation in *Saccharomyces cerevisiae* grown aerobically or anaerobically. *Microbiol.* **140**, 569-576.
- Stephen, D. W. S. and Jamieson, D. J. (1996). Glutathione is an important antioxidant molecule in the yeast, *Saccharomyces cerevisiae*. *FEMS Lett.* **141**, 207-212.
- Stephen, D. W. S., Rivers, S. L. and Jamieson, D. J. (1995). The role of the *YAP1* and *YAP2* genes in the regulation of the adaptive oxidative stress responses of *Saccharomyces cerevisiae*. *Mol. Microbiol.* **16**, 415-423.
- Stevenson, M. A. and Calderwood, S. K. (1990). Members of the 70-kilodalton heat-shock protein family contain a highly conserved calmodulin-binding domain. *Mol. Cell. Biol.* **10**, 1234-1238.
- Stewart, P. R. (1975). Analytical methods for yeast. *Meth. Cell Biol.* **7**, 125-127.
- Stokes, J. L. (1971). Influence of temperature on growth and metabolism of yeasts. In *The Yeasts*. 2nd ed. Vol. 2. pp. 119-134. Edited by Rose, A. H. and Harrison, J. S. Academic Press, London.
- Stone, D. E. and Craig, E. A. (1990). Self-regulation of 70-kilodalton heat shock proteins in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **10**, 1622-1632.
- Stryer, L. (1988). Biochemistry. 3rd Ed. W H. Freeman and Company, New York.
- Sur, I., Lobo, Z. and Maitra, P. K. (1994). Analysis of PFK3 – a gene involved in particulate phosphofructokinase synthesis reveals additional functions of *TPS2* in *Saccharomyces cerevisiae*. *Yeast* **10**, 199-209.
- Susek, R. E. and Lindquist, S. L. (1989). *hsp26* of *Saccharomyces cerevisiae* is related to the superfamily of small heat shock proteins but is without a demonstrable function. *Mol. Cell. Biol.* **9**, 5265-5271.
- Susek, R. E. and Lindquist, S. (1990). Transcriptional derepression of the *Saccharomyces cerevisiae* HSP26 gene during heat shock. *Mol. Cell. Biol.* **10**, 6362-6373.
- Swan, T. M. and Watson, K. (1997). Membrane fatty acid composition and membrane fluidity as parameters of stress tolerance in yeast. *Can. J. Microbiol.* **43**, 70-77.
- Switzer, R. C., Merril, C. R. and Shifrin, S. (1979). A highly sensitive silver stain for detecting proteins and peptides in polyacrylamide gels. *Anal. Biochem.* **98**, 231-237.
- Tanaka, K., Matsumoto, K. and Toh-e, A. (1988). Dual regulation of the expression of the polyubiquitin gene by cyclic AMP and heat shock in yeast. *EMBO J.* **7**, 495-502.

- Tansey, M. R. and Brock, T. D. (1972). The upper temperature limit for eucaryotic organisms. *Proc. Natl. Acad. Sci. USA* **69**, 2426-2428.
- Tansey, M. R. and Brock, T. D. (1978). Microbial life at high temperatures: Ecological aspects. In *Microbial life in extreme environments*. pp. 159-216. Edited by Kushner, D. J. Academic Press, London.
- Teichert, U., Mechler, H., Müller, H. and Wolf, D. H. (1989). Lysosomal (vacuolar) proteinases of yeast are essential catalysts for protein degradation, differentiation, and cell survival. *J. Biol. Chem.* **264**, 16037-16045.
- Thevelein, J. M. (1984). Regulation of trehalose metabolism in fungi. *Microbiol. Rev.* **48**, 42-59.
- Thevelein, J. M. (1991). Fermentable sugars and intracellular acidification as specific activators of the RAS-adenylate cyclase signalling pathway in yeast: The relationship to nutrient-induced cell cycle control. *Mol. Microbiol.* **5**, 1049-1074.
- Thevelein, J. M. and Hohmann, S. (1995). Trehalose synthase: guard to the gate of glycolysis in yeast? *Trends Biochem. Sci.* **20**, 3-10.
- Thiele, D. J. (1988). ACE1 regulates expression of the *Saccharomyces cerevisiae* metallothionein gene. *Mol. Cell. Biol.* **8**, 2745-2752.
- Tissières, A., Mitchell, H. K. and Tracy, V. M. (1974). Protein synthesis in salivary glands of *Drosophila melanogaster*: Relation to chromosome puffs. *J. Mol. Biol.* **84**, 389-398.
- Travassos, L. R. R. G. and Cury, A. (1971) Thermophilic enteric yeasts. *Ann. Rev. Microbiol.* **25**, 49-74.
- Trent, J. D. (1996). A review of acquired thermotolerance, heat-shock proteins and molecular chaperones in archaea. *FEMS Microbiol. Rev.* **18**, 249-258.
- Trent, J. D., Gabrielsen, M., Jensen, B., Neuhard, J. and Olsen, J. (1994). Acquired thermotolerance and heat shock proteins in thermophiles from the three phylogenetic domains. *J. Bacteriol.* **176**, 6148-6152.
- Trent, J. D., Osipiuk, J. and Pinkau, T. (1990). Acquired thermotolerance and heat shock in the extremely thermophilic archaebacterium *Sulfolobus* strain B12. *J. Bacteriol.* **172**, 1478-1484.
- Tuite, M. F., Bentley, N. J., Bossier, P. and Fitch, I. T. (1990). The structure and function of small heat shock proteins: Analysis of the *Saccharomyces cerevisiae* Hsp26 protein. *Anton. van Leeuwenhoek* **58**, 147-154.

- Ursic, D. and Culbertson, M. R. (1991). The yeast homolog to mouse Tcf-1 affects microtubule-mediated processes. *Mol. Cell. Biol.* **11**, 2629-2640.
- van Aelst, L., Hohmann, S., Bulaya, B., de Koning, W., Sierkstra, L., Neves, M. J., Luyten, K., Alijo, R., Ramos, J., Coccetti, P., Martegani, E., de Magelhães-Rocha, N. M., Brandão, R. L., Van Dijck, P., Vanhalewyn, M., Durnez, P., Jans, A. W. H. and Thevelein, J. M. (1993). Molecular cloning of a gene involved in glucose sensing in the yeast *Saccharomyces cerevisiae*. *Mol. Microbiol.* **8**, 927-943.
- van Bogelen, R. A., Kelley, P. M. and Neidhardt, F. C. (1987). Differential induction of heat shock SOS and oxidative stress regulons and accumulation of nucleotides in *Escherichia coli*. *J. Bacteriol.* **169**, 26-32.
- van Bogelen, R. A. and Neidhardt, F. C. (1990). Ribosomes as sensors of heat and cold shock in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **87**, 5589-5593.
- Vandercammen, A., François, J. and Hers, H.-G. (1989). Characterisation of trehalose-6-phosphate synthase and trehalose-6-phosphate phosphatase of *Saccharomyces cerevisiae*. *Eur. J. Biochem.* **182**, 613-620.
- van der Vaart, J. M., Caro, L. H., Chapman, J. W., Klis, F. M. and Verrips, C. T. (1995). Identification of three mannoproteins in the cell wall of *Saccharomyces cerevisiae*. *J. Bacteriol.* **177**, 3104-3110.
- van der Walt, J. P. (1957). Three new sporogenous yeasts from soil. *Anton. van Leeuwenhoek* **23**, 23-29.
- van der Walt, J. P. and Yarrow, D. (1984). The genus *Axiozyma* gen. nov. (Saccharomycetaceae). *South African J. Botany* **3**, 340-342.
- van Dijck, P., Colavizza, D., Smet, P. and Thevelein, J. M. (1995). Differential importance of trehalose in stress resistance in fermenting and nonfermenting *Saccharomyces cerevisiae* cells. *Appl. Environ. Microbiol.* **61**, 109-115.
- van Eden, W., Thole, J. E. R., Van der Zee, R., Van Embden, J. D. A., Hensen, E. J. and Cohen, I. R. (1988). Cloning of the mycobacterial epitope recognized by T lymphocytes in adjuvant arthritis. *Nature* **331**, 171-173.
- van Eden, W. and Young, D. B. (1996). *Stress Proteins in Medicine*. Marcel Dekker, New York.
- van Laere, A. (1989). Trehalose, reserve and/or stress metabolite? *FEMS Microbiol. Rev.* **63**, 201-210.

- van Uden, N. (1952). Zur Kenntnis von *Torulopsis pintolopesii* sp. nov. *Arch. Mikrobiol.* **17**, 199-208.
- van Uden, N. (1984a). Temperature profiles of yeasts. *Adv. Microbial Physiol.* **25**, 195-251.
- van Uden, N. (1984b). Effects of ethanol on the temperature relations of viability and growth in yeast. *Crit. Rev. Biotechnol.* **1**, 263-272.
- van Uden, N. and do Carmo-Sousa, L. (1957). Yeasts from the bovine caecum. *J. Gen Microbiol.* **16**, 385-395.
- van Uden, N. and Vidal-Leira, M. (1970). The genus *Torulopsis* Berlese. In *The Yeasts, a taxonomic study*. pp. 1235-1308. Edited by Lodder, J. North Holland Publ. Co., Amsterdam.
- Varela, J. C., van Beekvelt, C., Planta, R. J. and Mager, W. H. (1992). Osmostress-induced changes in yeast gene expression. *Mol. Microbiol.* **6**, 2183-2190.
- Varela, J. C. S., Praekelt, U. M., Meacock, P. A., Planta, R. J. and Mager, W. H. (1995). The *Saccharomyces cerevisiae* *HSP12* gene is activated by the High-Osmolarity glycerol pathway and negatively regulated by protein kinase A. *Mol. Cell. Biol.* **15**, 6232-6245.
- Vayda, M. E. and Yuan, M.-L. (1994). The heat shock response of an antarctic alga is evident at 5°C. *Plant Mol. Biol.* **24**, 229-233.
- Verduyn, C., van Dijken, J. P. and Scheffers, W. A. (1984). Colorimetric alcohol assays with alcohol oxidase. *J. Microbiol. Meth.* **2**, 15-25.
- Verma, R., Iida, H. and Pardee, A. (1988a). Identification of a novel stress-inducible glycoprotein in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **263**, 8569-8575.
- Verma, R., Iida, H. and Pardee, A. (1988b). Modulation of expression of the stress-inducible p118 of *Saccharomyces cerevisiae*. *J. Biol. Chem.* **263**, 8576-8582.
- Vishniac, H. S. (1993). The microbiology of Antarctic soils. In *Antarctic Microbiology* pp. 297-341. Edited by Friedman, E. I. Wiley-Liss, New York.
- Vishniac, H. S. (1996). Biodiversity of yeasts and filamentous microfungi in terrestrial Antarctic ecosystems. *Biodiversity and Conservation* **5**, 1365-1378.
- Vishniac, H. S. and Kurtzman, C. P. (1992). *Cryptococcus antarcticus* sp. nov. and *Cryptococcus albidosimilis* sp. nov., Basidioblastomycetes from Antarctic Soils. *Int. J. Sys. Bacteriol.* **42**, 547-553.

- Vogel, J. L., Parsell, D. A. and Lindquist, S. (1995). Heat-shock proteins HSP104 and HSP70 reactivate mRNA splicing after heat inactivation. *Curr. Biol.* **5**, 306-317.
- Vogel, J. P., Misra, L. M. and Rose, M.D. (1990). Loss of BiP/GRP78 function blocks translocation of secretory proteins in yeast. *J. Cell. Biol.* **110**, 885-895.
- Vogelstein, B. and Gillespie, D. (1979). Preparative and analytical purificationof DNA from agarose. *Proc. Natl. Acad. Sci. USA* **76**, 615-620.
- Vuoria, O., Kalkkinen, N. and Londesborough, J. (1993). Cloning of two related genes encoding the 56 kDa and 123 kDa subunits of trehalose synthase from the yeast *Saccharomyces cerevisiae*. *Eur. J. Biochem.* **216**, 849-861.
- Walton, E. F. and Pringle, J. R. (1980). Effect of growth temperature upon heat sensitivity in *Saccharomyces cerevisiae*. *Arch. Microbiol.* **124**, 285-287.
- Watson, K. (1980). Homeoviscous adaptation in psychophilic, mesophilic and thermophilic yeasts. In *Membrane Fluidity: Biophysical Techniques and Cellular Regulation*. pp. 349-363. Edited by Kates, M. and Kuksis, A. Humana Press, Clifton, New Jersey.
- Watson, K. (1984). Membrane lipid adaptation in yeasts. In *Membrane Fluidity: Biomembranes*. pp. 517-542. Edited by Kates, M. and Manson, L. A. Plenum Press, New York.
- Watson, K. (1987). Temperature relations. In *The Yeasts*. 2nd ed. Vol. 2. pp. 41-71. Edited by Rose, A. H. and Harrison, J. S. Academic Press, London.
- Watson, K. (1990). Microbial stress proteins. *Adv. Microbial. Physiol.* **31**, 183-223.
- Watson, K.. Arthur, H. and Blakey, M. (1980). Biochemical correlations among the thermophilic enteric yeasts *Torulopsis pintoipesii*, *Saccharomyces telluris*, and *Candida slooffii*. *J. Bacteriol.* **143**, 693-702.
- Watson, K., Arthur, H. and Morton, H. (1978). Thermal adaptation in yeast: Obligate psychrophiles are obligate aerobes and obligate thermophiles are facultative anaerobes. *J. Bacteriol.* **136**, 815-817.
- Watson, K., Arthur, H. and Shipton, W. A. (1976). *Leucosporidium* yeasts: Obligate psychrophiles which alter membrane-lipid and cytochrome composition with temperature. *J. Gen. Microbiol.* **97**, 11-18.
- Watson, K. and Cavicchioli, R. (1983). Acquisition of ethanol tolerance in yeast cells by a heat shock. *Biotechnol. Lett.* **5**, 683-687.

- Watson, K., Dunlop, G. and Cavicchioli, R. (1984). Mitochondrial and cytoplasmic protein syntheses are not required for heat shock acquisition of ethanol and thermotolerance in yeast. *FEBS Lett.* **172**, 299-302.
- Weitzel, G., Pilatus, U. and Rensing, L. (1987). The cytoplasmic pH, ATP content and total protein synthesis rate during heat-shock protein inducing treatments in yeast. *Exp. Cell Res.* **170**, 64-79.
- Welch, W. J. and Suhar, J. P. (1985). Morphological study of the mammalian stress response: Characterization of changes in cytoplasmic organelles, cytoskeleton, and nucleoli, and appearance of intracellular actin filaments in rat fibroblasts after heat shock treatment. *J. Cell. Biol.* **101**, 1198-1211.
- Wemmie, J. A., Szczypka, M. S., Thiele, D. J. and Moye-Rowley, W. S. (1994). Cadmium tolerance mediated by the yeast AP-1 protein requires the presence of an ATP-binding cassette transporter-encoding gene, YCF1. *J. Biol. Chem.* **269**, 32592-32597.
- Wiech, H., Buchner, M., Zimmermann, H., Zimmermann, R. and Jakob, U. (1993). Hsc 70, immunoglobulin heavy chain binding protein, and hsp 90 differ in their ability to stimulate transport of precursor proteins into mammalian microsomes. *J. Biol. Chem.* **268**, 7417-7421.
- Wiemken, A. (1990). Trehalose in yeast, stress protectant rather than reserve carbohydrate. *Anton. van Leeuwenhoek* **58**, 209-217.
- Wieser, R., Adam, G., Wagner, A., Schuller, C., Marchler, G., Ruis, H., Krawiec, Z. and Bilinski, T. (1991). Heat shock factor-independent heat control of transcription of the *CTT1* gene encoding the cytosolic catalase T of *Saccharomyces cerevisiae*. *J. Biol. Chem.* **266**, 12406-12411.
- Willinsky, G., Bang, H., Fisher, G. and Mahariel, M. (1992). Characterization of *cspB*, a *Bacillus subtilis* inducible cold shock gene affecting cell viability at low temperatures. *J. Bacteriol.* **174**, 6326-6335.
- Winkler, K., Kienle, I., Burgert, M., Wagner, J. C. and Holzer, H. (1991). Metabolic regulation of the trehalose content of vegetative yeast. *FEBS Lett.* **291**, 269-272.
- Wistow, G. (1990). Cold shock and DNA binding. *Nature* **344**, 823-824.
- Wolfe, K. H. and Lohan, A. J. E. (1994). Sequence around the centromere of *Saccharomyces cerevisiae* chromosome II: similarity of *CEN2* to *CEN4*. *Yeast* **10**, S41-S46.

- Wolff, S. P. and Dean, R. T. (1986). Fragmentation of proteins by free radicals and its effect on their susceptibility to enzyme hydrolysis. *Biochem. J.* **234**, 399-403.
- Wolffe, A. P., Tafuri, S., Ranjan, M. and Fainilari, M. (1992). The Y-box factors: a family of nucleic acid binding proteins conserved from *Escherichia coli* to man. *New Biologist* **4**, 290-298.
- Woodford, D. V., Parish, J. H. and Moradas-Ferreira, P. (1995). Hydrogen peroxide induces DNA damage in *Saccharomyces cerevisiae*. *Yeast* **11**, S149.
- Woods, D. R. and Jones, D. T. (1986). Physiological responses of *Bacteroides* and *Clostridium* strains to environmental stress factors. In *Advances in Microbial Physiology*. Edited by Rose, A. H. and Tempest, D. W. **28**, 1-64. Academic Press, London.
- Wu, A.-L. and Moye-Rowley, W. S. (1994). *GSH1*, which encodes  $\gamma$ -glutamylcysteine synthetase, is a target gene for yAP-1 transcriptional regulation. *Mol. Cell. Biol.* **14**, 5832-5839.
- Wu, A. L., Wemmie, J. A., Edgington, N. P., Goebl, M., Guevara, J. L. and Moye-Rowley, W. S. (1992). Yeast bZip proteins mediate pleiotropic drug and metal resistance. *J. Biol. Chem.* **268**, 18850-18858.
- Wynn-Williams, D. D. (1980). Seasonal fluctuations in microbial activity in Antarctic moss peat. *Biol. J. Linnean Soc.* **14**, 11-28.
- Yamada, Y. and Komagata, K. (1987). *Mrakia* gen. nov., a heterobasidomycetous yeast genus for the Q<sub>8</sub>-equipped, self-sporulating organisms which produce a unicellular metabasidium, formerly classified in the genus *Leucosporidium*. *J. Gen. Appl. Microbiol.* **33**, 455-457.
- Yang, X.-D. and Feige, U. (1992). Heat shock proteins in autoimmune disease. From causative antigen to specific therapy? *Experientia* **48**, 650-656.
- Yarrow, D. and Meyer, S. A. (1978). Proposal for amendment of the diagnosis of the genus *Candida* Berkout no. cons. *Int. J. Syst. Bacteriol.* **28**, 611-615.
- Ycas, M. (1956). A hereditary cytochrome deficiency appearing in yeast grown at elevated temperature. *Exp. Cell Res.* **10**, 746-748.
- Yonehara, M., Minami, Y., Kawata, Y., Nagai, J. and Yahara, I. (1996). Heat-induced chaperone activity of Hsp90. *J. Biol. Chem.* **271**, 2641-2645.
- Yost, H. J., Peterson, R. B. and Lindquist, S. L. (1990). RNA metabolism: Strategies for regulation in the heat shock response. *Trends Genet.* **6**, 223-227.

- Young, D. B., Lathigra, R., Hendrix, R., Sweetser, D. and Young, R. A. (1988). Stress proteins are immune targets in leprosy and tuberculosis. *Proc. Natl. Acad. Sci. USA* **85**, 4267-4270.
- Young, R. A. and Elliot, T. J. (1989). Stress proteins, infection and immune surveillance. *Cell* **59**, 5-8.
- Zeuthen, M. L. and Howard, D. H. (1989). Thermotolerance and the heat-shock response in *Candida albicans*. *J. Gen. Microbiol.* **135**, 2509-2518.
- Zitomer, R. S. and Lowry, C. V. (1992). Regulation of gene expression by oxygen in *Saccharomyces cerevisiae*. *Microbiol. Rev.* **56**, 1-11.