

Selective breeding as a mitigation tool for methane emissions from dairy cattle



Y. de Haas^{a,*}, R.F. Veerkamp^a, G. de Jong^b, M.N. Aldridge^a

^a Animal Breeding and Genomics, Wageningen University & Research, 6700 AH Wageningen, the Netherlands

^b CRV, 6800 AL Arnhem, the Netherlands

ARTICLE INFO

Article history:

Received 29 October 2020
Revised 21 January 2021
Accepted 25 January 2021
Available online 8 July 2021

Keywords:

Breeding goal
Genetic improvement
Greenhouse gases
Large-scale recording
Sustainable agriculture

ABSTRACT

The global livestock sector, particularly ruminants, contributes substantially to the total anthropogenic greenhouse gases. Management and dietary solutions to reduce enteric methane (**CH₄**) emissions are extensively researched. Animal breeding that exploits natural variation in **CH₄** emissions is an additional mitigation solution that is cost-effective, permanent, and cumulative. We quantified the effect of including **CH₄** production in the Dutch breeding goal using selection index theory. The current Dutch national index contains 15 traits, related to milk yield, longevity, health, fertility, conformation and feed efficiency. From the literature, we obtained a heritability of 0.21 for enteric **CH₄** production, and genetic correlations of 0.4 with milk lactose, protein, fat and DM intake. Correlations between enteric **CH₄** production and other traits in the breeding goal were set to zero. When including **CH₄** production in the current breeding goal with a zero economic value, **CH₄** production increases each year by 1.5 g/d as a correlated response. When extrapolating this, the average daily **CH₄** production of 392 g/d in 2018 will increase to 442 g/d in 2050 (+13%). However, expressing the **CH₄** production as **CH₄** intensity in the same period shows a reduction of 13%. By putting economic weight on **CH₄** production in the breeding goal, selective breeding can reduce the **CH₄** intensity even by 24% in 2050. This shows that breeding is a valuable contribution to the whole set of mitigation strategies that could be applied in order to achieve the goals for 2050 set by the EU. If the decision is made to implement animal breeding strategies to reduce enteric **CH₄** production, and to achieve the expected breeding impact, there needs to be a sufficient reliability of prediction. The only way to achieve that is to have enough animals phenotyped and genotyped. The power calculations offer insights into the difficulties that will be faced in trying to record enough data. Recording **CH₄** data on 100 farms (with on average 150 cows each) for at least 2 years is required to achieve the desired reliability of 0.40 for the genomic prediction.

© 2021 The Authors. Published by Elsevier B.V. on behalf of The Animal Consortium. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Implications

The global livestock sector, particularly ruminants, contributes substantially to the total anthropogenic greenhouse gases. Management and dietary solutions to reduce enteric methane emissions are extensively researched. Animal breeding that exploits natural variation in methane emissions is an additional mitigation solution that is cost-effective, permanent, and cumulative. By adding methane production to the Dutch breeding goal, selective breeding can reduce the methane intensity by 24% in 2050. This shows that breeding is a valuable contribution to the whole set of mitigation strategies that could be applied in order to achieve the goals for 2050 set by the EU.

Introduction

Livestock production is a significant contributor to greenhouse gas emissions globally at 5.6–7.5 Gigatonne of carbon dioxide equivalent (**GtCO_{2e}**) (Herrero et al., 2016). Methane (**CH₄**) produced via enteric fermentation by ruminants is the largest contributor to agricultural greenhouse gases at 1.6–2.7 **GtCO_{2e}** (Herrero et al., 2016). In the Netherlands, dairy cattle are the largest contributor to enteric **CH₄** with 49% of the total (Van der Maas et al., 2009).

Genetic selection of low **CH₄** emitting cows is a potential method to reduce the contribution of the dairy sector (Knapp et al., 2014). Unlike management and diet strategies which have been extensively researched (Beauchemin et al., 2008), genetics has the benefit that the effects of selection are additive and permanent. Selection experiments in other ruminants, including beef cattle (Bird-Gardiner et al., 2015), and sheep (Pinares-Patiño et al.,

* Corresponding author.

E-mail address: yvette.dehaas@wur.nl (Y. de Haas).

2013; Jonker et al., 2017), have shown a decrease in CH₄ production when selecting for low CH₄ lines.

Breeding programmes use selection index theory to ensure maximum gains and that trait changes are in the desired direction. The current selection indexes for the Dutch national breeding goal include 15 traits from the following categories; milk production, milk components, feed efficiency, health, reproduction, longevity, calving, and conformation (CRV, 2018). It is possible to add CH₄ as another category of traits to such selection indexes. Before a CH₄ trait can be added to a selection index, it needs to be: clearly defined, recordable, affordable, have phenotypic variation, be heritable, and the genetic correlations between other traits in the index need to be known (Hazel, 1943).

Methane intensity is a promising CH₄ trait to be included in a selection index (Lassen and Løvendahl, 2016; Kandel et al., 2018). The definition of CH₄ intensity is the amount of CH₄ produced per unit of milk produced (g CH₄/kg milk). It is more advantageous to have fewer cows that produce more milk per g of CH₄ (possible with selection for reduced CH₄ intensity), than to have more cows that produce less CH₄ but less milk, which could occur if selecting on CH₄ production alone (g CH₄/day). It has already been shown that, when assuming a fixed total milk production, selection for CH₄ intensity or increased productivity will increase an individual's total CH₄ but the overall CH₄ of the farming system is reduced (Van der Maas et al., 2009; Kandel et al., 2018).

Methane from cow's breath, recorded with non-dispersive infrared units or 'sniffers', have been investigated in small studies as an accurate method of recording CH₄ intensity (Lassen et al., 2012; Lassen and Løvendahl, 2016; Difford et al., 2018; Van Engelen et al., 2018). The sniffer method is more affordable, can be scaled up, and easily incorporated with current milking technologies, compared to other methods of recording CH₄ (Lassen and Løvendahl, 2016). Methane concentration recorded with sniffers has a phenotypic SD of between 65 and 137 ppm (Difford et al., 2018; González-Recio et al., 2020; López-Paredes et al., 2020; Olijhoek et al., 2020), and heritability ranges between 0.10 and 0.26 (Difford et al., 2020; López-Paredes et al., 2020; Olijhoek et al., 2020). Some genetic correlations have been published for CH₄ production and intensity, and traits already included in the Dutch national breeding goal selection index (Lassen and Løvendahl, 2016; Breider et al., 2019; Difford et al., 2020).

The effect of including CH₄ intensity in the selection index of the Dutch national breeding goal is not known. Recent literature suggests that if current selection indexes are used with no selection on CH₄ traits, then CH₄ production will increase and CH₄ intensity will decrease (González-Recio et al., 2020). With active selection for CH₄ intensity, CH₄ production will continue to increase but the rate of genetic improvement will be faster in reducing CH₄ intensity (Kandel et al., 2018; Flay et al., 2019). This would reduce the total contribution of dairy cattle to global greenhouse gas production (Van der Maas et al., 2009).

So far, sniffer devices have measured CH₄ for short periods of time, under experimental conditions, or with a limited number of animals or farms. Before a trait can be included in a selection index, there needs to be reasonable confidence in the accuracy of estimated breeding values (EBVs) (Mrode, 2014). With an increasing rate of genotyping and the difficulty in recording CH₄ at an individual cow level, it is probable that any future CH₄ trait will rely on genomic prediction. Genomic prediction will facilitate a shorter time until adoption, assuming there is sufficient accuracy. There is a gap in the literature in regards to how many animals and farms, sniffer recorded CH₄ would need to be measured on, before an acceptable accuracy for genomic prediction is reached, and for the trait to be included in a selection index.

Our objective for this study was to determine the potential effects of including CH₄ production as a trait in the selection index

of the Dutch national breeding objective. Preliminary results of this objective have been published in an abstract form (de Haas et al., 2019). Furthermore, determine how many cows and sniffer devices would be needed to achieve an accuracy suitable to include CH₄ as a trait in such an index.

Material and methods

Dutch national breeding goal

Breeding programmes use selection index theory to ensure maximum gains and that trait changes are in the desired direction. The current selection indexes for the Dutch national breeding goal include 15 traits from the following categories; milk production, milk components, feed efficiency, health, reproduction, longevity, calving, and conformation (CRV, 2018). Table 1 shows the correlations between the traits in the Dutch breeding goal, as well as the heritabilities and the number of daughters with records for that trait.

Genetic parameters for methane production

There are not many papers that have published correlations between CH₄ production and other traits in the breeding goal, but from the literature we obtained a heritability of 0.21 for enteric CH₄ production (Lassen and Løvendahl, 2016), and genetic correlations of 0.4 with milk lactose, fat, protein and DM intake (DMI) (Lassen and Løvendahl, 2016; González-Recio et al., 2020). Correlations between enteric CH₄ production and other traits in the breeding goal were set to zero. Phenotypic SD for CH₄ production was set to 50 g/d.

The Dutch population means were 392 g/d for enteric CH₄ production (Niu et al., 2018), and 9 000 kg milk in 305 days (CRV, 2019). This corresponds to a CH₄ intensity of 15.9 g/d per kg of milk.

Selection index calculations

The spreadsheet with desired gains of Van der Werf (2020) was used to add enteric CH₄ production to the Dutch breeding goal, with six scenarios:

- 1) Adding CH₄ production to the breeding goal with no weight and no records on daughters of sires (=current trend);
- 2) Adding CH₄ production to the breeding goal with no economic weight and increasing number of records on daughters of sires;
- 3) Adding CH₄ production to the breeding goal with an economic weight based on the carbon price and increasing number of records on daughters of sires;
- 4) Adding CH₄ production to the breeding goal with the restriction of no genetic gain in the CH₄ produced per cow;
- 5) Adding CH₄ production to the breeding goal with an active selection on reducing the CH₄ production per cow;
- 6) Simulating a breeding goal that only focusses on reducing the enteric CH₄ production per cow, using a maximum amount of available information.

Scenarios 4 and 5 were achieved by adjusting the economic weights that were put on CH₄ production per cow.

Genetic trend

From the spreadsheet with desired gains of van der Werf (2020), genetic gains for all traits, including the enteric CH₄ production, were obtained. Based on the annual overviews that are

Table 1

Heritabilities of (on diagonal) and genetic correlations (below diagonal) between the traits in the Dutch national breeding goal for dairy cattle and number of daughters (#daughs) with records for that trait (adapted from: https://www.crv4all-international.com/wp-content/uploads/2018/06/E_20-NVL_apr2018_en.pdf).

Traits	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	#daughs
1. Lactose	0.43															125
2. Fat	0.38	0.58														125
3. Protein	0.88	0.58	0.50													125
4. Longevity	0.36	0.35	0.42	0.12												125
5. Udder Health	-0.03	-0.02	-0.06	0.36	0.09											125
6. Int-first-last ¹	-0.34	-0.24	-0.29	0.25	0.27	0.08										125
7. Calving Interval	-0.44	-0.33	-0.37	0.11	0.21	0.85	0.15									100
8. Udder conformation	-0.08	-0.04	-0.10	0.11	0.27	-0.05	0.00	0.34								75
9. Feet_Leg	0.02	0.04	0.05	0.25	0.21	0.00	0.00	0.35	0.17							75
10. direct_calving_ease	0.07	0.15	0.11	0.24	0.15	0.20	0.24	0.00	0.00	0.07						200
11. maternal_calving_ease	0.00	0.00	0.00	0.16	0.09	0.25	0.24	0.10	0.10	0.19	0.05					75
12. direct_vitality	0.05	0.09	0.02	0.14	0.05	0.10	0.14	0.00	0.00	0.60	0.14	0.04				400
13. maternal_vitality	-0.04	-0.07	0.03	0.16	0.07	0.32	0.24	0.00	0.00	0.11	0.34	-0.16	0.09			140
14. Claw health	0.00	0.15	0.07	0.33	0.09	0.10	0.14	0.15	0.65	0.16	0.06	0.03	0.10		0.18	15
15. saved feed costs	0.20	0.35	0.30	0.50	-0.03	-0.10	-0.30	-0.09	-0.29	0.41	-0.20	0.17	-0.05	0.11	0.25	20

¹ Int-first-last: interval between first and last insemination.

provided by the national herdbook CRV, Arnhem, the Netherlands (CRV, 2020), the achieved gains per year are known, and based on the obtained and achieved gain for kg fat production, we scaled the obtained gain for the enteric CH₄ production back to yearly genetic trend.

Carbon price

The economic value of CH₄ production used for the selection indexes was based on the shadow price of carbon dioxide (CO₂). The economic value was based on the expected shadow price for CO₂ for 2025 obtained from literature (DBEIS, 2017). The shadow price for CO₂ used for the calculation of the shadow price of CH₄ production was obtained from the UK Government report on “Updated short-term traded carbon values used for modelling purposes” (DBEIS, 2017). This was €36.19/tCO₂e, current exchange rate at the time of analyses.

The following formula was then used to calculate the economic values of CH₄ production;

$$\text{Economic value} = -1 * (\text{shadow price of CO}_2 * 28/1\,000\,000) * 365$$

where the multiplication by -1 was due to the direction of selection on CH₄ production being negative, shadow price of CO₂ was as described above, 28 represents the global warming potential of CH₄ and was used to convert from CO₂ shadow price to CH₄ shadow price, division by 1 000 000 converts from grams to tonnes, and multiplication by 365 was because the index calculates gain per year and CH₄ data are in grams per day, in order to align the economic value of CH₄ production with those for the other traits in the breeding goal that are also expressed on an annual basis. This resulted in an economic value of -€0.37 g/d for a year.

Genomic prediction

For scenarios 3 to 6, where there is a desire, or financial incentive, to limit or reduce CH₄ intensity, selection with genomic prediction was tested. To include genomic selection in the index, a pseudo-trait was added. This trait had a heritability of 0.99, and a genetic correlation with CH₄ production of 0.63. This genetic correlation was used based on the requirement that for a trait to be included in the national breeding goal, the reliability of prediction should be equal to or greater than 0.40 (Koenen E., Personal communication, 2019). For selection with genomic prediction, we assumed no daughters were phenotyped for the CH₄ production trait, but genotyped cows had their own performance for enteric CH₄ emission recorded.

Power calculations for recording enteric methane

The parameters for a typical dairy cow population were used to determine the number of animals, farms, sniffer devices, and the number of measurements recorded with the sniffer devices, required for a suitable reliability of genomic prediction. The parameters included; the number of markers (60 000), effective population size (Ne, 120), average chromosome length (L, 100 cM), number of chromosomes (k, 30). The effective number of chromosome segments (Me, 7 200) was calculated as 2NeLk. The total number of individuals with CH₄ recorded in the reference population (T) ranged between 1 000 and 15 000 cows. The accuracy of genomic prediction (r_{gg}) was estimated following the first equation of Daetwyler et al. (2008):

$$r_{gg} \hat{=} \sqrt{\frac{\lambda h_o^2}{\lambda h_o^2 + 1}}$$

where h_o^2 is the observed heritability (0.21), and λ is the ratio of the number of phenotypic observations and the number of loci associated (Me/T). Reliability of genomic prediction was then estimated as r_{gg}^2 .

Different recording scenarios were considered over 3 years of recording, with 100 or 50 sniffer devices available (Table 2). For both 100 and 50 sniffer devices, the units were installed for either 3 years of repeated records on the original farm, 1 year of recording on one farm and then 2 years of recording on a second farm, or every year, the unit is moved to a new farm. Each farm was assumed to have a herd size of 150 genotyped cows with a replacement rate of 0.3. Only one automatic milking station was equipped with a sniffer device per farm, but all 150 cows would have access to the sniffer.

Depending on the manufacturer, sniffer devices provide a CH₄ measurement approximately every 10 s. It was assumed that the measurements during one visit to the milking robot would be used to calculate a CH₄ record for that visit. Due to the lower accuracy of sniffers compared to other devices, it is unlikely that fitting CH₄ for each visit or even daily would be useful. A trait could be calculated from the available recorded visits, the simplest would be a mean of these visit records over some period of time. As the sniffer devices are installed inside milking robots, the longest period for one phenotypic observation would be one full lactation period. This would provide one phenotypic observation per year. There are biological differences during lactation stages which affect CH₄ production (Van Engelen et al., 2018), therefore, calculating one phenotypic

Table 2
Summary of scenarios tested and the predicted reliability of an individual's genotype based on the mean number of repeated records per cow.

Sniffer installation	Repeated records	Number of			Cows on all farms	Total records on all farms	Records per cow	Reliability of GEBV ¹
		Unique farms	Unique cows per farm	Mean records per farm				
Three years of repeated records per farm								
100 sniffer devices								
Year 1	No	100	150	150	15 000	15 000	1.0	0.20
Year 2	Yes	100	195	300	19 500	30 000	1.5	0.26
Year 3	Yes	100	240	450	24 000	45 000	1.9	0.29
50 sniffer devices								
Year 1	No	50	150	150	7 500	7 500	1.0	0.20
Year 2	Yes	50	195	300	9 750	15 000	1.5	0.26
Year 3	Yes	50	240	450	12 000	22 500	1.9	0.29
Device moved once after first year								
100 sniffer devices								
Year 1	No	100	150	150	15 000	15 000	1.0	0.20
Year 2	No	200	150	150	30 000	30 000	1.0	0.20
Year 3	Yes	200	195	225	34 500	45 000	1.3	0.24
50 sniffer devices								
Year 1	No	50	150	150	7 500	7 500	1.0	0.20
Year 2	No	100	150	150	15 000	15 000	1.0	0.20
Year 3	Yes	100	195	225	17 500	22 500	1.3	0.24
Device move to a new farm every year								
100 sniffer devices								
Year 1	No	100	150	150	15 000	15 000	1.0	0.20
Year 2	No	200	150	150	30 000	30 000	1.0	0.20
Year 3	No	300	150	150	45 000	45 000	1.0	0.20
50 sniffer devices								
Year 1	No	50	150	150	7 500	7 500	1.0	0.20
Year 2	No	100	150	150	15 000	15 000	1.0	0.20
Year 3	No	150	150	150	22 500	22 500	1.0	0.20

¹ GEBV: genomic estimated breeding value.

observation per lactation period (i.e., early, mid, late) could be useful, this would provide three repeated records per year (assuming lactation stage was fitted as a fixed effect and not a separate trait). It was assumed that at least 100 days per period would be used in the calculation of one observation, allowing for: errors, days where the sniffer device needs maintenance, potential that the cow does not visit the robot with the device installed, and reproductive management. Shorter periods could be more useful and provide more repeated records, with two observations per lactation period, six trait observations per year would be available with at least 50 days included in the calculation of each observation. Finally using observations of just 10 days, which is in line with the previous sniffer studies (Lassen and Løvendahl, 2016; López-Paredes et al., 2020), would provide at least 30 observations per year.

Recording on the same farm for multiple years, or using shorter recording periods per phenotypic observation, has the potential to improve the reliability of prediction. With repeated records, the reliability of prediction for an individual's genotype was estimated following the method of Van Vleck (1993), as implemented by Tenghe et al. (2018):

$$\text{Reliability of individual's phenotype} = \frac{mh^2}{(m-1)t+1}$$

where m is the number of records per cow, h^2 is the heritability (0.21) used in the selection index calculations, and t is the repeatability (0.35, 0.50, or 0.65). Reliability of genomic prediction was then estimated as previously with the Daetwyler et al. (2008) equation with h_o^2 replaced with the reliability of the individual's phenotype.

Results

Selection index theory

Selection index calculations show how much the traits are predicted to change per year. This is plotted in Fig. 1 for both CH₄ production (g/d) and CH₄ intensity (CH₄ production expressed per kg of milk). It shows that average annual CH₄ production per cow will




Fig. 1. Simulated genetic trends in enteric methane production (solid line) and methane emission intensities (methane emission per kg milk; dashed line) with the current breeding goal for dairy cattle.

steadily increase by 1.5 g/d as a correlated response to selection for the current breeding goal. However, the CH₄ intensity drops. When extrapolating these, the average daily CH₄ production of 392 g/d will increase to 442 g/d in 2050 (+13%). However, in the same period, the CH₄ intensity will reduce by 13%.

Further reductions can be achieved when actively selecting lower methane-emitting cows, by adding more weight on CH₄ in the national breeding goal (Fig. 2). By putting economic weight on CH₄ production in the breeding goal, selective breeding can reduce the CH₄ intensity by 24% in 2050. This shows that breeding is a valuable contribution to the whole set of mitigation strategies that could be applied in order to achieve the goals for 2050 set by the EU. Selecting actively against CH₄ would result in healthy, fertile, long-living cows that emit less CH₄. Actively selecting against CH₄ emission, however, requires large-scale recording of individual CH₄ emissions.

Comparison of the scenarios (Table 3) shows that when no economic weight is put on CH₄ production, adding 10, 20, 50 or 100 daughters with CH₄ records does not have a large impact on the genetic trend. Any small impact on the genetic trend in this scenario (no economic weight) is probably due to small changes in selection pressure caused by the traits that do have an economic value and are correlated with CH₄, and the small amount of extra information the daughters provide. For the proceeding scenarios where there is an incentive for selecting CH₄, genomic prediction provided similar results as sires with 20 progeny. Adding an economic weight on CH₄ production based on the carbon price does slow down the increase in CH₄ production as a correlated response with ~1 g/d (i.e., difference scenario 1 (5.79 g/d) and scenario 2 (4.72 g/d) in Table 3) in 3.75 years, but still, the CH₄ production will increase per year, and the CH₄ intensity will decrease. With a breeding goal aiming for no gain in CH₄, the genetic trend for milk production drops by 70–130 kg, depending on the number of daughters with records for CH₄ production per sire. Achieving no gain in CH₄ can only be realized with high economic values of –1 to –1.8, which corresponds to carbon prices between 100 and 175€/tCO₂e. With a breeding goal aiming for a reduction in CH₄ production, but still taking into account the other breeding goal traits; i.e., selecting for healthy, fertile, long-living cows that emit less CH₄, the genetic trend for milk production drops by 170–280 kg, depending on the number of daughters with records for CH₄ production per sire. The economic values will have to be –2 to –3.6, which corresponds to carbon prices between 195 and 350€/tCO₂e. Finally, an unrealistic breeding goal was assumed with

only a focus on CH₄ reduction, to show what breeding could potentially achieve. This maximal reduction in CH₄ production was determined based on a scenario with no negative genetic trend in milk production when sires had at least 100 daughters with records. A genetic trend of –12.75 g/d CH₄ production in 3.75 years can then be achieved, resulting in a 29% reduction in 2050 (Fig. 2).

Power calculations

To achieve a reliability of genomic prediction above 0.40 (Fig. 3) would require approximately 25 000 genotyped cows to have a CH₄ record. When considering using sniffer devices over multiple years, the same number of measurements is recorded regardless of the strategy (22 500 records for 50 sniffer devices and 45 000 records for 100 sniffer devices), but the number of cows recorded is different. Capturing repeated records by leaving 50 sniffer devices on the same 50 farms for 3 years limits the number of cows recorded (12 000 cows), and extends the time until the required threshold would be reached. Only if the 50 sniffer devices were moved to new farms each year (150 total), would there be close to enough cows with a CH₄ record (22 500), even with a high trait repeatability of 0.65, the reliability of prediction (0.39) is still only approaching the required 0.40. With 100 sniffer devices, a reliability of 0.40 is achieved in the second year regardless of whether sniffer devices are moved or not. Having repeated records on 100 farms is the only scenario where the number of cows recorded (19 500) is fewer than 25 000 but still offers sufficient reliability (0.41).

So far only one observation per cow per lactation was considered. Reducing the observation period increases the number of repeated records per cow, thereby increasing the reliability of genomic prediction (Fig. 3). With a trait repeatability of 0.35 or 0.50, the scenarios with 100 sniffer devices achieved the threshold in the first year when using observations from three lactation periods, 50 days, or 10 days (3, 6, and 30 observations in a year, respectively). At the higher trait repeatability of 0.65, only three lactation periods required an additional year of recording before reaching a reliability of 0.40. When considering the 50 farms and a trait repeatability of 0.35, both observation periods of 50 and 10 days achieved the 0.40 reliability threshold after 2 years of recording, and three observations approached the threshold (0.37) but did not achieve it until the third year. However, this is a considerable improvement compared to the scenario with only one observation a year. Similar results were observed with the higher trait repeatabilities.

Discussion

In this study, we showed that with the current Dutch breeding goal, the genetic trend in CH₄ production is increasing, but the genetic trend in CH₄ intensity is decreasing. Actively selecting on lower emitting dairy cows will decrease this trend even further. In order to actively select on lowering the enteric CH₄ emission of dairy cattle by breeding, data on 100 farms (with on average 150 cows each) need to be collected for at least 2 years, in order to achieve the desired reliability of the genomic prediction.

Impact of breeding

Technical solutions to reduce enteric CH₄ emissions have been, and continue to be, extensively researched. Animal breeding exploits natural animal variation in CH₄ emissions, and we have shown that it is an additional mitigation strategy that is cost-effective, permanent, and cumulative. A continuous reduction in CH₄ intensity through breeding for low(er) emitting cows will help




Fig. 2. Simulated genetic trends in methane emission intensities (methane emission per kg milk) of dairy cattle in three situations: current trend (solid), combined selection for methane and other relevant breeding goal traits (small dash), theoretical maximum when exclusively focussing on methane (large dash).

Table 3

Genetic trends per generation interval of 3.75 years in enteric methane (CH₄) production (g/d) and in milk production (kg), of dairy cattle with the implicit economic value of CH₄ and the maximal reduction in methane intensity (g CH₄/kg milk) that could be achieved in 2050 compared to 2017 with this scenario.

Scenario		Genetic trend CH ₄ production (g/d)	Genetic trend milk production (kg)	Economic value CH ₄	Max reduction in CH ₄ intensity (%)
1	current breeding goal: no weight, 0 daughters	5.79	307.06	0.00	13
2a	no weight, records on 10 daughters	5.66	306.85	0.00	13
2b	no weight, records on 20 daughters	5.58	306.72	0.00	13
2c	no weight, records on 50 daughters	5.47	306.54	0.00	14
2d	no weight, records on 100 daughters	5.39	306.41	0.00	14
3	economic weight of carbon price (0 daughters)	4.72	285.25	-0.37	14
3a	economic weight of carbon price (10 daughters)	4.33	284.37	-0.37	14
3b	economic weight of carbon price (20 daughters)	4.08	283.82	-0.37	15
3c	economic weight of carbon price (50 daughters)	3.70	282.98	-0.37	15
3d	economic weight of carbon price (100 daughters)	3.43	282.39	-0.37	16
3e	economic weight of carbon price (genomic prediction)	4.29	284.99	-0.37	14
4	no genetic gain (0 daughters)	0.00	178.55	-1.82	15
4a	no genetic gain (10 daughters)	0.00	202.93	-1.47	17
4b	no genetic gain (20 daughters)	0.00	214.28	-1.30	17
4c	no genetic gain (50 daughters)	0.00	227.72	-1.10	18
4d	no genetic gain (100 daughters)	0.00	235.06	-0.99	19
4e	no genetic gain (genomic prediction)	0.00	212.83	-1.36	17
5	active selection on CH ₄ reduction (0 daughters)	-5.79	26.65	-3.65	15
5a	active selection on CH ₄ reduction (10 daughters)	-5.79	75.16	-2.98	19
5b	active selection on CH ₄ reduction (20 daughters)	-5.79	98.22	-2.66	21
5c	active selection on CH ₄ reduction (50 daughters)	-5.79	126.03	-2.27	23
5d	active selection on CH ₄ reduction (100 daughters)	-5.79	141.37	-2.06	24
5e	active selection on CH ₄ reduction (genomic prediction)	-5.79	97.52	-2.73	21
6	selection on maximal CH ₄ reduction (0 daughters)	-12.75	-197.76	-7.85	11
6a	selection on maximal CH ₄ reduction (10 daughters)	-12.75	-114.72	-5.93	20
6b	selection on maximal CH ₄ reduction (20 daughters)	-12.75	-75.61	-5.15	23
6c	selection on maximal CH ₄ reduction (50 daughters)	-12.75	-25.64	-4.27	27
6d	selection on maximal CH ₄ reduction (100 daughters)	-12.75	1.58	-3.81	29
6e	selection on maximal CH ₄ reduction (genomic prediction)	-12.75	-74.69	-5.26	23




Fig. 3. Reliability of genomic prediction as a predictor of the true breeding value for methane emission of dairy cattle with 50 or 100 sniffer devices available (left or right, respectively), and different recording scenarios, estimated following Daetwyler et al. (2008). Circle; Repeated records with the same farms recorded on each year, Triangle; Repeated records with the same farms in the first and second year but sniffer devices are moved to new farms in the third year, Square; Sniffer devices are moved to a new farm every year with no repeated records.

in reaching the targets set at the Paris COP meeting in 2015 (Framework Convention on Climate Change - United Nations., 2015).

The calculated impact of breeding depends on the chosen parameters. For Dutch data, no parameter estimations exist yet, so we collated information from the literature. A heritability of 0.21 is well within the range of reported heritabilities in dairy cattle, sheep or beef cattle (Pickering et al., 2015b; Brito et al., 2018). A recent review of Lassen and Difford (2020) collated genetic correlations between CH₄ emission traits and existing selection index traits in dairy cattle. The genetic correlations with milk production traits ranged between 0.37 and 0.61, with a low sSE (0.07) on the lowest estimate large SE on the highest estimate (0.32) (Lassen and Løvendahl, 2016; Difford et al., 2018; Breider et al., 2019). Genetic correlations with feed efficiency traits were estimated by Difford et al. (2020) for two populations, and shown to be inconsistent across both countries and trait definitions for feed efficiency, ranging between -0.69 and 0.69 with large SEs up to -0.4. Our assumed correlations on 0.4 with milk production traits and -0.4 with saved feed costs fit well within the range of published results. Lassen and Difford (2020) reviewed also the genetic correlations with other breeding goal traits, such as reproduction and health, and concluded that selection for reduced CH₄ emissions has minimal consequences on these traits, but analyses of larger datasets are needed to confirm or deny the genetic correlation structure of other traits. This is also confirmed in a Spanish study on data from 1 500 animals from 14 commercial farms (López-Paredes et al., 2020). Assuming a genetic correlation of zero, as we did, does therefore fit with the current knowledge.

When looking at a dairy system level, however, Wall et al. (2010) showed that selection for improved fitness traits (e.g., lifespan, health, fertility) will help to reduce emissions by reducing wastage of animals. Improving lifespan in dairy cows will reduce greenhouse gas emissions of the system by reducing the number of followers required to maintain the herd at a given size. Also, Van Middelaar et al. (2015) showed that improving the longevity of cows with one genetic SD is important when aiming for mini-

mizing the greenhouse gas emissions. Garnsworthy (2004) estimated, via modelling, that if cow fertility was restored to 1 995 levels from 2 003 levels, CH₄ emissions from the dairy industry would reduce by 10% to 15%. Therefore, correlated responses of national breeding goals can be expected, but the genetic correlation structure still has to be confirmed (or denied) in large studies.

Breeding goal

Breeding has the potential to reduce enteric CH₄ emissions from dairy cows, but how it should be placed in the context of breeding goals is still under debate. Breeding goal traits to genetically reduce CH₄ emissions in dairying are (1) CH₄ production as a mass flux rate per day (in litres or grams per day), (2) methane yield, which is CH₄ production divided by feed intake (e.g., CH₄ production per kg DMI, (3) methane intensity per unit product (e.g., CH₄ production per kg milk yield), and (4) residual methane production (e.g., CH₄ regressed on DMI, BW and milk yield) (de Haas et al., 2017). The European climate goals are to reduce the overall greenhouse gas emission, and therefore, CH₄ production is the most appropriate breeding goal trait, and not any of the ratio traits. Expressing it as a ratio, is, however, a useful metric to describe groups of animals; e.g. per breed, per herd, per country (Gerber et al., 2013), but the goal should be to lower the CH₄ emission per animal.

Current international dairy breeding goals involve the improvement of multiple traits simultaneously, usually compiled in a total merit index to ensure balanced breeding. Total merit indices consist of many traits, and enteric CH₄ might in the (near) future be one of them. Selecting based on a total merit index avoids establishing extreme divergent selection lines for e.g. low and high methane emitters. Selecting one trait can result in unforeseen, and possibly unwanted, correlated effects, like the rumen differences shown in two divergent selection lines of sheep in New Zealand (Bain et al., 2013). They showed that the rumen content, volume and surface area differed significantly between high and low emitting sheep, with low emitting sheep having a smaller rumen than high emitting sheep.

Traits in a total merit index are weighed to maximize economic gain, and those weights are either based on economic values or on desired gains. At this moment, however, no carbon-pricing scheme exists for dairy cattle farming in the Netherlands. We therefore used the expected shadow price for CO₂ for 2025 obtained from literature (DBEIS, 2017) as a proxy economic value for enteric CH₄ emissions. It was shown that this carbon price was not enough to realize a negative genetic trend for enteric CH₄ emissions of dairy cattle; in fact, the carbon price had to be 10 times higher (36€ vs 350€). It is therefore questionable whether carbon pricing alone is sufficient to mitigate climate change (Rosenbloom et al., 2020). The economic value was not only based on the expected shadow price for CO₂, but also on the global warming potential (GWP) of CH₄. For this, the GWP100 was used, which refers to the GWP over 100 years. However, recent studies have modified the use of GWP so that it accounts for the differences between short- and long-lived gases, which is referred to as GWP* (Cain et al., 2019; Lynch et al., 2020). Enteric CH₄ emissions of cattle are classified as short-lived gases, these gases do not build up in the atmosphere, so their warming potential is shorter lived than of long-lived gases (like fossil CO₂). This results in a much higher GWP for CH₄ than 28 CO₂-equivalents in the short term, and a lower GWP after that (Lynch et al., 2020). On average, the GWP* of CH₄ will be lower than the GWP100. When using GWP*, the calculated economic value for CH₄ production will be lower with the same expected shadow price for CO₂, which will reduce the impact of genetic selection.

Large-scale recording of enteric methane

Genomic prediction offers the shortest period of time and similar rates of gain (compared to progeny testing) before selection for lower CH₄ production can begin. If the decision is made to implement genomic breeding strategies to reduce enteric CH₄ production, and to achieve the expected breeding impact, there needs to be a sufficient reliability of prediction. The only way to achieve that is to have enough animals in a reference population phenotyped and genotyped. The power calculations offer insights into the difficulties that will be faced in trying to record enough data, but also provide optimism with potential methods of increasing the reliability of prediction and reducing the time period before selection can begin. Importantly, it is possible to obtain sufficient reliability within a relatively short period of time.

Before a trait is included in national evaluations, the reliability of prediction needs to be sufficiently high for accurate predictions and confidence in the values. In this example, we assumed a trait would not be included unless a threshold of 0.40 was achieved. By recording CH₄ on 100 farms, this threshold could be achieved in just 2 years, with phenotypic information on 19 500 cows collected in that time period. Of course, there are many factors that increase or decrease the required cows, farms, and time periods, but this is a useful indicator as no matter which parameters are used, a very large number of cows and farms will need to be measured.

There are now many methods of recording CH₄ at a cow level, but the options for recording at such a large scale are limited. Respiration chambers, while the most accurate method, have a low throughput, and a high cost (Hammond et al., 2015). The use of devices that measure CH₄ and flux, such as GreenFeed™ (Hammond et al., 2015), has been very beneficial for increasing the number of cows recorded and providing a reliable measurement. However, on such a large scale, they would not be logistically or economically viable. In most scenarios, such devices record 30 cows during a 2 week period (Hammond et al., 2015; Sorg et al., 2018). Therefore, multiple devices would be required per farm, or management strategies and infrastructure changes implemented to record on all cows. The use of handheld lasers would provide a means of increasing the number of cows recorded, however, they only offer a spot sample, require a large amount of labour, and have a lower accuracy (Pickering et al., 2015a; Sorg et al., 2018). It is unlikely that the use of reference gases such as SF₆ would be used at a large scale, due to animal welfare concerns caused by the invasive nature of the current methods (Deighton et al., 2014).

Sniffers are currently the only device that provides the convenience required for large-scale recording, at a reasonable cost point, while maintaining sufficient reliability of measurements, but there are still concerns regarding the limited validation of the method (Difford et al., 2018; Sorg et al., 2018; Garnsworthy et al., 2019). Sniffer devices could provide the largest amount of cows with CH₄ records, but there is a possibility of supplementing them with the other devices. It would be easiest to combine data sources that provide CH₄ measurements with the same units, for sniffers that would be CH₄ concentration (in ppm). The largest difficulty would be ensuring that the same trait is being measured. Sniffers offer a method of long-term recording, capturing data at all points of the lactation curve. While it would be possible to combine data sources from different methods, they do not capture the variation caused by various fixed effects, it is unknown what effect this will have on the reliability of prediction.

If other data from different devices are considered, a national breeding programme could also be supplemented with international data. This is already done with other traits, but relies on the fact that countries have a good genetic linkage, and enough

animals recorded with similar reliabilities of genomic EBVs (Dürr and Philipsson, 2012; Van Eenennaam et al., 2014). At the moment, CH₄ recording across countries is a patchwork of recording methods, data set sizes, and production systems. No country currently has enough data to meet the requirements that would be needed for sharing. This could be overcome if countries decided to collaborate in a large-scale recording programme and combining resources. At the moment, no such programme exists and countries continue to work independently. It is most likely that a small number of national breeding programmes implement some form of large-scale recording and connections are made between them at a later date.

If data sources from other methods, or countries, are not used to obtain a sufficient reliability of prediction, there are still ways that the amount of recording can be reduced. One method to achieve a 0.40 reliability of prediction is to increase the reliability of an individual's recorded breeding value. One method of doing this is adding repeated records. So far, we have only discussed a conservative one record for a full lactation period, or one potential repeated record per year. The test period could be reduced to the lactation period, to 50 days, or even shorter to 10 day periods, which would align with other recording methods and other traits.

Increasing the number of repeated measurements is the most convenient method to increase reliability of prediction, a larger increase can be achieved by recording on more cows and farms. With the conservative test period of one record per lactation, recording on new farms is beneficial if the number of sniffer devices is limited by availability. With 50 sniffer devices available and recording on a new farm each year, the threshold is almost reached within 3 years. If it is appropriate to reduce the test period, and there was an ability to record on new farms after each period, the threshold would be reached after four recording periods, or within 2 to 3 years. If the number of sniffer devices is above 100, there is no need to move the sniffers and repeated records are sufficient.

The uncertainty regarding test periods is just one limitation of the power analysis. Other assumptions were also made, such that all animals recorded were also genotyped. The rate of genotyping continues to increase, for these results to be relevant, the farms selected for measuring would need to be targeted based on if the herd is genotyped, otherwise an additional cost is required to have them genotyped. That would be achievable with the current systems; otherwise, the time to the threshold would be extended. Another assumption was farm size, 150 cows per farm were used for this analysis, more important is the total number of cows recorded. Assuming the mean herd size is at least 150, this would not be an issue and could be used as another target alongside genotyping when selecting farms to be included in recording.

There are some assumptions in the index and power calculation that could underestimate the rate of gain when using genomic prediction. With a sufficient reference population, genomic selection can begin earlier and would have shorter generation intervals without a need for daughters to mature. Importantly, we did not consider that some sires could still have progeny tested daughters with genomic selection.

The relevance of this work is a demonstration of the fact that, for an enteric CH₄ trait to be included in a national breeding goal, there needs to be a large and targeted recording scheme. The parameters used were either means or a range of values. For example, a lower or higher heritability could have been used instead of the literature mean. However, the effect of selecting other values within the range of published estimates would have a limited impact on the final outcome, a very large number of farms and cows would still need to be recorded.

Conclusions

The global livestock sector, particularly ruminants, contributes substantially to total anthropogenic greenhouse gas emissions. Management and dietary solutions to reduce enteric CH₄ emissions are extensively researched. Animal breeding that exploits natural variation in CH₄ emissions is an additional mitigation solution that is cost-effective, permanent, and cumulative. By adding CH₄ production to the Dutch breeding goal, selective breeding can reduce the CH₄ intensity by 24% in 2050. This shows that breeding is a valuable contribution to the whole set of mitigation strategies that could be applied in order to achieve the goals for 2050 set by the EU. If the decision is made to implement animal breeding strategies to reduce enteric CH₄ production, and to achieve the expected breeding impact, there needs to be a sufficient reliability of prediction. The only way to achieve that is to have enough animals phenotyped and genotyped. The power calculations offer insights into the difficulties that will be faced in trying to record enough data. Recording CH₄ data on 100 farms (with on average 150 cows each) for at least 2 years is required to achieve the desired reliability of 0.40 for the genomic prediction.

Ethics approval

This study is based on a data simulation (with input from literature), therefore, no data was collected on animals and no ethics statement is needed.

Data and model availability statement

None of the data were deposited in an official repository. Data and model are available upon request, the software used is available to reviewers.

Author ORCIDs

Y. de Haas: <https://orcid.org/0000-0002-4331-4101>
 R.F. Veerkamp: <https://orcid.org/0000-0002-5240-6534>
 G. de Jong: <https://orcid.org/0000-0001-9365-5089>
 M.N. Aldridge: <https://orcid.org/0000-0002-9033-3081>

Author contributions

Conceptualization: **Yvette de Haas (YdH), Roel F. Veerkamp (RFV), Gerben de Jong (GdJ), Michael N. Aldridge (MNA)**; Methodology: **YdH, RFV, GdJ, MNA**; Software: **MNA**; Formal analysis: **YdH, RFV, GdJ, MNA**; Visualization: **YdH, MNA**; Writing – original draft: **YdH, MNA**; Writing – review & editing: **YdH, RFV, GdJ, MNA**; Funding acquisition: **YdH, RFV**. All authors have read and agreed to the published version of the manuscript.

Declaration of interest

The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Acknowledgements

The authors thank Erwin Koenen and René van der Linde (CRV, Arnhem, the Netherlands) for their input in the discussions.

Financial support statement

This study was financially supported by the Dutch Ministry of Economic Affairs (TKI Agri & Food project 16022) and the Breed4-Food partners Cobb Europe, CRV, Hendrix Genetics and Topigs Norsvin.

Transparency Declaration

This article is part of a supplement entitled 'Sustainable livestock systems for high-producing animals', supported by the European Federation of Animal Science (EAAP) and the World Association for Animal Production (WAAP).

References

Bain, W.E., Bezuidenhout, L., Jopson, N.B., Pinares-Patiño, C.S., McEwan, J.C., 2013. Ruminal differences between sheep identified as being low or high emitters of greenhouse gas. In: Proceedings of the 20th Association for the Advancement of Animal Breeding and Genetics, 20-23 October 2013, Napier, New Zealand, pp. 376-378.

Beauchemin, K., Kreuzer, M., O'mara, F., McAllister, T., 2008. Nutritional management for enteric methane abatement: a review. *Australian Journal of Experimental Agriculture* 48, 21-27.

Bird-Gardiner, T., Donoghue, K.A., Arthur, P.F., Herd, R.M., Hegarty, R.F., 2015. Divergent selection for methane yield in beef cattle. In: Proceedings of the 21st Association for the Advancement of Animal Breeding and Genetics, 28-30 September 2015, Lorne, Australia, pp. 122-125.

Breider, I.S., Wall, E., Garnsworthy, P.C., 2019. Heritability of methane production and genetic correlations with milk yield and body weight in Holstein-Friesian dairy cows. *Journal of Dairy Science* 102, 7277-7281.

Brito, L.F., Schenkel, F.S., Oliveira, H.R., Cánovas, A., Miglior, F., 2018. Meta-analysis of heritability estimates for methane emission indicator traits in cattle and sheep. In: Proceedings of the 11th World Congress on Genetics Applied to Livestock Production, Volume Challenges-Environmental, 11-16 February 2018, Auckland, New Zealand, p. 740.

Cain, M., Lynch, J., Allen, M.R., Fuglestedt, J.S., Frame, D.J., Macey, A.H., 2019. Improved calculation of warming-equivalent emissions for short-lived climate pollutants. *Nature Partner Journals Climate and Atmospheric Science* 2, 1-7.

CRV, 2018. Statistical Indicators. Retrieved on 18 July 2019 from https://www.crv4all-international.com/wp-content/uploads/2018/06/E_20-NVI_apr2018_en.pdf.

CRV, 2019. Bedrijven en koeien in cijfers - Nederland. Retrieved on 14 October 2020 from <https://publicaties.cooperatie-crv.nl/bedrijven-en-koeien-in-cijfers.html>.

CRV, 2020. Jaarstatistiek. Retrieved on 23 October 2020 from <https://www.crv4all.nl/downloads/prestaties/jaarstatistiek/>.

Daetwyler, H.D., Villanueva, B., Woolliams, J.A., 2008. Accuracy of predicting the genetic risk of disease using a genome-wide approach. *PLoS one* 3, e3395.

DBEIS, 2017. Updated energy and emissions projections: 2017. Retrieved on 23 October 2020 from <https://www.gov.uk/government/publications/updated-energy-and-emissions-projections-2017>.

de Haas, Y., Koenen, E., Veerkamp, R.F., 2019. Selective breeding as a mitigation tool for methane intensity of dairy cattle. In: Proceedings of the 7th Greenhouse Gas and Animal Agriculture Conference, 4-8 August 2019, Iguassu Falls, Brazil, p. 43.

de Haas, Y., Pszczola, M., Soyeurt, H., Wall, E., Lassen, J., 2017. Invited review: phenotypes to genetically reduce greenhouse gas emissions in dairying. *Journal of Dairy Science* 100, 855-870.

Deighton, M.H., Williams, S.R.O., Hannah, M.C., Eckard, R.J., Boland, T.M., Wales, W.J., Moate, P.J., 2014. A modified sulphur hexafluoride tracer technique enables accurate determination of enteric methane emissions from ruminants. *Animal Feed Science and Technology* 197, 47-63.

Difford, G.F., Løvendahl, P., Veerkamp, R.F., Bovenhuis, H., Visker, M.H.P.W., Lassen, J., de Haas, Y., 2020. Can greenhouse gases in breath be used to genetically improve feed efficiency of dairy cows?. *Journal of Dairy Science* 103, 2442-2459.

Difford, G.F., Olijhoek, D.W., Hellwing, A.L.F., Lund, P., Bjerring, M.A., de Haas, Y., Lassen, J., Løvendahl, P., 2018. Ranking cows' methane emissions under commercial conditions with sniffers versus respiration chambers. *Acta Agriculturae Scandinavica, Section A—Animal Science* 68, 25-32.

Dürr, J., Philipsson, J., 2012. International cooperation: The pathway for cattle genomics. *Animal Frontiers* 2, 16-21.

Flay, H.E., Kuhn-Sherlock, B., Macdonald, K.A., Camara, M., Lopez-Villalobos, N., Donaghy, D.J., Roche, J.R., 2019. Hot topic: Selecting cattle for low residual feed intake did not affect daily methane production but increased methane yield. *Journal of Dairy Science* 102, 2708-2713.

Framework Convention on Climate Change - United Nations, 2015. Adoption of the Paris agreement. In Proceedings of United Nations Framework Convention on Climate Change, 30 November - 11 December 2015, Paris, France, pp. 1-27.

Garnsworthy, P.C., 2004. The environmental impact of fertility in dairy cows: a modelling approach to predict methane and ammonia emissions. *Animal Feed Science and Technology* 112, 211-223.

Garnsworthy, P.C., Difford, G.F., Bell, M.J., Bayat, A.R., Huhtanen, P., Kuhla, B., Lassen, J., Peiren, N., Pszczola, M., Sorg, D., 2019. Comparison of methods to measure methane for use in genetic evaluation of dairy cattle. *Animals* 9, 837.

Gerber, P.J., Steinfeld, H., Henderson, B., Mottet, A., Opio, C., Dijkman, J., Falucci, A., Tempio, G., 2013. Tackling climate change through livestock: a global assessment of emissions and mitigation opportunities. Food and Agriculture Organization of the United Nations, Rome, Italy.

González-Recio, O., López-Paredes, J., Ouatahar, L., Charfeddine, N., Ugarte, E., Alenda, R., Jiménez-Montero, J.A., 2020. Mitigation of greenhouse gases in dairy cattle via genetic selection: 2. Incorporating methane emissions into the breeding goal. *Journal of Dairy Science* 103, 7210-7221.

Hammond, K.J., Humphries, D.J., Crompton, L.A., Green, C., Reynolds, C.K., 2015. Methane emissions from cattle: estimates from short-term measurements using a GreenFeed system compared with measurements obtained using respiration chambers or sulphur hexafluoride tracer. *Animal Feed Science and Technology* 203, 41-52.

Hazel, L.N., 1943. The genetic basis for constructing selection indexes. *Genetics* 28, 476-490.

Herrero, M., Henderson, B., Havlík, P., Thornton, P.K., Conant, R.T., Smith, P., Wiersma, S., Hristov, A.N., Gerber, P., Gill, M., 2016. Greenhouse gas mitigation potentials in the livestock sector. *Nature Climate Change* 6, 452-461.

Jonker, A., Hickey, S., Pinares-Patiño, C., McEwan, J., Olinga, S., Díaz, A., Molano, G., MacLean, S., Sandoval, E., Harland, R., 2017. Sheep from low-methane-yield selection lines created on alfalfa pellets also have lower methane yield under pastoral farming conditions. *Journal of Animal Science* 95, 3905-3913.

Kandel, P., Vanderick, S., Vanrobays, M.-L., Soyeurt, H., Gengler, N., 2018. Consequences of genetic selection for environmental impact traits on economically important traits in dairy cows. *Animal Production Science* 58, 1779-1787.

Knapp, J.R., Laur, G.L., Vadas, P.A., Weiss, W.P., Tricarico, J.M., 2014. Invited review: Enteric methane in dairy cattle production: Quantifying the opportunities and impact of reducing emissions. *Journal of Dairy Science* 97, 3231-3261.

Lassen, J., Difford, G.F., 2020. Genetic and genomic selection as a methane mitigation strategy in dairy cattle. *Animal* 14, 473-483.

Lassen, J., Løvendahl, P., 2016. Heritability estimates for enteric methane emissions from Holstein cattle measured using noninvasive methods. *Journal of Dairy Science* 99, 1959-1967.

Lassen, J., Løvendahl, P., Madsen, J., 2012. Accuracy of noninvasive breath methane measurements using Fourier transform infrared methods on individual cows. *Journal of Dairy Science* 95, 890-898.

López-Paredes, J., Goiri, I., Atxaerandio, R., García-Rodríguez, A., Ugarte, E., Jiménez-Montero, J.A., Alenda, R., González-Recio, O., 2020. Mitigation of greenhouse gases in dairy cattle via genetic selection: 1. Genetic parameters of direct methane using noninvasive methods and proxies of methane. *Journal of Dairy Science* 103, 7199-7209.

Lynch, J., Cain, M., Pierrehumbert, R., Allen, M., 2020. Demonstrating GWP*: a means of reporting warming-equivalent emissions that captures the contrasting impacts of short- and long-lived climate pollutants. *Environmental Research Letters* 15, 044023.

Mrode, R.A., 2014. Linear models for the prediction of animal breeding values. Cabi Publishing, Wallingford, UK.

Niu, M., Kebreab, E., Hristov, A.N., Oh, J., Arndt, C., Bannink, A., Bayat, A.R., Brito, A.F., Boland, T., Casper, D., 2018. Prediction of enteric methane production, yield, and intensity in dairy cattle using an intercontinental database. *Global Change Biology* 24, 3368-3389.

Olijhoek, D.W., Difford, G.F., Lund, P., Løvendahl, P., 2020. Phenotypic modeling of residual feed intake using physical activity and methane production as energy sinks. *Journal of Dairy Science* 103, 6967-6981.

Pickering, N.K., Chagunda, M.G.G., Banos, G., Mrode, R., McEwan, J.C., Wall, E., 2015a. Genetic parameters for predicted methane production and laser methane detector measurements. *Journal of Animal Science* 93, 11-20.

Pickering, N.K., Oddy, V.H., Basarab, J., Cammack, K., Hayes, B., Hegarty, R.S., Lassen, J., McEwan, J.C., Miller, S., Pinares-Patiño, C.S., 2015b. Animal board invited review: genetic possibilities to reduce enteric methane emissions from ruminants. *Animal* 9, 1431-1440.

Pinares-Patiño, C.S., Hickey, S.M., Young, E.A., Dodds, K.G., MacLean, S., Molano, G., Sandoval, E., Kjestrup, H., Harland, R., Hunt, C., 2013. Heritability estimates of methane emissions from sheep. *Animal* 7, 316-321.

Rosenbloom, D., Markard, J., Geels, F.W., Fuenschilling, L., 2020. Opinion: Why carbon pricing is not sufficient to mitigate climate change—and how "sustainability transition policy" can help. *Proceedings of the National Academy of Sciences* 117, 8664-8668.

Sorg, D., Difford, G.F., Mühlbach, S., Kuhla, B., Swalve, H.H., Lassen, J., Strabel, T., Pszczola, M., 2018. Comparison of a laser methane detector with the GreenFeed and two breath analysers for on-farm measurements of methane emissions from dairy cows. *Computers and Electronics in Agriculture* 153, 285-294.

Tenghe, A.M.M., Bouwman, A.C., Berglund, B., de Koning, D.J., Veerkamp, R.F., 2018. Improving accuracy of bulls' predicted genomic breeding values for fertility using daughters' milk progesterone profiles. *Journal of Dairy Science* 101, 5177-5193.

Van der Maas, C.W.M., Brandes, L.J., Baas, K., Van den Born, G.J., Geilenkirchen, G., Te Molder, R., Nijdam, D.S., Olivier, J.G.J., Peek, C.J., Van Schijndel, M.W., 2009. Greenhouse Gas Emissions in the Netherlands 1990-2007. National Inventory Report 2009. Netherlands Environmental Assessment Agency, Bilthoven, the Netherlands.

- van der Werf, J. 2020. Teaching Software used for Quantitative Genetics at University of New England. Retrieved on 14 October 2020 from <https://jvanderw.une.edu.au/software.htm>.
- Van Eenennaam, A.L., Weigel, K.A., Young, A.E., Cleveland, M.A., Dekkers, J.C.M., 2014. Applied animal genomics: results from the field. *Annual Review Animal Bioscience* 2, 105–139.
- Van Engelen, S., Bovenhuis, H., van der Tol, P.P.J., Visker, M.H.P.W., 2018. Genetic background of methane emission by Dutch Holstein Friesian cows measured with infrared sensors in automatic milking systems. *Journal of Dairy Science* 101, 2226–2234.
- Van Middelaar, C.E., Berentsen, P.B.M., Dijkstra, J., Van Arendonk, J.A.M., De Boer, I.J. M., 2015. Effect of feed-related farm characteristics on relative values of genetic traits in dairy cows to reduce greenhouse gas emissions along the chain. *Journal of Dairy Science* 98, 4889–4903.
- Van Vleck, L.D., 1993. Selection index and introduction to mixed model methods for genetic improvement of animals: the green book. CRC Press, Boca Raton, FL, USA.
- Wall, E., Simm, G., Moran, D., 2010. Developing breeding schemes to assist mitigation of greenhouse gas emissions. *Animal* 4, 366–376.