

Calls and vocal behaviour of the Black Falcon *Falco subniger*

S. J. S. Debus¹, A. L. Bauer¹ and F. W. van Gessel²

¹Division of Zoology, University of New England, Armidale, New South Wales 2351

(Corresponding author: sdebus@une.edu.au)

²64 Dorothy Avenue, Woy Woy, New South Wales 2256 (Email:

fvangessel@optusnet.com.au)

Received: 10 May 2017

Sound-recordings of some calls of the Black Falcon *Falco subniger* (adult female cackle calls and begging whines, adult male creaking call) are described and presented as sound spectrograms, along with quantification of calling rates of both sexes at four active nests near Tamworth, NSW. Females were more vocal than males at all stages of the breeding cycle. Black Falcon cackle calls are similar to those of the Peregrine Falcon *F. peregrinus*, though more guttural, owing to a lower fundamental frequency of ~1 kHz in females, compared with ~2 kHz in female Peregrine Falcons. The Black Falcon is less vocal than the Peregrine, and breeding males use the whining call much less frequently than do Peregrine males.

INTRODUCTION

The vocalisations of the Black Falcon *Falco subniger* were described in general terms, with sound spectrograms, by Marchant and Higgins (1993) and Jurisevic (1998), although it was difficult to align and categorise these according to standard terminology of the cackle, creaking call and wailing call of large falcons

(e.g., as defined by Carlier 1995 and Leonardi *et al.* 2013). Subsequent studies on the Black Falcon tried to match observed call types with the aforementioned literature on that species (Debus *et al.* 2005; Debus and Olsen 2011; Debus and Tsang 2011; Debus and Zuccon 2013). However, it is apparent that some confusion remained, until Charley *et al.* (2014) and Whelan *et al.* (2016) provided somewhat more accurate interpretations. Baylis *et al.* (2015) noted that spectrographic metrics of the calls of Australian falcons require further study and comparison, of specific call types (e.g. cackle, creaking call, wail) in specific contexts across species, and that it remains to conduct a spectrographic study on the Black Falcon. Such would extend the prior (incomplete) studies on this species, and be potentially useful in systematics and in interpreting behavioural studies.

In 2015–16 we (SD and AB) studied the breeding biology and behaviour of four pairs of Black Falcons (Debus *et al.* 2017), partly modelling our approach (e.g., quantification of calling rates) on studies of the Lanner Falcon *F. biarmicus* (Leonardi *et al.* 2013) and Red-headed Falcon *F. chicquera* (Naoroji 2011). In the process, one of us (FVG) obtained sound-recordings of some typical calls. A comparable study has been conducted on the Grey Falcon *F. hypoleucos* (Ley and Tynan 2016), with spectrograms of its cackle and whining calls provided by Baylis *et al.* (2015). The Lanner and Black Falcons are heirofalcons ('great' or 'desert' falcons), the Grey Falcon is sister to the heirofalcon/Peregrine Falcon *F. peregrinus* split, and the Red-headed Falcon is sister to that divergence (Fuchs *et al.* 2015), thus all are related to some degree. As falcon vocalisations are evidently innate (Cade 1982; Jurisevic 1998), and there is no suggestion of individual or regional variation in the calls of the

Black Falcon (Marchant and Higgins 1993), the results presented here may be applicable continent-wide, although further study would be fruitful.

METHODS

The study area was the Tamworth district of northern inland New South Wales, and the subject pairs are here numbered 1 to 4 corresponding with the biological study (see Debus *et al.* 2017). Adult vocalisation types were categorised according to previous studies on hierofalcons (including the Black Falcon) and the Peregrine, i.e. the cackle, creaking call (or *ee-chip*) and food-begging wail or whine ('whine' being considered more apt than 'wail'; e.g. Marchant and Higgins 1993; Leonardi *et al.* 2013; Charley *et al.* 2014; Baylis *et al.* 2015; Ley and Tynan 2016). Vocalisations were quantified as the number of separate calling events (bouts) rather than individual syllables (which were highly variable within a calling bout), a 'bout' being a sequence that was separated from others by a distinct pause at least the duration of a syllable of that call type.

The principles for interpreting sound spectrograms, as described elsewhere (Baylis *et al.* 2015), apply here. That is, spectrograms are graphical representations of the structure, pitch and tempo of bird calls, the vertical axis showing frequency (pitch) and the horizontal axis showing time elapsed. The fundamental frequency is the lowest value (trace) in the spectrogram, and the dominant frequency is the frequency band that contains the greatest acoustic energy.

Recordings of Black Falcon calls were made at two active nests by FVG on 31 July–1 August (Pair 2) and 23–24 September 2016 (Pair 4), using three open Sennheiser K6 modular ME66 microphones and parabolic dish (50-cm diameter fitted with an ME64 microphone) connected to a digital Sound Devices 702 recorder and two Tascam DR44WL recorders at each site (set up ~130 m from each nest). The recordings were made using a sampling rate of 96 kHz at 24 bits as WAV files (Waveform Audio File Format, being uncompressed audio files). To generate the spectrograms, the recordings were edited and down-sampled to 44.1 kHz at 16 bits (i.e. CD quality) for use with Raven Lite 1.0. The fast Fourier transform (FFT) was set at 512, otherwise settings were Raven Lite 1.0 default settings (e.g. window type is Hann, setting unstated for dB depth). The recordings will be made available to *bona fide* researchers, but not publicly (e.g. through Sound Cloud), to reduce the risk of over-use and misuse of call-playback.

In processing the sound-recordings for presentation of spectrograms herein (Figures 1–3) at the same scale (standardised to 3.5 sec. in length), all spectrograms were made using Raven Pro 1.4, FFT 1024, Hann window, overlap 50%, then converted to greyscale in Photoshop for publication. No low-cut filter was used on the sound-recordings, but for producing the spectrograms a filter was used to delete the very loud and low frequencies (60–70 Hz) made by aircraft noise and wind (i.e. environmental noise). The Black Falcon spectrograms are from typical samples of multiple recordings of multiple individuals (see above), which we regard as representative of the respective call types of that species (based on SD's experience of Black Falcon calls interstate, e.g. Queensland and Victoria, and of a further two

breeding pairs not sampled by FVG in this study). Therefore, the recorded calls are likely to be within the range of individual variation.

RESULTS

Vocalisation types

Breeding adult Black Falcons of both sexes uttered the three main call types (Table 1), examples being the male's creaking call (Figure 1), the female's whining call (Figure 2) and the female's cackling call (Figure 3). All are broad-band calls with most acoustic energy in the 2–3.5 kilohertz range. Creaking calls by the male were usually an abrupt 'chip' or 'chik' sound repeated, sometimes disyllabic (fundamental frequency ~800 Hz, dominant frequencies ~1.7–3.8 kHz; Figure 1); those of the female were deeper and more guttural, occasionally disyllabic ('ee-chip' type). Female whining calls were long, slightly rising notes (~0.5–1 sec duration and pulse rate, fundamental frequency ~ 1.9 kHz, dominant frequencies 2–3.5 kHz; Figure 2). The whining call of one female during the incubation period (Figure 2A) was at times more rapid and intense than that of another during the nestling period (Figure 2B). Either sex occasionally uttered a single low moan, deeper in pitch than the whine, when approaching the nest tree. Female cackle calls were a series of slow, hoarse cackles with a guttural quality, the notes somewhat clipped, uttered at a slightly faster rate than the whining call (cackle fundamental frequency ~ 1 kHz, dominant frequencies ~1.9–2.7 kHz; Figure 3), and more guttural than the male's cackle. Male cackle calls sounded somewhat more Peregrine-like than the female's. A male fledgling, when captured, uttered a shrill, 'screaming' version of the cackle (whereas

a captured female fledgling uttered a repeated, deep resonant whining call: Debus *et al.* 2005).

Vocal behaviour

Female Black Falcons were more vocal than males at all stages of the breeding cycle, and the frequency of the three main call types also changed through the cycle (Table 1). Creaking calls, used by the male as nest-advertisement (to the female in the courtship period) and food-delivery calls, by the female in social contexts (e.g. nest selection and defence, pre-copulatory posture) and by both sexes in intraspecific conflicts, were most frequent in the pre-laying (especially) and incubation periods. Whining calls were frequently used by females as food-begging calls throughout the cycle, but particularly in the incubation period. Cackle calls became frequent in the incubation and especially the nestling period, during interspecific nest defence (notably against Wedge-tailed Eagles *Aquila audax*) and especially by the female (the main defender of the nest and young: Debus *et al.* 2017) (Table 1). Creaking calls, particularly during intraspecific conflicts and especially by the female, often became long rattling or chattering series (e.g. up to at least 20 syllables). Similarly, the female's whining calls when begging to the male, especially if he was present but had no prey, often became prolonged, insistent series (e.g. Figure 2A). Thus, the frequency of bouts of these call types by the female during the pre-laying to nestling periods does not fully reflect their intensity. For instance, when begging intently the female's whining could persist for up to four bouts per minute for up to 20 syllables per bout, for up to five minutes in duration before pausing.

DISCUSSION

This study clarifies that the Black Falcon's three main loud calls are the cackle, the creaking call and the whining call, often heard around the nest or (the cackle) when defending the area against eagles. The long series of creaking calls run together are what have been called chattering, rattling or chuckling calls elsewhere (e.g. Debus *et al.* 2005; Charley *et al.* 2014; Whelan *et al.* 2016). The 'greeting trill' of Charley *et al.* (2014) was a soft, musical version of the extended creaking series, and the calls of interacting juveniles in Barnes and Debus (2012) were the cackle. The Black Falcon's calls, including softer 'ticking' or clucking variants of the creaking call at the nest, are similar to equivalent calls of the Lanner Falcon (see Leonardi 2015). As for the Lanner Falcon and Grey Falcon (Leonardi *et al.* 2013; Ley and Tynan 2016), females are more vocal, and use the whining call much more frequently, than males (Table 1).

The results of this study enable classification of the various descriptions and/or spectrograms in Marchant and Higgins (1993) and Jurisevic (1998) according to standard terminology for falcons. The adult Black Falcon call types as numbered by Marchant and Higgins (pp. 303–304) are: (1) first sentence = cackle, second sentence = creaking call; (2) = low moan; (3) = begging whine; (4) first sentence = maternal cluck, second sentence = single cackle note; (5) = male creaking call in aerial display ('falsetto' version, labelled Sonagram A); (7) first sentence = begging whine, final clause = creaking call; Sonagrams B and C are juvenile creaking call (B) and whine (C). Similarly, the Black Falcon spectrograms in Figure 2 of Jurisevic (1998) are: (a, b) male and female whining calls; (c, d) cackle calls; and (e) a distress chitter. The

adult Peregrine Falcon spectrograms in Figure 1 of Jurisevic (1998) appear to be: (a, b) cackle calls; (c, d) whining calls; (e) creaking calls; and (f) a single cackle note. It is apparent from this study and that of Jurisevic (1998) and Baylis *et al.* (2015) that the Black Falcon's creaking and cackle calls are somewhat deeper (slightly lower in pitch, i.e. in fundamental and dominant frequencies) than those of the Peregrine Falcon (~1 kHz vs ~2 kHz, respectively), and the cackle call is also uttered rather more slowly than typical Peregrine cackles.

The relative frequency of use of the creaking and whining calls by the Black Falcon sexes, and their change in relative frequency of use through the breeding cycle, are fairly consistent with the pattern in other well-known hierofalcons (cf. Potapov and Sale 2005; Leonardi *et al.* 2013). These aspects contrast with the more vocal Peregrine Falcon, in which there is no difference between the sexes in the use of the creaking call in the pre-laying period, or in the whining call throughout the cycle (Carlier 1995). However, only one of three Black Falcon nests at Tamworth succeeded, fledging a single underweight male (Debus *et al.* 2017), so vocal rates might differ at more successful nests under better weather and prey conditions (as suggested by Charley *et al.* 2014). These comparisons are still rather incomplete; it remains to obtain sound-recordings of the male Black Falcon's cackling call, the female's creaking call, and the long series of creaking calls by either sex, and subject them to similar analysis as here.

ACKNOWLEDGEMENTS

We here gratefully acknowledge the assistance provided to the main biological study on Black Falcons at Tamworth, notably support of Alice Bauer's UNE Zoology Honours project (see Debus *et al.* 2017). Thanks to Paul M^cDonald, Giovanni Leonardi and Alan Lill for comments on a draft. This study was conducted under UNE Animal Ethics authority AEC16-02.

REFERENCES

- Barnes, C. P. and Debus, S. J. S. (2012). A snapshot in the post-fledging period of the Black Falcon. *Australian Field Ornithology* **29**: 86–88.
- Baylis, T., van Gessel, F. W. and Debus, S. J. S. (2015). Some vocalisations of the Grey Falcon *Falco hypoleucos*. *Corella* **39**: 73–76.
- Cade, T. J. (1982). 'The Falcons of the World'. (Collins: London.)
- Carlier, P. (1995). Vocal communication in Peregrine Falcons *Falco peregrinus* during breeding. *Ibis* **137**: 582–585.
- Charley, D., Lutter, H. and Debus, S. J. S. (2014). Breeding behaviour and prey of Black Falcons, *Falco subniger*, including food-caching. *South Australian Ornithologist* **40**: 11–30.
- Debus, S. J. S. and Olsen, J. (2011). Some aspects of the biology of the Black Falcon *Falco subniger*. *Corella* **35**: 29–36.
- Debus, S. J. S. and Tsang, L. R. (2011). Notes on Black Falcons *Falco subniger* breeding near Tamworth, New South Wales. *Australian Field Ornithology* **28**: 13–26.
- Debus, S. J. S. and Zuccon, A. E. (2013). Observations on hunting and breeding behaviour of the Black Falcon *Falco subniger*. *Sunbird* **43**: 12–26.
- Debus, S. J. S., Bauer, A. L. and Mitchell, G. I. (2017). Breeding biology, behaviour and foraging ecology of the Black Falcon *Falco subniger* near Tamworth, New South Wales. *Corella* **41**: ?-?.
- Debus, S. J. S., Hatfield, T. S., Olde, G. S. and Rose, A. B. (2005). Breeding behaviour and diet of a pair of Black Falcons *Falco subniger* in northern New South Wales. *Australian Field Ornithology* **22**: 165–181.

- Fuchs, J., Johnson, J.A. and Mindell, D.A. (2015). Rapid diversification of falcons (Aves: Falconidae) due to expansion of open habitats in the late Miocene. *Molecular Phylogenetics and Evolution* **82**: 166–182.
- Jurisevic, M. A. (1998). Comparison of vocalisations of Australian falcons and elanine kites. *Emu* **98**: 1–12.
- Leonardi, G., Amato, M., Brogna, A., Cipriano, M., D'Angelo, R., Dipasquale, G., Mannino, V., Ossino, A. and Andreotti, A. (2013). Effectiveness of vocal activities of the Lanner Falcon (*Falco biarmicus*) during breeding. *Journal of Ornithology* **154**: 83–89.
- Ley, A. and Tynan, B. (2016). Observations on nesting Grey Falcons, *Falco hypoleucos*. *South Australian Ornithologist* **41**: 49–64.
- Marchant, S. and Higgins, P. J. (Eds) (1993). 'Handbook of Australian, New Zealand and Antarctic Birds, Volume 2: Raptors to Lapwings'. (Oxford University Press: Melbourne.)
- Naoroji, R. (2011). Breeding of the Red-headed Falcon *Falco chicquera* in Saurashtra, Gujarat, India. *Forktail* **27**: 1–6.
- Potapov, E. and Sale, R. (2005). 'The Gyrfalcon'. (Yale University Press: New Haven, CT, USA.)

Table 1

Frequency of adult Black Falcon vocalisation types (calling bouts per hour of observation time) in stages of the breeding cycle: pre-laying period to laying/start of incubation (Pair 1, May–July 2015); incubation period (Pairs 2 and 3 pooled, July–September 2016); nestling period (Pairs 3 and 4 pooled, September–October 2016).

	Creaking		Whining		Cackle	
	Male	Female	Male	Female	Male	Female
Pre-lay (94.5 h)	0.15	0.97	0.03	1.47	0	0.07
Incubation (134.75 h)	0.11	0.29	0.01	2.69	0.01	0.24
Nestling (50.83 h)	0.04	0	0	0.94	0.06	0.37

Figure 1. *Spectrogram of creaking calls of adult male Black Falcon (a) at nest, and (b) as approaching nest with prey, Pair 2 (female incubating). Tamworth (NSW), July–August 2016; spectrograms prepared using Raven Lite 1.0.*

Figure 2. *Spectrogram of whining calls of adult female Black Falcon as male approaching nest with prey: (a) incubation period, Pair 2, and (b) nestling period, Pair 4. Tamworth (NSW), August–September 2016; spectrograms prepared using Raven Lite 1.0.*

Figure 3. *Spectrograms of two examples of cackling call of adult female Black Falcon defending nestling against Wedge-tailed Eagle, Pair 4. Tamworth (NSW), September 2016; spectrograms prepared using Raven Lite 1.0.*